

THE ALDERNEY PLANT LIST

This list last updated 10.01.2008

THE PLANT LIST

No.	Scientific name	English name	Finder/date
Note; Individual squares are not generally shown for abundant, common, or frequent plants			
* = 1999 list of Endangered or Scarce species			
# = rare species in Alderney (not usually endangered) Often frequ			

PTERIDOPHYTA

Lycopodiopsida

Selaginellaceae

1	Selaginella kraussiana	#	Kraus's clubmoss	JP 1993
---	------------------------	---	------------------	---------

Isoetaceae

2	Isoetes histrix	# *	Land Quillwort (RDBk)	EDM 1902
---	-----------------	-----	-----------------------	----------

Equisetopsida

Equisetaceae

3	Equisetum arvense		Field Horsetail	EDM 1901
4	Equisetum palustre		Marsh Horsetail	CCB 1838
5	Equisetum telmateia	#	Great Horsetail	CRPA 1900

Ophioglossaceae

6	Ophioglossum vulgatum	#	Adder's-tongue	JES 1969
7	Ophioglossum lusitanicum	# *	Least Adder's-tongue (RDBk)	EDM 1906

Pteropsida

Osmundaceae

8	Osmunda regalis	#	Royal Fern	WMT 1967
---	-----------------	---	------------	----------

Adiantaceae

9	Adiantum capillus-veneris	# *	Maidenhair Fern (Scarce)	WTA 1865
---	---------------------------	-----	--------------------------	----------

Polypodiaceae

10	Polypodium vulgare	#	Polypody	CCB 1838
11	Polypodium interjectum		Intermediate Polypody	RMW 1962

Dennstaedtiaceae

(Formerly hypolepidaceae)

12	Pteridium aquilinum		Bracken	CCB 1838
----	---------------------	--	---------	----------

Aspleniaceae

13	Phyllitis scolopendrium		Hart's-tongue Fern	CCB 1838
14	Asplenium adiantum-nigrum		Black Spleenwort	CCB 1838

THE PLANT LIST

Frequency	Fertile period	Squares where found and notes
-----------	----------------	-------------------------------

nt or com mon in Britain

1 site		8, Large patch 10 x 1.5m along base of church wall NW (NW corner)
--------	--	---

vr	1 to 6	5,10, 14 2 new sites on E coast found 6/1995, JB, MK
----	--------	--

lc	3 to 7	Generally distributed in damp places. Separate
----	--------	--

lf	4 to 8	4,7,8,9,10,13. Cones on vegetative shoots.
----	--------	--

2 sites	3 to 5	12, Veg stem to 1.5m. Fruiting stems 25cm, brown, (appear first)
---------	--------	--

1 site	5 to 8	5
--------	--------	---

Only record, probably an error

1 site	5 to 9	12. 1 large plant established about 40+ years
--------	--------	---

Only other record CH

vr	5 to 11	All CI records doubtful, now referred to next species.
----	---------	--

c	6 to 12	Frequent on old mortared walls and in most squares
---	---------	--

a	7 to 10	
---	---------	--

a	1 to 12	
---	---------	--

c	6 to 12	Petiole black
---	---------	---------------

THE PLANT LIST

No.	Scientific name		English name	Finder/date
15	<i>Asplenium obovatum</i>	# *	Lanceolate Spleenwort (Scarce)	CCB 1838
16	<i>Asplenium marinum</i>		Sea Spleenwort	CCB 1838
17	<i>Asplenium trichomanes</i>		Maidenhair Spleenwort	EDM 1901
18	<i>Asplenium ruta-muraria</i>		Wall-rue	EDM 1901
19	<i>Ceterach officinarum</i>	#	Rusty-back Fern	EDM 1901
Woodsiaceæ			Formerly Athyriaceæ	
20	<i>Athyrium filix-femina</i>		Lady-fern	CCB 1838
Dryopteridaceæ			Formerly Aspidiaceæ	
21	<i>Polystichum setiferum</i>		Soft Shield-fern	FLS 1960
22	<i>Phanerophlebia falcatum</i>	#	House Holly-fern	PL 1976
23	<i>Dryopteris filix-mas</i>		Male-fern	CCB 1838
24	<i>Dryopteris affinis</i> ssp. <i>borreri</i>		Golden-scaled Male-fern	DMcC 1957
25	<i>Dryopteris dilatata</i>		Broad Buckler-fern	EDM 1901
Blechnaceæ				
26	<i>Blechnum spicant</i>	# *	Hard-fern	EDM 1901
Azollaceæ				
27	<i>Azolla filiculoides</i>	#	Water Fern	JES 1972

THE PLANT LIST

Frequency	Fertile period	Squares where found and notes
r	6 to 10	Cliffs, esp W. end of slides in Bluestone Bay
lf	3 to 10	4,5. Also in offshore island Forts
o	6 to 9	4,7,8. Mainly on old mortared walls
o	1 to 12	4,5,7,8,10,12. Especially old walls in town
vr	1 to 12	4,7,8. A few small colonies on old walls
o	2 to 12	Scattered
r	7 to 10	7,8, 9 In G.C.D. 1932
1 site	11 to 1	7. Several specimens stream bank near Watermill,
f	1 to 12	(spread by 2002)
lf	1 to 12	3,4,5,7,8,9,10,11,13
r	5 to 10	7,8,12. (Also in quarry with the Osmunda)
vr	6 to 8	6,7,12. Last record 1937
4 sites	6 to 9	1,4,5,7 in Corblets Quarry & Platte Saline ponds. Found with spores 22.8.88



Azolla in spore

THE PLANT LIST

No.	Scientific name	English name	Finder/date
-----	-----------------	--------------	-------------

Gymnospermæ

Mostly planted but some self seed

Pinopsida

Pinaceæ

28	<i>Picea abies</i>	# Norway Spruce	BB 1987
29	<i>Larix decidua</i>	# European Larch	CH 1987
30	<i>Cedrus deodora</i>	# Deodar	CH 1987
31	<i>Cedrus libani</i>	# Cedar-of-Lebanon	BB1987
32	<i>Pinus sylvestris</i>	# * Scots Pine (Scarce)	CH 1987
33	<i>Pinus nigra</i> ssp. <i>nigra</i>	Austrian Pine	CH 1987
34	<i>Pinus nigra</i> ssp. <i>laricio</i>	Corsican Pine	BB 1986
35	<i>Pinus pinaster</i>	Maritime Pine	BB 1987
36	<i>Pinus radiata</i>	Monterey Pine	DMcC 1975
37	<i>Pinus attenuata</i>	# Knobcone Pine	BB 1987
38	<i>Tsuga heterophylla</i>	# Western Hemlock	RW 1987

Cupressaceæ

39	<i>Cupressus macrocarpa</i>	Monterey Cypress	BB 1986
40	X <i>Cupressocyparis leylandii</i>	Leyland Cypress	BB 1987
41	<i>Chamæcyparis lawsoniana</i>	Lawson's Cypress	BB 1987
42	<i>Thuja occidentalis</i>	# White Cedar	CH 1987
43	<i>Thuja plicata</i>	# Western Red-cedar	BB 1985

Araucariaceæ

44	<i>Araucaria araucana</i>	# Monkey-puzzle	BB 1986
----	---------------------------	-----------------	---------

Taxaceæ

45	<i>Taxus baccata</i>	# Yew	BB 1986
----	----------------------	-------	---------

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
-----------	------------------	-------------------------------

Mostly planted

vr	5 to 6	
r	3 to 6	8. 3 specimens to 13m. others smaller
vr	12 to 1	9
vr	10 to 12	
r	3 to 7	3,5,7,8,13
f	4 to 7	3,8
f	4 to 6	3,8. Specimen to 16.5.m
o	4 to 7	3,7,8,10,13. Plantation Newtown Road.
c	2 to 7	Specimens to 14.5m. Most frequent planting
1 site	5 to 6	8. One stand in Newtown Road
1 site	4 to 5	8. Island Hall garden
f	5 to 6	3,7,8,9. Trees to 20.5m badly damaged in hurricane 10/87
o	5 to 6	Much used in hedging
f	3 to 4	3,7,9,13. Mostly in gardens
vr		8
1 site	3	8. Island Hall garden. Tree cut down 2006
o	6	8. Several in Gardens
r	2	8. Only in gardens and Parish churchyard

THE PLANT LIST

No.	Scientific name	English name	Finder/date
 Magnoliopsidæ			
1. Magnoliidæ			
Magnoliopsida			
 Lauraceæ			
46	<i>Laurus nobilis</i>	Bay	BB 1987
 Nymphaeaceæ			
47	<i>Nymphaea alba</i>	White Water-lily	BB 1986
48	<i>Nuphar lutea</i>	# * Yellow Water-lily	BB 1989
 Ceratophyllaceæ			
49	<i>Ceratophyllum submersum</i>	# Soft Hornwort	BB1987
 Ranunculaceæ			
50	<i>Caltha palustris</i>	# * Marsh Marigold	AJR 1966
51	<i>Consolida ajacis</i>	# * Larkspur	EDM 1901
52	<i>Clematis vitalba</i>	Traveller's-joy	CMR 1950
53	<i>Ranunculus acris</i>	Meadow Buttercup	CCB 1838
54	<i>Ranunculus repens</i>	Creeping Buttercup	EDM 1900
55	<i>Ranunculus bulbosus</i>	Bulbous Buttercup	CCB 1838
56	<i>Ranunculus sardous</i>	# Hairy Buttercup	BB 1987
57	<i>Ranunculus parviflorus</i>	Small-flowered	EDM 1901
58	<i>Ranunculus lingua</i>	# Greater Spearwort	CCB 1838
59	<i>Ranunculus flammula</i>	Lesser Spearwort	EDM 1901
60	<i>Ranunculus ficaria</i> agg.	Lesser Celandine	EDM 1901
61	<i>Ranunculus ficaria</i> ssp. <i>ficaria</i>	Lesser Celandine	BB 1996
62	<i>Ranunculus ficaria</i> ssp.	Lesser Celandine	BB 1996
63	<i>Ranunculus ficaria</i> ssp. <i>bulbifer</i>	Lesser Celandine	BB 1996
64	<i>Ranunculus ficaria</i> ssp.	Lesser Celandine	BB 1996
65	<i>Ranunculus hederaceus</i>	# Ivy-leaved Crowfoot	CCB 1838
66	<i>Ranunculus trichophyllus</i>	Three-leaved Water-cro~	EDM 1901
67	<i>Thalictrum minus</i>	# Lesser Meadow-rue	MM 1985

THE PLANT LIST

Frequency Flowering period Squares where found and notes

(Formerly Dicotyledonidæ)

r	3 to 6	4,8, Large specimens, Vicarage, St. Vignalis, Berry's Quarry
2 sites	5 to 9 6 to 7	4,10 10. Not refound, ? error for <i>Nymphoides peltata</i> .
1 site	7 to 9	4. Lvs. light green, segments scarcely toothed
vr	4 to 7	7. Now seems to be confined to domestic ponds
vr	6 to 7	8. Occasional garden escape. Only other record 1957
vlf	5 to 10	2,3,8,9, 12 Newtown, Rochers, Braye Dunes
c	12 to 9	
c	1 to 12	
c	12 to 8	
vr	3 to 4	7
o	4 to 6	4,5,7,9, 10,14
vr	6 to 7	4,7,12,13 Refound BB 1991. Previous GCD
o	4 to 8, 11	7,11,13. In damp valleys
la	1 to 6	10. Small fls., 25mm, long stems, leaves spreading, 7. Small fls. short stems, congested leaves, no bulbils 7. Small fls, short stems, with bulbils 7. Large fls 35mm. No bulbils
vr	6 to 9	6,7, Vau Pommier. Still there AB/AKJ 1934. Only records
vlc	5 to 6	4, 5. Mannez & Longis ponds. Berry's Quarry pool
vr	6 to 8	3,5

THE PLANT LIST

No.	Scientific name	English name	Finder/date
Berberidaceæ			
68	<i>Berberis vulgaris</i>	# Barberry	BB 1988
69	<i>Berberis darwinii</i>	# Darwin's Barberry	BB1989
Papaveraceæ			
70	<i>Roemeria hybrida</i>	# Violet Horned Poppy	VKM 1956
71	<i>Papaver somniferum</i>	Opium Poppy	EDM 1901
72	<i>Papaver rhoeas</i>	Common Poppy	CCB 1838
73	<i>Papaver dubium</i>	Long-headed Poppy	EDM 1901
74	<i>Papaver hybridum</i>	Rough Poppy	CCB 1838
75	<i>Glaucium flavum</i>	Yellow Horned-poppy	CCB 1838
76	<i>Glaucium corniculatum</i>	# Red Horned-poppy	TB 1903
77	<i>Eschscholzia californica</i>	Californian Poppy	DMcC 1953
Fumariaceæ			
78	<i>Pseudofumaria lutea</i>	# Yellow Corydalis	BB 1990
79	<i>Fumaria capreolata</i>	White Ramping-fumitory	CCB 1838
80	<i>Fumaria bastardii</i>	# Tall Ramping-fumitory	EDM 1901
81	<i>Fumaria muralis</i> ssp. <i>boraiei</i>	Common	EDM 1901
82	<i>Fumaria officinalis</i>	Common Fumitory	CCB 1838
Most of today's Elms were originally planted, a few have survived the Dutch Elm disease as suckers			
Ulmaceæ			
83	<i>Ulmus glabra</i> ssp. <i>glabra</i>	# Wych Elm	CCB 1838
84	<i>Ulmus glabra</i> ssp. <i>montana</i>	# Smooth-leaved Elm	CCB 1838
85	<i>Ulmus x hollandica</i>	# Dutch Elm	CCB 1838
86	<i>Ulmus procera</i>	# English Elm	CCB 1838
87	<i>Ulmus minor</i> ssp. <i>minor</i>	# Small-leaved Elm	EDM 1901
88	<i>Ulmus minor</i> ssp. <i>angustifolia</i>	# Cornish Elm	1957
89	<i>Ulmus minor</i> ssp. <i>sarniensis</i>	# Jersey Elm	?
Cannabaceæ			
90	<i>Cannabis sativa</i>	# Hemp	EDM 1902
91	<i>Humulus lupulus</i>	# Hop	EDM 1901

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
vr	1 to 8	10, Longis Bay Determined by DMcC 6/88
vr	10 to 6	4, Corblets Bay, either side of road
		Only record
		10. Specimen confirmed by FLS 1967. S.T.P. Nr. Fort Essex
o	5 to 11	2,5,7,8
f	5 to 12	1,2,3,7,9
lf	6 to 9	6,10,11,12
o	5 to 8	7,8,10,14 Near Fort Albert FLS 1967 S.T.P.
lc	5 to 11	1,2,3,7,12
?	?	7 1 plant at Platte Saline. Only record
o	5 to 11	2,5,7. Occasional garden escape
vr	5 to 8	7,8 Gutter in town, wall at Platte Saline. Garden
r	4 to 8	2 (1967), 7 (1989), 14 (1838), cliff paths (1995)
vr	7 to 8	7. Refound BB 1989. previous record ABJ 1938
c	1 to 12	Most Alderney Fumitory plants are this sp.
o	5 to 10	9,12,13. On cultivated ground. On Les Rochers F.L.S. 1967 S.T.P.
vr	2 to 3	7,8,10,12
vr	3 to 5	8. ABJ 1932 as <i>U. nitens</i> 'A row in Reuter's Valley'
r	2 to 3	7,8. All the remaining elms need critical
r	2 to 3	Probably Babington's <i>U. suberosa</i> & Marquand's <i>U.</i>
r	3 to 5	8
r	3 to 5	Determined by RM 1957 as <i>U. cornubiensis</i> var.
f	3 to 5	7,8. Suckers only now
?	?	7,8, Only other record, a single plant 1987, destroyed
vr	7 to 8	8. Refound RW 1987. Also recorded BSBI 1995

THE PLANT LIST

No.	Scientific name	English name	Finder/date
Moraceæ			
		Probably all planted	
92	<i>Ficus carica</i>	# Fig	BB 1987
Urticaceæ			
93	<i>Urtica dioica</i>	Common Nettle	CCB 1838
94	<i>Urtica urens</i>	Small Nettle	CCB 1838
95	<i>Parietaria judaica</i>	Pellitory-of-the-Wall	EDM 1901
96	<i>Soleirolia soleirolii</i>	Mind-your-own-business	FLS 1956
Juglandaceæ			
		Planted	
97	<i>Juglans regia</i>	# Walnut	NVM 1966
Fagaceæ			
		Mostly planted	
98	<i>Fagus sylvatica</i>	Beech	DMcC 1957
99	<i>Castanea sativa</i>	# Sweet Chestnut	EDM 1901
100	<i>Quercus cerris</i>	# Turkey Oak	DMcC 1957
101	<i>Quercus ilex</i>	Evergreen Oak (native)	CH 1987
102	<i>Quercus petræa</i>	# Sessile Oak	BB 1987
103	<i>Quercus robur</i>	# Pedunculate Oak	CCB 1838
Betulaceæ			
		Mostly planted	
104	<i>Betula pendula</i>	Silver Birch	EDM 1901
105	<i>Alnus glutinosa</i>	Alder	DMcC 1975
106	<i>Carpinus betulus</i>	# Hornbeam	RW 1986
107	<i>Corylus avellana</i>	Hazel (probably native)	RW 1987
Nyctaginaceæ			
108	<i>Mirabilis jalapa</i>	# Marvel-of-Peru	BB 1987
Aizoaceæ			
109	<i>Aptenia cordifolia</i>	Heart-leaf Ice-plant	BB 1987
110	<i>Lampranthus roseus</i>	Rosy Dew-plant	MR 1978
111	<i>Carpobrotus acinaciformis</i>	Sally-my-Handsome	DMcC 1959
112	<i>Carpobrotus edulis</i>	Hottentot-fig	JO 1953
113	<i>Carpobrotus glaucescens</i>	Angular Sea-fig	BB 1988
113a	<i>Tetragonia tetragonioides</i>	# New Zealand Spinach	JP2006

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
r	5 to 6	4,7,8. Berry's Quarry. Path to Clonque. Scattered in gardens
a	4 to 10	
f	5 to 9	
c	1 to 12	
lf	6 to 8	6,7,8,10. Moist places & old gardens. Wall bottoms
1 site	5	8 Single specimen in Island Hall garden
o	4 to 5	7,8 Including copper beeches
o	5 to 7	7,8,10. Scattered
o	4 to 7	8 plus scattered small specimens
f	4 to 7	Frequent planting, self-seeds
vr	4 to 7	7,8,9
r	4 to 7	2,7,8,9,10,13
o	3 to 6	7,8,9,10,13
o	1 to 5	6, 7,8,9,12,13 Watered valleys, self-seeds
vr	4 to 5	8, 2 in Churchyard
vr	1 to 4	4,10.
vr	7 to 10	Occasional persisting garden weed or escape
vr	3 to 12	2,8, On old walls
o	1 to 12	6,10,13. (S. Cliffs, first noted July 1978)
o	3 to 7	4,5,7,12,13. Fls. 8-10cm. pink/ purple. Lvs curved, thickest near top
f	1 to 12	Old forts and cliffs. Fls 4-9cm pink/purple or yellow. Lvs equal, thick
lf	3 to 10	2,4,5,6,7. Fls. 3.5-6cm, pink/purple, petals white at base
vr	?-11	9. Barrackmaster's Lane nr. Devereux. 1 plant 1st record 20.11.06

THE PLANT LIST

No.	Scientific name	English name	Finder/date
Chenopodiaceæ			
114	<i>Chenopodium polyspermum</i>	Many-seeded Goosefoot	FS/CR 1958
115	<i>Chenopodium vulvaria</i> # *	Stinking Goosefoot (RDB)	?
116	<i>Chenopodium murale</i>	Nettle-leaved Goosefoot	CCB 1838
117	<i>Chenopodium ficifolium</i>	Fig-leaved Goosefoot	EDM 1901
118	<i>Chenopodium opulifolium</i> #	Grey Goosefoot	GCD 1906
119	<i>Chenopodium album</i>	Fat-hen	CCB 1838
120	<i>Atriplex prostrata</i>	Spear-leaved Orache	EDM 1901
121	<i>Atriplex prostrata</i> x <i>A. glabriuscula</i> #		EMJ 1967
122	<i>Atriplex glabriuscula</i>	Babington's Orache	EDM 1901
123	<i>Atriplex littoralis</i> #	Grass-leaved Orache	CCB 1838
124	<i>Atriplex patula</i>	Common Orache	CCB 1838
125	<i>Atriplex laciniata</i>	Frosted Orache	EDM 1901
126	<i>Atriplex portulacoides</i> #	Sea-purslane	WC 1838
127	<i>Beta vulgaris</i> ssp. <i>maritima</i>	Sea-beet	CCB 1838
128	<i>Salsola kali</i>	Prickly Saltwort	CCB 1838

Amaranthaceæ

129	<i>Amaranthus retroflexus</i>	Common Pigweed	DMcC 1957
-----	-------------------------------	----------------	-----------

Portulacaceæ

130	<i>Claytonia sibirica</i> #	Pink Purslane	BB 1987
130a	<i>Portulaca oleracea</i> #	Common Purslane	ML 2001
131	<i>Montia fontana</i> ssp. <i>fontana</i>	Blinks	EDM 1901
131a	<i>Montia fontana</i> ssp. <i>variabilis</i>		EDM 1901
132	<i>Montia fontana</i> ssp. <i>chondrosperma</i> #		AB/AKJ 1934
133	<i>Montia fontana</i> ssp. <i>amporitana</i> #		1962

Caryophyllaceæ

134	<i>Arenaria serpyllifolia</i> ssp. <i>serpyllifolia</i>	Thyme-leaved Sandwort	CCB 1838
135	<i>Arenaria serpyllifolia</i> ssp. <i>leptoclados</i>	Lesser Thyme-leaved Sandwort	EDM 1901
135a	<i>Moerhingia trinervia</i> #	Three-leaved Sandwort	BB 1996
136	<i>Honckenya peploides</i>	Sea Sandwort	EDM 1901
137	<i>Stellaria media</i>	Common Chickweed	CCB 1838
138	<i>Stellaria pallida</i>	Lesser Chickweed	ABJ 1934

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
vr	6 to 9	7. Refound BB 1995 Platte
	7 to 9	Marked as ? errors in WFG Recorded in Atlas 1962
f	5 to 10	Waste places
r	6 to 9	7,8,14
		No site. Only record
c	6 to 10	Gardens & cultivated ground
f	7 to 9	1,2,3,5,8,11. Most shores
?		'On beaches'. Only
o	6 to 10	1,2,3,4,6,7. Sandy shores
	7 to 8	Possibly the only record
f	6 to 10	
f	7 to 10	1,2,3,4,5. Beaches, just about HWM
r	6 to 7	5,6,7,11,12,13. Rarely at beach level here, mostly on cliff faces
c	5 to 9	
o	6 to 9	2,3,4,5. At top of sandy beaches
r	8 to 9	2,8, 10, 12, 13 Newtown 30.8.57, La Marette 3 plants 19.9.2000
vr	4 to 7	7. Occasional persisting garden weed
vr	7 to 8	10, gravel at The Old Barn, 27.8.2001. 2nd. JP Essex Fm. 9.11.06
vlf	4 to 10	5,8,10,11,13. Seeds smooth & very shiny
r		Recorded as <i>Montia fontana</i> var. <i>rivularia</i> Det. by SMW 1963as <i>ssp chondrosperma</i> . Seeds have dull tubercles Det. by SMW 1963 as <i>ssp. intermedia</i> . Seeds 3 rows pointed
f	1 to 11	4,7,10. Common garden weed in sandy areas. Infl. diffuse at fruiting
o	5 to 8	5,7,8,11. Slender, inflorescence diffuse at fruiting
r	5 to 6	7. First record 27.5.96
lc	4 to 8	1,2,3,4,5,13. Sand dunes
c	1 to 12	
	3 to 5	5. (As <i>S. apetalla</i> var. <i>glabella</i>). Mannez. 1st record in BIs

THE PLANT LIST

No.	Scientific name	English name	Finder/date
139	<i>Stellaria graminea</i> #	Lesser Stichwort	JO 1953
140	<i>Cerastium tomentosum</i>	Snow-in-Summer	CMR 1950
141	<i>Cerastium fontanum</i> ssp. <i>vulgare</i>	Common Mouse-ear	CCB 1838
142	<i>Cerastium fontanum</i> ssp. <i>holosteoides</i>	Common Mouse-ear	CCB 1838
143	<i>Cerastium glomeratum</i>	Sticky Mouse-ear	EDM 1901
144	<i>Cerastium diffusum</i>	Sea Mouse-ear	CCB 1838
145	<i>Cerastium semidecandrum</i>	Little Mouse-ear	CCB 1838
146	<i>Moenchia erecta</i>	Upright Chickweed	EDM 1901
147	<i>Sagina nodosa</i>	Knotted Pearlwort	CCB 1838
148	<i>Sagina subulata</i>	Heath Pearlwort	CCB 1838
149	<i>Sagina procumbens</i>	Procumbent Pearlwort	CCB 1838
150	<i>Sagina apetala</i> ssp. <i>apetala</i>	Annual Pearlwort	EDM 1901
151	<i>Sagina apetala</i> ssp. <i>erecta</i>	Common Pearlwort	CCB 1838
152	<i>Sagina maritima</i>	Sea Pearlwort	CCB 1838
153	<i>Scleranthus annuus</i> #	Annual Knawel	CCB 1838
154	<i>Herniaria glabra</i> #	Smooth Rupturewort	CCB 1838
155	<i>Herniaria ciliolata</i> ssp. <i>ciliolata</i> #	Fringed Rupturewort (RDBk)	EDM 1901
156	<i>Polycarpon tetraphyllum</i> *	Four-leaved Allseed (RDBk)	CCB 1838
157	<i>Spergula arvensis</i>	Corn Spurrey	CCB 1838
158	<i>Spergularia rupicola</i>	Rock Sea-spurrey	EDM 1901
159	<i>Spergularia rubra</i>	Sand-spurrey	CCB 1838
160	<i>Spergularia bocconeii</i>	Greek Sea-spurrey	PR 1979
161	<i>Lychnis flos-cuculi</i>	Ragged-robin	CCB 1838
162	<i>Agrostemma githago</i> # *	Corncockle	CCB 1838
163	<i>Silene nutans</i> # * (Scarce)	Nottingham Catchfly	EDM 1901
164	<i>Silene vulgaris</i>	Bladder Champion	EDM 1901
165	<i>Silene uniflora</i>	Sea Champion	CCB 1838
166	<i>Silene armeria</i>	Sweet William Catchfly	CJ 1993
167	<i>Silene latifolia</i>	White Champion	CCB 1838
168	<i>Silene x hampeana</i>	Pink Champion	JDG 1937
169	<i>Silene dioica</i>	Red Champion	CCB 1838
170	<i>Silene gallica</i> * (Scarce)	Small-flowered Catchfly	CCB 1838
171	<i>Silene gallica</i> var. <i>quinquevulnera</i>	Variegated Catchfly	EDM 1901
172	<i>Silene conica</i> #	Sand Catchfly	EDM 1901

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
	5 to 8	5, 6 Only other record FF 1967
o	4 to 9	4,7,8,9. Garden escape
c	3 to 6	Petals (5) roughly equal sepals. Hairs sparse/dense, not glandular
c	4 to 9	1,2,3,4,5. Less hairy than above ssp. Lvs narrower
f	1 to 12	Petals (5) equal sepals. Hairs abundant some glandular. Sticky
c	1 to 12	Dark-green. Petals (4) shorter than sepals. Abundant glandular hairs
c	3 to 5	Petals (5) shorter than sepals. Abundant hairs. Dries up by June
vr	4 to 6	5, 10. Mannez & La Tchue. 4-merous, capsule has 8 teeth
o	4 to 9	3,4,5,6,7,8. Petals twice as long as sepals (5). Perennial
c	1 to 10	Petals slightly shorter than sepals (5). Perennial
f	5 to 9	2,5,7,8,9,12. Petals minute or absent, sepals 4 or 5.. Perennial
f	3 to 10	2,4,5,7,8. Petals minute/absent, sepals 4, erect in fruit seed >.3mm
lf	3 to 10	7,8. As 152, sepals patent in fruit, seeds<3mm. Lf point .1-.4mm
o	5 to 9	4,5,6,7,8,10. Similar to above 2 sspp, but leaf point <.1mm
vr	4 to 6	9,10,12,14. Sq. 12, 1908, 1951. Refound Sq. 9 BB 1996)
		3,4,5,14. Records ? for next species. Bibette Head JDG
	7 to 8	10. Evergreen perennial, stems hairy on one side Last seen 1956
lf	3 to 11	1,3,5,7,12,13,14. (var. diphyllum recorded JDG 1937)
lc	3 to 12	7,11,12,13. Cultivated ground. var. nana found GCD 1907
c	5 to 10	2,3,6,7. Perennial. Fls. >8mm, stems glandular hairy
lc	5 to 10	2,6,7,11. Old walls and sandy tracks. Ann/bien, upper bracts=lvs.
		11. Trois Vaux. Upper bracts much shorter than leaves
o	4 to 10	7,11,12,13
vr	5 to 8	2,7,12 Refound BB
r	5 to 6	5, 7 12. Petal limb with conspicuous acute scales at base
lf	5 to 6	Scattered in many squares
la	1 to 12	
vr	7 to 8	13
lf	2 to 12	Fls dioecious, calyx 10-veined male, 20-veined female.
c	1 to 12	Flower colour very variable
lc	1 to 12	Most coastal squares
f	4 to 8, 11	4,5,9,10,12,14. Petals yellowish-white
o	6 to 8	1, 10,11, 12. Petals with red blotches
vr	5 to 7	4,5,7,8. Very rare since late 1960s. 2 plants refound ML 1991

THE PLANT LIST

No.	Scientific name	English name	Finder/date
173	<i>Saponaria officinalis</i>	# Soapwort	BHR 1956
174	<i>Petrorhagia nanteuilii</i>	# Childing Pink	OJ 1985
175	<i>Dianthus armeria</i>	# * Deptford Pink (Scarce)	EDM 1900
Polygonaceæ			
176	<i>Persicaria amphibia</i>	Amphibious Bistort	CCB 1838
177	<i>Persicaria maculosa</i>	Redshank	CCB 1838
178	<i>Persicaria lapathifolia</i>	Pale Persicaria	EDM 1901
179	<i>Persicaria hydropiper</i>	# Water-pepper	EDM 1901
180	<i>Fagopyrum esculentum</i>	# Buckwheat	EDM 1901
181	<i>Polygonum oxyspermum</i>	# Ray's Knotgrass	WH 1881
182	<i>Polygonum arenastrum</i>	Equal-leaved Knotgrass	ABJ 1932
183	<i>Polygonum aviculare</i>	Knotgrass	CCB 1838
184	<i>Fallopia japonica</i>	Japanese Knotweed	DMcC 1957
185	<i>Fallopia baldschuanica</i>	Russian Vine	BB 1987
186	<i>Fallopia convolvulus</i>	Black Bindweed	CCB 1838
187	<i>Rheum x hybridum</i>	# Rhubarb	DMcC 1963
188	<i>Rumex acetosella</i>	Sheep's Sorrel	CCB 1838
189	<i>Rumex acetosa</i>	Common Sorrel	CCB 1838
190	<i>Rumex hydrolapathum</i>	# Water Dock	BB 1987
191	<i>Rumex crispus</i>	Curled Dock	CCB 1838
192	<i>Rumex x pseudopulcher</i>	# A hybrid Dock	DMcC 1953
193	<i>Rumex conglomeratus</i>	Clustered Dock	EDM 1901
194	<i>Rumex sanguineus</i> var. <i>viridis</i>	Wood Dock	EDM 1901
195	<i>Rumex sanguineus</i> var. <i>sanguineus</i>	Wood Dock	BB 1987
196	<i>Rumex rupestris</i>	# * Shore Dock (RDBk)	DMcC 1958
197	<i>Rumex pulcher</i>	Fiddle Dock	EDM 1901
198	<i>Rumex obtusifolius</i>	Broad-leaved Dock	CCB 1838
Plumbaginaceæ			
199	<i>Limonium vulgare</i>	# Common Sea-lavender	AS 1965
199a	(<i>Limonium auriculae-ursifolium</i>)	Broad-leaved Sea-lavender	
200	<i>Limonium normanicum</i>	# Alderney Sea-lavender	CRPA 1900
201	<i>Limonium binervosum</i>	Rock Sea-lavender	??R 1842

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
		No site. Only record No site. Only record 13 Only other record ABJ 1932
lc	6 to 9	1,2,4,5,7. Wet places and ponds
lc	6 to 11	Cultivated ground. Common garden weed
o	7 to 11	7,10,11,12. Cultivated ground
	7 to 9	7. Possibly the only record Rochers Hill & Reuter's valley. Only record 4,7. Platte Saline & Corblets. Only records
vlc	6 to 10	Forming dense mat to 30cm across, common on sandy tracks.
c	6 to 11	Robust, spreading to 2m. Sandy areas. Frequent garden weed
r	6 to 10	3,7,8. Garden escape. Refound BB
r	6 to 11	7,8,9. Persisting garden escape. Old buildings
o	6 to 11	7,9,11,12. Sandy areas, cultivated land and gardens Probably the only record
la	4 to 11	
c	4 to 9	
1 site	7 to 9	5 Only record
c	5 to 7	Only record
o	3 to 8	4,7. Moist places
f	5 to 9	4,7,8,11,12,13. Green-veined. Woods and hedges, damp places
o	5 to 9	4,5,7,8,11,12,13. Red-veined
	4-8, 11	Only record
f	4-8, 11	All squares. Widespread in small numbers
c	4 to 8	
vr	7 to 8	8. Garden escape Refound BB (as <i>S. lychnidifolia</i>) Alderney plants now all referred to next spp.
vr	5 to 8	5 5 to 7 leaf veins. One colony, variable
r	5 to 8	5,6,10 1 to 3 leaf veins. Finder Rogers (? initials)

THE PLANT LIST

No.	Scientific name	English name	Finder/date
202	<i>Armeria maritima</i>	Thrift	CCB 1838
203	<i>Armeria arenaria</i>	# Jersey Thrift	FLS 1984
Clusiaceæ		Formerly Hypericaceæ	
204	<i>Hypericum androsæmum</i>	Tutsan	NJ 1974
204a	<i>Hypericum perforatum</i>	Perforate St. John's-wort	LP 2005
205	<i>Hypericum tetrapterum</i>	Square-stalked St. J's-wort	CCB 1838
206	<i>Hypericum humifusum</i>	Trailing St. John's-wort	CCB 1838
207	<i>Hypericum liniifolium</i> # (RDBk)	Toadflax-leaved St. J's-wort	EDM 1901
208	<i>Hypericum pulchrum</i>	# Slender St. John's-wort	CCB 1838
209	<i>Hypericum elodes</i>	# Marsh St. John's-wort	CCB 1838
Tiliaceæ		Probably all planted	
210	<i>Tilia platyphyllos</i> # * (Scarce)	Large-leaved Lime	CH 1987
211	<i>Tilia x europea</i>	# Lime	CH 1987
212	<i>Tilia cordata</i>	# Small-leaved Lime	RW/CH 1987
Malvaceæ			
213	<i>Malva moschata</i>	Musk-mallow	EDM 1901
214	<i>Malva sylvestris</i>	Common Mallow	CCB 1838
215	<i>Malva neglecta</i>	# Dwarf Mallow	CCB 1838
216	<i>Lavatera arborea</i>	Tree-mallow	CCB 1838
Cistaceæ			
217	<i>Tuberaria guttata</i>	Spotted Rock-rose	CCB 1838
Violaceæ			
218	<i>Viola odorata</i>	# Sweet Violet	RW/BB 1987
219	<i>Viola riviniana</i>	Common Dog-violet	EDM 1901
220	<i>Viola canina</i>	? Heath Dog-violet	CCB 1838
221	<i>Viola arvensis</i>	Field Pansy	EDM 1901
Tamaricaceæ			
222	<i>Tamarix gallica</i>	Tamarisk	EDM 1901

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
la	3 to 1	Occasionally found with white flowers <i>FLS was an expert on Jersey flora, but ? an error. Only record</i>
vr	6 to 11	'7,8,9,11. Fontaine David 9.9.74
1 site	6 to 7	3, Fort Albert Glacis. Still there BB 1.7.2006
lf	4 to 10	6,7,8,11,12,13. Wet places
o	5 to 9	5,6,7,10,11,12,13,14
1 site now	6 to 8	10,11,12,13,14. Single remaining colony thriving. 56 plants in 1995
vr	5 to 9	6,7,10,12 11 Last record 1901
vr	6 to 7	Planted
vr	6 to 7	Planted
r	6 to 7	Planted
f		7,9,10 Sq 5 LP 2006 Previous record, 2 plants Blaye AB/AKJ 1932
c	5 to 12	
vr	6 to 9	4,5,8,10. Refound JDG
o	5 to 11	1,2,3,4,5,7,8,10. Specimen to 9'4" high, Braye Hill 1989
2 sites	4 to 7	12,13. 1995 over c. 200 yds of cliffs. Drops petals by noon
vr	9 to 6	9. In the Ash wood in Water Lanes
a	1 to 12	<i>Possibly in error for previous species Only record</i>
o/vla	5 to 10	7,8,11,12,13. Cultivated ground
o	1 to 12	2,3,4,5,7,8. Scattered in old hedges. Some over 100 yrs old

THE PLANT LIST

No.	Scientific name	English name	Finder/date
Frankeniaceæ			
223	<i>Frankenia laevis</i>	# * Sea Heath (Scarce)	JNCW 1956
Cucurbitaceæ			
224	<i>Cucurbita pepo</i>	# Marrow	NJ 1974
Salicaceæ			
225	<i>Populus alba</i>	White Poplar	EDM 1901
226	<i>Populus x canescens</i>	Grey Poplar	ABJ 1933
227	<i>Populus nigra</i>	# Black Poplar	EDM 1901
228	<i>Populus x canadensis</i>	Hybrid Black-poplar	?
229	<i>Populus candicans</i>	Balm of Gilead	OJ 1987
230	<i>Salix pentandra</i>	# Bay Willow	BB 1987
231	<i>Salix fragilis</i>	Crack Willow	CCB 1838
232	<i>Salix alba</i>	White Willow	BB 1987
233	<i>Salix purpurea</i>	# Purple Willow	EDM 1901
234	<i>Salix daphnoides</i>	# Violet Willow	MM 1986
235	<i>Salix viminalis</i>	# Osier	EDM 1901
236	<i>Salix x stipularis</i>	# Eared Osier	CCB 1838
237	<i>Salix x sericans</i>	# Broad-leaved Willow	EDM 1901
238	<i>Salix x smithiana</i>	# Silky-leaved Osier	EDM 1901
239	<i>Salix caprea</i>	# Goat Willow	EDM 1901
240	<i>Salix cinerea</i>	# Grey Willow	EDM 1901
241	<i>Salix cinerea ssp. oleifolia</i>	Rusty Sallow	CMR 1949
242	<i>Salix aurita</i>	# Eared Willow	CCB 1838
243	<i>Salix babylonica</i>	# Weeping Willow	CH 1987
244	<i>Salix matsudana</i>	# Contorted Willow	RW 1986
Brassicaceæ		Formerly Cruciferæ	
245	<i>Sisymbrium orientale</i>	# Eastern Rocket	CMR 1953
246	<i>Sisymbrium officinale</i>	Hedge Mustard	CCB 1838
247	<i>Alliaria petiolata</i>	Garlic Mustard	CCB 1838
248	<i>Arabidopsis thaliana</i>	# Thale Cress	CMR 1954
249	<i>Erysimum cheiri</i>	Wallflower	FLS 1960
250	<i>Hesperis matronalis</i>	# Dame's-violet	AS 1965

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
	7 to 8	
		Waste heap at Banquage. Probably the only record
f	2 to 3	Scattered. Suckers freely. Much planted by States
o	3	4,5,9,13. Up to 12.5m in Val du Saue
o	2 to 5	8,10. Up to 22m.
vr	2 to 5	10. Another record BB 1989
o	3	7,8,10. Up to 9m. Some seedlings found. Suckers
vr	6 to 7	5. Twigs red-brown, hairless, glossy
o	4 to 5	6,7,8,10,11,13. Twigs very brittle
vr	4 to 7	4,5,7,9,10,13. Twigs usually yellowish-redish, downy
vr	3 to 5	10. Twigs hairless, usually purplish-brown
vr	2 to 5	10. Twigs violet-brown. glossy with dense
r	2 to 4	4,5,7,10. Twigs yellow-brown
		Assisted by W. Borer. No site. 7. Leaves densely grey pubescent underneath.
vr	3 to 4	7,8,9 7,8,12. Mouriaux & Reuter's Valley. Only other Vau
	3 to 4	Only record, probably in mistake for next species
lc	3 to 4	Common species in Alderney. Lvs. stiff rust-coloured
		12. Not present in other C.Is. in 1838. Last record 1901
o	3 to 4	8. Probably hybrid with <i>S. alba</i> . Planted in gardens
vr	3 to 6	7,13. Planted Val du Saue & Platte Saline
	6 to 8	13. (As <i>Erysimum orientale</i> .) a few plants Val du Saue Only recor
f	4 to 12	3,7,9,10,13.
o	3 to 8	3,7,8,9,10,13
r	4 to 6	3,7,8,10.track to Frying Pan 8.5.2007
o	1 to 12	5,8. Tops of old walls in town
vr	5 to 7	Mostly a garden weed. Recorded inAtlas 1962

THE PLANT LIST

No.	Scientific name	English name	Finder/date
251	<i>Malcolmia maritima</i>	# Virginia Stock	DMcC 1973
252	<i>Matthiola incana</i>	* Hoary Stock (RDBk)	EV 1929
253	<i>Barbarea intermedia</i>	# Medium-flowered Winter-cress	DMcC 1963
254	<i>Barbarea verna</i>	# American Winter-cress	EDM 1901
255	<i>Rorippa nasturtium-aquaticum</i>	Water-cress	CCB 1838
256	<i>Rorippa x sterilis</i>	# Hybrid Water-cress	DMcC 1958
257	<i>Rorippa microphylla</i>	# Narrow-fruited Water-cress	DMcC 1958
258	<i>Armoracia rusticana</i>	Horse-radish	CCB 1838
259	<i>Cardamine flexuosa</i>	Wavy Bitter-cress	EDM 1901
260	<i>Cardamine hirsuta</i>	Hairy Bitter-cress	EDM 1901
260a	<i>Cardamine pratensis</i>	Ladies' Smock	BB 2002
261	<i>Arabis hirsuta</i>	Hairy Rock-cress	CCB 1838
262	<i>Aubrieta deltoides</i>	# Aubretia	BB 1987
263	<i>Lunaria annua</i>	Honesty	DMcC 1957
264	<i>Bertoroa incana</i>	# Hoary Alison	FJH 1884
265	<i>Lobularia maritima</i>	Sweet Alison	WH 1881
266	<i>Erophila verna</i>	Common Whitlow-grass	EDM 1901
267	<i>Cochlearia officinalis</i>	Common Scurvy-grass	EDM 1901
268	<i>Cochlearia danica</i>	Danish Scurvy-grass	CCB 1838
269	<i>Capsella bursa-pastoris</i>	Shepherd's-purse	CCB 1838
270	<i>Teesdalia nudicaulis</i>	Shepherd's Cress	EDM 1901
271	<i>Thlaspi arvense</i>	# Field Penny-cress	BB 1987
272	<i>Lepidium sativum</i>	# Garden Cress	EDM 1903
273	<i>Lepidium heterophyllum</i>	# Smith's Pepperwort	EDM 1901
274	<i>Lepidium draba ssp. draba</i>	Hoary Cress	EDM 1901
275	<i>Lepidium ruderales</i>	# Narrow-leaved Water Pepper	EDM 1901
276	<i>Coronopus squamatus</i>	Swine-cress	CCB 1838
277	<i>Coronopus didymus</i>	Lesser Swine-cress	EDM 1901
278	<i>Diplotaxis tenuifolia</i>	Perennial Wall-rocket	CCB 1838
279	<i>Diplotaxis muralis</i>	Annual Wall-rocket	EDM 1901
280	<i>Brassica oleracea</i>	# * Sea Cabbage (Scarce)	GCD 1932
281	<i>Brassica napus</i>	Rape	CCB 1838
282	<i>Brassica rapa</i>	Turnip	Atlas 1962
283	<i>Brassica nigra</i>	# Black Mustard	HCP 1969

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
vr	5 to 7	Garden escape
lc	1 to 12	1,2,3,4,5,7,10. With white flowers only, at Corblets Quarry
	5 to 8	9
vr	1 to 12	7,8,9,12. Nowadays only a garden escape
lf	5 to 12	7,8,11,12,13
r	5 to 9	7
r	5 to 8	7 (Refound BB 1993. Clonque Quarry)
o	5 to 10	4,7,10 7,8,9,10,13. Claimed refound FH 1981, no voucher. Doubtful
c	2 to 8	
1 site lf	4	Large area along stream bed at Ladysmith. 1st. island record
f	2 to 8	4,5,8,9
r	1 to 8	4,10. Garden escape
o	3 to 6	8,9,10 Braye Bay Possibly the only record
c	1 to 12	All squares except cliffs
lf	3 to 4	4,5,7,8
r	3 to 6	2,3,4,5,6,7,12,13
c	2 to 7	Coastal only
o	1 to 12	Scattered plants
lc	2 to 4	4,5,10,11,12,13
vr	5 to 7	12,13 Butes Hill. Only record Fort Tourgis Only other records AB/AKJ 1933 'increased', 1 plant
r	3 to 6	4,7 By Hammond Memorial & Platte Saline jct Picaterre 2, Crabby Bay 20 plants 1899 & 1900. Only records
o	5 to 7	7,8. Stem glabrous. Petals longer than sepals
c	1 to 12	Vigorous garden weed. Strong smell if crushed. Petals shorter or 0
la	4 to 1	1,2,3,4,7. Branched leafy stem
vo	6 to 10	3,7,8. Stem only leafy near base In 'Comital Flora' "ALDERNEY". Appears to be the only record
o	4 to 8	11,12,13. Buds overtop open fls. Leaves clasping stem at base
r	1 to 12	4,7,12. Open fls overtop buds. Stem leaves clasping
r	4 to 9	2,3,11. Leaves petiolate. Determined SMW Only record

THE PLANT LIST

No.	Scientific name	English name	Finder/date
284	<i>Sinapis arvensis</i>	Charlock	CCB 1838
285	<i>Sinapis alba</i>	White Mustard	CCB 1838
286	<i>Eruca vesicaria</i> ssp. <i>sativa</i> #	Garden Rocket	RW 1980
287	<i>Coincya monensis</i> ssp. <i>recurvata</i> #	Wallflower Cabbage	CCB 1838
288	<i>Hirschfeldia incana</i>	Hoary Mustard	CCB 1838
289	<i>Cakile maritima</i>	Sea Rocket	CCB 1838
290	<i>Crambe maritima</i>	Sea-kale	WH 1881
291	<i>Raphanus raphanistrum</i> ssp. <i>raphanistrum</i>	Wild Radish	EDM 1901
292	<i>Raphanus raphanistrum</i> ssp. <i>maritimus</i>	Sea Radish	CCB 1838
293	<i>Raphanus sativus</i> #	Garden Radish	DMcC 1962
Resedaceæ			
294	<i>Reseda luteola</i>	Weld	CCB 1838
295	<i>Reseda lutea</i> #	White Mignonette	CCB 1838
Ericaceæ			
296	<i>Calluna vulgaris</i>	Heather	CCB 1838
297	<i>Erica cinerea</i>	Bell Heather	CCB 1838
Primulaceæ			
298	<i>Primula vulgaris</i>	Primrose	EDM 1901
299	<i>Primula veris</i>	Cowslip	CMR 1954
300	<i>Cyclamen hederifolium</i> #	* Cyclamen (RDBk)	BH 1988
301	<i>Anagallis tenella</i>	Bog Pimpernel	CCB 1838
302	<i>Anagallis arvensis</i> ssp. <i>arvensis</i>	Scarlet Pimpernel	CCB 1838
303	<i>Anagallis arvensis</i> ssp. <i>foemina</i>	Blue Pimpernel #	EDM 1901
304	<i>Anagallis minima</i> #	Chaffweed	EDM 1901
305	<i>Glaux maritima</i> #	Sea-milkwort	EDM 1901
306	<i>Samolus valerandi</i>	Brookweed	CCB 1838
Hydrangeaceæ			
307	<i>Hydrangea macrophylla</i> #	Hydrangea	BB 1987

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
c	1 to 12	Cultivated or bare ground. Lvs ± pinnately lobed, terminal largest
r	6 to 8	Lvs. deeply pinnately lobed, terminal the v. largest. Long fruit beak Whitish petals violet edged. 1st C.I. record. Only Alderney record 3. In GCD 1932 'Native'. Refound HCP 1969, Determined by SMW
la	1 to 12	1,2,3,4,5,6,7. Fruit c. 10mm adpressed to stem, beak 3-5mm
lf	6 to 9	1,2,3,4,5,6. Sandy bays above strand line. Fls pale
vlf	3 to 10	1,2,4,5. Young lvs green or mauve. Was scarce, spreading
o	3 to 11	2,3,4,5,7,8,9,11. Fruit segments cylindrical
la	1 to 12	1-7,9,10. Fls often white. Biennial/perennial. Fruit segments globose Only record
r	5 to 9	2,3,4,5,7,8,9,10. Stem stiff, erect, Petals yellow, 3 carpels. Biennial
vr	5 to 8	3,8. Stem branched, ascending. Petals white, 4 carpels. Annual
lf	4 to 12	4,5,6,10,11,12
lc	1 to 12	5,6,7,9,10,11,12,13. With white flowers, EDM 1908
o	1 to 12	6,7,8,9,10,11. Rarely in any quantity
vr	3 to 7	9,10. Stranger's Cemetery only now
vr	8 to 10	4,5,7,8
vr	5 to 8	11,12. Over 50m. in Baxter's Valley 1996
c	1 to 12	Several colour forms, pale, flesh, red and blue. Fls close in afternoon
vr		12. Recorded as var azure. Others CMR 1950, AS 1955, AT 1981
	6 to 7	6,12. Only other records ABJ 1934, CMR 1954
r	6 to 10	5 Rock ledges near Limonium (and on Burhou)
o	4 to 8	4,5,11,12,13
vr	5 to 2	8,13. Garden escape

THE PLANT LIST

No.	Scientific name	English name	Finder/date
Grossulariaceæ			
308	<i>Escallonia macrantha</i>	Escallonia	JCNW 1956
309	<i>Ribes nigrum</i> #	Black Currant	BB 1988
310	<i>Ribes sanguineum</i> #	Flowering Currant	BB 1988
311	<i>Ribes uva-crispa</i> #	Gooseberry	MMcW 1962
Crassulaceæ			
312	<i>Crassula helmsii</i>	New Zealand Pigmy-weed	ML 1983
313	<i>Crassula pubescens</i> #	Jersey Pigmy-weed	BB1991
314	<i>Umbilicus rupestris</i>	Navelwort	CCB 1838
315	<i>Aeonium x velutinum</i>		BB 1991
316	<i>Sedum spectabile</i>	Butterfly Stonecrop	MM 1987
317	<i>Sedum telephium</i> #	Orpine	EDM 1901
318	<i>Sedum spurium</i> #	Caucasian-stonecrop	DMcC 1963
319	<i>Sedum acre</i>	Biting Stonecrop	CCB 1838
320	<i>Sedum album</i>	White Stonecrop	DMcC 1953
321	<i>Sedum anglicum</i>	English Stonecrop	CCB 1838
Saxifragaceæ			
322	<i>Bergenia</i> spp. (? <i>crassifolia</i>) #	Elephant's-ears	BB 1987
323	<i>Saxifraga x urbium</i> #	Londonpride	DMcC 1963
324	<i>Saxifraga tridactylites</i>	Rue-leaved Saxifrage	EDM 1901
Rosaceæ			
327	<i>Rubus caesius</i>	Dewberry	EDM 1901
328	<i>Rubus fruticosus</i> agg.	Blackberry. (Many subspecies)	CCB 1838
329	<i>Rubus</i> series <i>rhamifolia</i>	Blackberry	CCB 1838
330	<i>Rubus leucostachys</i>	Blackberry	GCD 1906
331	<i>Rubus corylifolius</i>	Blackberry	GCD 1906
332	<i>Rubus rusticanus</i>	Blackberry	GCD 1907
333	<i>Potentilla fruticosa</i>	Shrubby Cinquefoil	BB 1987
325	<i>Filipendula vulgaris</i> #	Dropwort	DMcC 1963
326	<i>Rubus idaeus</i>	Raspberry	EDM 1901
334	<i>Potentilla anserina</i>	Silverweed	CCB 1838
335	<i>Potentilla erecta</i>	Tormentil	CCB 1838

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
o	1 to 12	2,4,5,7,8,10. Much used as hedging. self-seeds regularly
vr	3 to 5	4,9, Longis Road among pines. Berry's quarry
vr	3 to 6	6,10 Bastion of Fort Clonque. Longis Bay
vr	2 to 5	8. Ash wood in Water Lanes BB
vlf	3 to 9	4,5,10. Tank on Essex, Mannez & Corblets. New Bailiwick record.
	5 to 8	5 Only record
a	4 to 9	Up to 90cm at Val du Saue
vr	5 to 9	5, Determined DG, Kew 1996
vr	8 to 10	7. Garden escape
		4. Only record
		Only record
c	5 to 9	Seems to prefer alkaline sandy areas
o	5 to 8	2,4,5,6,7,8,9,10
c	5 to 8	2-9,13. Seems to prefer slightly acid soils
vr	11 to 8	4. Garden escape
vr	5 to 7	Garden escape Only record
lf	3 to 5	4,5,8
c	4 to 8	All leaves in 3s. Fls white, earliest. Few large drupelets in fruit
a	5 to 11	Most leaves in 5s. fls pinkish, later. Fruits more, smaller, drupelets
vr	6 to 7	Garden escape
	6	Only record
		10. In GCD 1932 "Alderney"
f	5 to 10	7,11,12. Some plants silver both sides of leaf
c	5 to 11	2,4,5,6,7,10,11,12,13 Mainly cliffs and quarries

THE PLANT LIST

No.	Scientific name	English name	Finder/date
336	<i>Potentilla x mixta</i>	Hybrid Cinquefoil	CMR 49-63
337	<i>Potentilla reptans</i>	Creeping Cinquefoil	CCB 1838
338	<i>Potentilla sterilis</i>	Barren Strawberry	EDM 1901
339	<i>Fragaria ananassa</i>	Garden Strawberry	DMcC 1973
339a	<i>Fragaria vesca</i>	Wild Strawberry	BB1997
340	<i>Geum urbanum</i> #	Herb Bennet	EDM 1901
341	<i>Agrimonia eupatoria</i>	Agrimony	EDM 1901
342	<i>Sanguisorba minor</i> ssp. <i>minor</i>	Salad Burnet #	JM 1959
343	<i>Aphanes arvensis</i>	Parsley-piert	CCB 1838
344	<i>Aphanes australis</i>	Lesser Parsley-piert	JO 1953
345	<i>Rosa multiflora</i>	Many-flowered Rose	DMcC 1953
346	<i>Rosa pimpinellifolia</i>	Burnet Rose	CCB 1838
347	<i>Rosa x involuta</i> #	(<i>R. pimpinell. x R. sherardii</i>)	EDM 1901
348	<i>Rosa rugosa</i>	Japanese Rose	CMR 1950
348a	<i>Rosa x alba</i> #	White Rose of York	BB 1990
349	<i>Rosa stylosa</i>	Short-styled Field-rose	EDM 1901
350	<i>Rosa canina</i>	Dog-rose	CCB 1838
351	<i>Rosa tomentosa</i> #	Harsh Downy-rose	EDM 1901
352	<i>Rosa mollis</i> #	Soft Downy-rose	FF 1967
353	<i>Rosa rubiginosa</i>	Sweet-briar	EDM 1901
354	<i>Rosa micrantha</i> #	Small-flowered Sweet-br	EDM 1901
355	<i>Rosa wicheriana x Shirley Hibberd</i>	Alberic Barbier Rose #	NVM 1967
356	<i>Rosa wicheriana x Gabriel Luiget</i>	Dorothy Perkins Rose	BB 1988
357	<i>Rosa wicheriana x R. setigera</i>	American Pillar Rose #	BB 1986
357a	<i>Prunus cerasifera</i>	Cherry Plum	BB 1997
358	<i>Prunus spinosa</i>	Blackthorn	CCB 1838
359	<i>Prunus domestica</i> ssp. <i>domestica</i>	Wild Plum	DMcC 1957
360	<i>Prunus domestica</i> ssp. <i>insititia</i>	Bullace	BB 1996
361	<i>Prunus x fruticans</i>	Hybrid between <i>P.spin x P.dom</i>	?
362	<i>Prunus avium</i> #	Gean	RW 1987
362a	<i>Prunus cerasus</i> #	Dwarf Cherry	BB 1997
363	<i>Pyrus pyraeaster</i> #	Wild Pear	CJ 1980
364	<i>Pyrus communis</i> # *	Pear (RDBk)	DMcC1960
365	<i>Malus sylvestris</i>	Crab Apple	EDM 1901
366	<i>Malus domestica</i> #	Apple	BB 1989
367	<i>Sorbus aucuparia</i> #	Rowan	BB 1987

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
f	6 to 9	7,11,12.
c	6 to 10	Fixed dune areas
o	2 to 5	7,8,12. Next CMR 1954 Refound JB 1992
vr	2, 5 to 12	Garden escape or throwout
vr	6 to 7	8. N. wall of Parish Churchyard. 2 plants in fl. & fr. 1st island record
o	2 to 8	2,7,8,9. 'Damp places
vr	5 to 9	4, 8, 9 E. rim Battery Quarry
	5 to 8	2,4,9. Garden escape Only record
r	3 to 6	5,7,11
a	2 to 5	4,5,6,7,9,10,11,13,14
1 site	6 to 8	10 Refound BB 1989 at The Old
r	5 to 9	3,4,12,13 on walls bounding Blaye. (Braye Common, BB1988 only)
		10. Only record
r	4 to 11	5,7,8,10. Much used as hedging. Suckers and seeds
1 site	7 to 9	11. Large shrub in hedge, Grande Blaye nr. Airport
o	6 to 8	3,4,5,7,10,11,12,13 Refound JES 1971, BB 1992
o	6 to 8	2-14
		7. Only record
		7. Only record
o	6 to 7	4,10,11,12,13,14
o	6 to 8	3,4,5,7,12,13,14 Refound BB 1992
1 site	1 to 12	7 Rim of sand pit for 25+ years. Surviving in hedge 1991-2007
4 sites	7 to 9	6,7,11,13. Vau Pommier, S. cliffs. Probably planted 30+ years ago
2 sites	6 to 10	2,6. Bank by "Pilot's View" Crabby for 40+ years. Probably planted
r	2 to 4	8,9. LA Vallée and Val Fontaine. Several trees suckering
c	2 to 5	7,9-14
r	2 to 5	8, 9 specimens nearer type. Most others seem to be hybrids
1 site ?	2 to 5	12. Small copse in Le Grand Val
la	2 to 6	Many plants seem to be sterile. Few or no fruits.
1 site		9? Single specimen
vr	6 to 7	8. Les Venelles, Hedge suckering. Mostly destroyed in 2005
1 site	3 to 5	2 Single specimen in York Hill Quarry
1 site	3 to 5	4 Single specimen in Mannez Quarry
vr	4 to 6	4,5,7, 8,,9,12. A few old survivors round ruins
vr	4 to 6	4,5,7,8,9,12. In roadside verges or scrub. Possibly from seed
vr	5 to 6	13

THE PLANT LIST

No.	Scientific name	English name	Finder/date
368	<i>Sorbus intermedia</i>	# Swedish Whitebeam	BB 1987
369	<i>Cotoneaster simonsii</i>	# Himalayan Cotoneaster	CMR 1958
370	<i>Crataegus pedicellata</i>	# Pear-fruited Cockspur-thorn	BB 1992
371	<i>Crataegus monogyna</i>	Hawthorn	CCB 1838
372	<i>Crataegus laevigata</i>	? Midland Hawthorn	CCB 1838
Fabaceæ		Formerly Leguminosæ	
373	<i>Robinia pseudacacia</i>	# False Acacia	BB 1987
374	<i>Onobrychis viciifolia</i>	# Sainfoin	EDM 1901
375	<i>Anthyllis vulneraria</i> agg.	Kidney Vetch	CCB 1838
376	<i>Anthyllis vulneraria</i> ssp. <i>vulneraria</i>	var. <i>vulneraria</i>	ET 1963
377	<i>Anthyllis vulneraria</i> ssp. <i>vulneraria</i>	var. <i>langei</i>	ET 1963
378	<i>Anthyllis vulneraria</i> ssp. <i>maritima</i>	#	1971
379	<i>Lotus corniculatus</i>	Common	CCB 1838
380	<i>Lotus pedunculatus</i>	Greater	CCB 1838
381	<i>Lotus subbiflorus</i>	Hairy Bird's-foot-trefoil;	CCB 1838
382	<i>Lotus angustissimus</i>	Slender Bird's-foot-trefoil	CCB 1838
383	<i>Ornithopus perpusillus</i>	Bird's-foot	CCB 1838
384	<i>Ornithopus pinnatus</i>	Orange Bird's-foot	MD 1881
385	<i>Vicia cracca</i>	# Tufted Vetch	EDM 1901
386	<i>Vicia villosa</i>	# Fodder Vetch	EDM 1900
387	<i>Vicia hirsuta</i>	Hairy Tare	CCB 1838
388	<i>Vicia tetrasperma</i>	# Smooth Tare	EDM 1901
389	<i>Vicia sativa</i> ssp. <i>nigra</i>	Narrow-leaved Vetch	EDM 1901
390	<i>Vicia sativa</i> ssp. <i>sativa</i>	Common Vetch	CCB 1838
391	<i>Vicia lathyroides</i>	Spring Vetch	CMR 1952
392	<i>Vicia bithynica</i>	# * Bithynian Vetch (Scarce)	DMcC 1963
393	<i>Vicia faba</i>	# Broad Bean	DMcC 1963
394	<i>Lathyrus pratensis</i>	# Meadow Vetchling	EDM 1901
395	<i>Lathyrus grandiflorus</i>	# Two-flowered	BB 1988
396	<i>Lathyrus latifolius</i>	# Broad-leaved	PHAS 1969
397	<i>Lathyrus aphaca</i>	# * Yellow Vetchling Scarce)	JES 1970
398	<i>Ononis reclinata</i>	# * Small Restharrow RDBk)	CCB 1828
399	<i>Ononis reclinata</i> f. <i>albiflorus</i>	# White Small-restharrow RDBk)	OJ 1979
400	<i>Ononis repens</i>	Common Restharrow	CCB 1838

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
vr	5	Planted by States in several squares
vr		
1 site	5 to 6	8. Single specimen in Old Churchyard. Fls 3-4weeks later than next
c	4 to 6	Next record almost certainly meant for this sp. In Flora Armoricaine. error, but recorded by RW & CH June 1986
vr	3 to 5	2,8 Churchyard & 1 tree along stream side. Platte Saline, (planted)
c	2 to 11	4,7,8. Butes Only other record 1908
r		Determined by JC
lc		Determined by JC. Stace says most CI plants are this ssp. Recorded in Flora Armoricaine
a	4 to 11	
o	5 to 10	3,4,5,7,9,11,12,13. Mostly in damp places
la	4 to 9	Cliffs and low coasts
o	6 to 8	5,10,11,12,13,14. Refound OJ 1982, was rare, now occasional
vla	4 to 7	4,5,9,10,11,12,13,14
lf	5 to 7	4,5,9,10,11,12,13,14
vr	5 to 8	4,7,10,11,12
c	4 to 7	9. Only other records CPH 1902, ABJ 1932
vr	6 to 7	2-7fls. 2-seeded pods
c	4 to 10	5,7,12,13. 1-2fls, 4-seeded pods
c	1 ro 11	Plant very slender often procumbent
r	4 to 6	Robust plant. Many white flowered plants at Crabby Bay 1979
2 sites	4 to 8	5,10,11,12. Annual to 20cm. Mannez quarry, BB June 1995
vr		2,4. Robust plant , fls purple & white Along cliff edge, Crabby Bay
vr	6 to 8	7. Garden escape
vr	6 to 9	4,9,10. Perennial Flowers yellow, stem angled but not winged
vr	6 to 9	2,5,8. Fls. >25mm, pinkish/purple
vr	6 to 9	5,8,9. Fls 15-30mm, pink, brighter than previous sp.
vr		Annual. Mature lvs reduced to tendrils. Refound JP & Soc G. 1986
vr	5 to 6	4,5. Not seen 1988-92 Refound coast MM 1993, Mannez hill
vr	5 to 6	5 Refound OJ 1979, BB 1988
c	4 to 10	Occasionally with white flowers

THE PLANT LIST

No.	Scientific name	English name	Finder/date
401	<i>Melilotus altissimus</i>	# Tall Melilot	JDG 1937
402	<i>Melilotus albus</i>	White Melilot	GCD 1906
403	<i>Melilotus officinalis</i>	Ribbed Melilot	EDM 1903
404	<i>Melilotus indicus</i>	Small Melilot	EDM 1899
405	<i>Medicago lupulina</i>	Black Medick	CCB 1838
406	<i>Medicago sativa</i> ssp. <i>sativa</i>	Lucerne	EDM 1901
406a	<i>Medicago sativa</i> ssp. <i>falcata</i>	# Sickle Medick	LP 2006
407	<i>Medicago polymorpha</i>	Toothed Medick	CCB 1838
408	<i>Medicago arabica</i>	Spotted Medick	CCB 1838
409	<i>Trifolium ornithopodioides</i>	Bird's-foot Clover	EDM 1901
410	<i>Trifolium repens</i>	White Clover	CCB 1838
411	<i>Trifolium occidentale</i>	* Western Clover (Scarce)	RWW 1962
412	<i>Trifolium hybridum</i>	# Alsike Clover	BHR 1956
413	<i>Trifolium glomeratum</i>	* Clustered Clover(Scarce)	EDM 1901
414	<i>Trifolium suffocatum</i>	Suffocated Clover	EDM 1901
415	<i>Trifolium fragiferum</i>	# Strawberry Clover	EDM 1901
416	<i>Trifolium fragiferum</i> ssp. <i>bonannii</i> *	Strawberry Clover (Southern)	JH 1932
417	<i>Trifolium campestre</i>	Hop Trefoil	CCB 1838
418	<i>Trifolium dubium</i>	Lesser Trefoil	CCB 1838
419	<i>Trifolium micranthum</i>	Slender Trefoil	CCB 1838
420	<i>Trifolium pratense</i>	Red Clover	CCB 1838
421	<i>Trifolium medium</i>	# Zigzag Clover	JO 1953
422	<i>Trifolium stellatum</i>	# Starry Clover	ABJ/HKAS
423	<i>Trifolium incarnatum</i> ssp. <i>incarnatum</i> #	Crimson Clover	EDM 1901
424	<i>Trifolium striatum</i>	Knotted Clover	CCB 1838
425	<i>Trifolium bocconeii</i> (RDBk) *	Twin-headed Clover #	DMcC 1965
426	<i>Trifolium scabrum</i>	Rough Clover	CCB 1838
430	<i>Laburnum anagyroides</i>	# Laburnum	BB 1987
430a	<i>Spartina juncea</i>	# Spanish Broom	BB 1989
431	<i>Cytisus scoparius</i> ssp. <i>scoparius</i>	Broom #	CCB 1838
432	<i>Cytisus scoparius</i> ssp. <i>maritimus</i>	Prostrate Broom	M'Nab 1850
427	<i>Trifolium arvense</i>	Hare's-foot Clover	CCB 1838
428	<i>Trifolium subterraneum</i>	Subterranean Clover	EDM 1901
429	<i>Lupinus arboreus</i>	Tree Lupin	DMcC 1957
433	<i>Ulex europaeus</i>	Gorse	CCB 1838

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
r	5 to 8	2,3,4,8. Biennial/perennial. Fls standard, wings, keel all equal
o	6 to 9	2,3,7. Refound NVM, Crabby/Platte Saline,
lf	5 to 9	2,3,7,8. Annual, Fls. keel shorter
o	5 to 10	2,3,4,5,7,8,12. Fls. standard longer
c	4 to 10	Annual or SLperennial. Fls 10-50. Fruit single turn, ripe net black
f	4 to 10	Sandy parts. Flower colour varies from v. dark purple to white
vr	5 to 8	4. Campsite road to Bibette Head 1 large plant. 1st. island record
r	5 to 8	3,4,8,11,12. Fls. 1-5. Fruit flat, 1½-6 turns, spiny
a	3 to 9	Fls 1-4. Fruit globose spiral, 3-7 turns, spines grooved ½way
o	4 to 8	4,5,6,7,10,11,12,13,14. Fls 1-4 in raceme, pedicels >1mm
a	1 to 11	Fls globose, peduncles up to 20cm, a month later than next sp.
lc	3 to 8	Smaller, peduncles short. Cliff and dune edges or within 100m sea 4,7. Fls usually pink & white. Only other record FF 1967, Longis
o	6 to 8	4,5,13,14. Cliffs. Fls 4-5mm purple. 2-seeded fruit encl. by calyx
o	3 to 6	4,5,10-14. Plant 5cm across, whitish 3mm fls mostly near root
vr	6 to 7	4,7,11,13. Marshy places. Pink, fls 5-7mm, Status now considered doubtful (Stace)
vf	1, 4 to 8	Bare places. Yellow racemes globose 10-15mm with 20-40 fls.
c	4 to 10	Racemes 6-8mm with 5-20 fls
o	4 to 8	All ex. 1. Racemose c5mm with 2-6 fls
c	4 to 12	Colour varies considerably pinkish-purple, pale pink, occ. cream
vr	4 to 8	7,8. Rhizomatous, fls terminal in globose red-purple raceme Short turf, Platte Saline. Only record 4,10, Above Hanging Rock Only other record JDG 1937
o	5 to 7	5,7,9,10,11,12,13,14. Coastal. Pink racemes sessile, axils & terminal
	5 to 6	Trois Vaux Only record
f	5 to 8	Cliff tops, dune edges, short turf. Pubescent, fls white/pale
o	5 to 6	8,9. Entirely in gardens
vr	4 to 6	9. Church Glebe land on Les Rochers 20.6.89
r	4 to 7, 12	4,11,13,14. Mainly on cliff tops
lc	3 to 6, 9	11,12,13,14. Cliffs
lf	3 to 6	Mainly cliffs & bare ground. Whitish terminal racemes, cylindrical
f	3 to 8	1,5,6, & Cliffs. Raceme axillary 2-5 whitish fertile fls, sterile calices
o	3 to 9	Sandy places close to sea. Occasionally with white flowers
a	1 to 12	Mainly on cliffs & rough places. Flowering peak in May

THE PLANT LIST

No.	Scientific name	English name	Finder/date
434	<i>Ulex gallii</i>	Western Gorse	EDM 1901
435	<i>Ulex minor</i>	? Dwarf Gorse	CCB 1838
Elæagnaceæ			
436	<i>Hippophae rhamnoides</i>	# * Sea-Buckthorn (Scarce)	BB 1986
437	<i>Elæagnus commutata</i>	# Elæagnus species	BB 1996
438	<i>Elæagnus glabra</i>	Elæagnus species	OJ1977
439	<i>Elæagnus ebbingei</i>	Elæagnus species	BB1988
Haloragaceæ			
440	<i>Myriophyllum verticillatum</i>	Whorled Water-milfoil	BB 1987
441	<i>Myriophyllum aquaticum</i>	# Parrot's-feather	BB1988
442	<i>Myriophyllum spicatum</i>	Spiked Water-milfoil	DMcC 1957
Gunneraceæ			
443	<i>Gunnera tinctoria</i>	# Giant-rhubarb	BB 1987
Lythraceæ			
443b	<i>Lythrum salicaria</i>	Purple Loosestrife	LP 2007
Myrtaceæ			
443a	<i>Myrtus ugni</i>	# Myrtle	HH 1997
Onagraceæ			
444	<i>Epilobium hirsutum</i>	Great Willowherb	JO 1953
445	<i>Epilobium parviflorum</i>	Hoary Willowherb	EDM 1901
446	<i>Epilobium montanum</i>	Broad-leaved Willowherb	DMcC 1958
447	<i>Epilobium lanceolatum</i>	Spear-leaved Willowherb	EDM 1901
448	<i>Epilobium tetragonum</i> ssp. <i>tetragonum</i>	Square-stalked W. herb	CCB 1838
449	<i>Epilobium tetragonum</i> ssp. <i>lamyi</i>	Square-stalked W. herb	BB 1987
450	<i>Epilobium obscurum</i>	Short-fruited Willowherb	DMcC 1958
451	<i>Epilobium ciliatum</i>	American Willowherb	BB 1987
452	<i>Epilobium palustre</i>	Marsh Willowherb	BB 1992
453	<i>Chamerion angustifolium</i>	Rosebay Willowherb	DMcC 1953
454	<i>Oenothera glazioviana</i>	Large-flowered Evening-primrose	JO 1953
454a	<i>Oenothera cambrica</i>	Small-flowered Evening-primrose	ML 2005
455	<i>Oenothera stricta</i>	# * Fragrant Evening-primrose	EDM 1901
456	<i>Fuchsia magellanica</i>	Fuchsia	PHAS 1970

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
c	2 to 12	BEWARE many rabbit-cropped cushiony bushes are <i>U. europeus</i>
?	7 to 9	Most small-spined plants seem to be <i>U. gallii</i> . Needs investigation
vr	2 to 4	2,8. Occasional hedging near sea, probably all planted
vr	6 to 7	13, Deciduous, suckering at Val du Saue. Conf. DG, Kew
o	10 to 12	4,5, Widely planted as salt-resistant hedge. Seedlings often found
r	10 to 12	7,10,13. Self seeds. Lvs. silver spotted above, shining silver below
vlc	5 to 9	4, (Corblet's quarry). Leaves in whorls of 5.
r	5 to 11	4,5,7. Longis, Corblets & Mannez ponds. Naturalised
vlc	6 to 7	4,5,6. Mannez & Corblets qu. Lvs. in whorls of 4.
1site	5 to 8	13. In Val du Saue. Planted at least 30 years ago. Det . by DMcC
1 site	6 to 8	8 c. 200 plants in corner of reseeded field S. side Longis Road
1 site	6 to 7	5. "Tides" headland. Established plant growing from under a rock
f	6 to 10	2,4-9,11,13. Stem to 1.8m. Fls 12mm bright pink/purple, white
o	5 to 10	Wet places. Stem to 75cm. Fls 6mm pale pinkish/purple
o	6 to 9	4,7,8,10. Fls 9mm pink. Lvs rounded at base. Likes shady spots
o	6 to 9	Also occasional garden weed. Lvs narrow elliptic, with petiole
o	6 ro 9	7. Stem sq, Fls 6mm plae purplish/pink. Lvs ±sessile,
r	6 to 9	5,7. As previous ssp but lvs with short petioles
	6 to 8, 11	9,11. Refound Trois Vaux, BB 7.1989
r	6 to 9	4,5,7,8,13. Fls. 5mm pinkish-purple. Lvs rounded at base
r	6 to 8	5,7. Fls 4-5mm pale pink. Lvs narrow at base
o	7 to 9	3,6,7,8,9,10
vlf	6 to 11	3,8,9. Green parts of stems & fruits with red bulbous-based hairs
o	6 to 10	1,4,7. Platte Saline on tipped soil. Sepals green. Red hairs ±
r	5 to 12	1,7,8,9. Petals turn reddish on withering. No red bulbous-based
o	1 to 12	2,5,6,7,8,10. Crabby Bay. Det. TGT & AOC

THE PLANT LIST

No.	Scientific name	English name	Finder/date
457	<i>Circaea lutetiana</i>	Enchanter's-nightshade	EDM 1901
Santalaceæ			
458	<i>Thesium humifusum</i>	Bastard-toadflax	CCB 1838
Celastraceæ			
459	<i>Euonymus japonicus</i>	Evergreen Spindle	FLS 1956
Aquifoliaceæ			
460	<i>Ilex aquifolium</i>	# Holly	BB 1987
Euphorbiaceæ			
461	<i>Mercurialis annua</i>	Annual Mercury	CCB 1838
462	<i>Euphorbia peplis</i> (now extinct)	Purple Spurge (RDBk)	JS 1838
463	<i>Euphorbia helioscopia</i>	Sun Spurge	CCB 1838
464	<i>Euphorbia lathyris</i>	# Caper Spurge	FLS 1977
465	<i>Euphorbia peplis</i>	Petty Spurge	CCB 1838
466	<i>Euphorbia portlandica</i>	Portland Spurge (Scarce)	CCB 1838
467	<i>Euphorbia paralias</i>	Sea Spurge	CCB 1838
468	<i>Euphorbia cyparissias</i>	# Cypress Spurge	BB 1987
Linaceæ			
469	<i>Linum bienne</i>	Pale Flax	CCB 1838
470	<i>Linum usitatissimum</i>	# * Flax	EDM 1902
471	<i>Linum catharticum</i>	Fairy Flax	CCB 1838
472	<i>Radiola linoides</i>	# Allseed	EDM 1901
Polygalaceæ			
473	<i>Polygala vulgaris</i> ssp. <i>vulgaris</i>	Common Milkwort #	EDM 1901
474	<i>Polygala vulgaris</i> ssp. <i>collina</i>	Common Milkwort	CCB 1838
475	<i>Polygala serpyllifolia</i>	Heath Milkwort	DMcC 1957
Hippocastanaceæ			
		Mostly planted	
476	<i>Aesculus hippocastanum</i>	Horse-chestnut	DMcC 1957
477	<i>Aesculus carnea</i>	Red Horse-chestnut	BB 1987
478	<i>Aesculus indica</i>	# Indian Horse-chestnut	BB 1987

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
o	6 to 8	7,8,9,13. Petals 2, white/pale pink. Stamens 2
o	5 to 11	1,4,5,7,8,9. Seems to be increasing again. Lvs & stem olive-green.
f	4 to 6	2,4,5,7,8,9,10. Much planted as hedging, seedlings seen. occasionally
r	3 to 5	6,7,8,13. Self seeds in gardens. also planted by States
c extinct	1 to 12	Dioecious 1,2. Habitat destroyed by storm 1977. Not seen since. Now extinct in BIs.
lf	1 to 12	Annual to 50cm.
r	6 to 7	5,7,8, Probably garden escape. Biennial 1-2m.
c	1 to 12	Annual to 30cm.
c	2 to 10	All round coast. Perennial. Stems usually redish, lvs yellow-green
lf	4 to 9	1,2,3,4. Dune edges. Perennial to 60cm. Lvs narrow, succulent
r	1 to 12	2,7,8. 'Garden Escape. Rhizomatous perennial. To 60cm, lvs linear
lf	3 to 11	2,3,4,5,7,9,13. Short sandy turf. Fls 10mm, pale blue
	5 to 8	Grown as a crop in 16th century. Last record 1902
o	5 to 8	4,5,9,10. Fls 5mm, white
r	5 to 7, 11	5, 7,10,12. Annual. Fls 1mm, petals 8. Capsule 1mm, 8 valves
r	5 to 8	1,7,9,10. Raceme has 10-40 florets. Sepals as wide as capsule
5	5 to 8	5,10,12,13. GCD 1932 As P. dubium. Sepals narrower
o	4 to 7	3,4,5,7,10, 11. Raceme has 3-10 florets
o	3 to 6	8,10. Planted. Specimens to 18m. Churchyard, Valley Gdns
o	4 to 6	8. Planted, Specimens to 9m
1 site	6 to 7	Planted by H.M. Queen in Royal Connaught Square in 1957

THE PLANT LIST

No.	Scientific name	English name	Finder/date
Aceraceæ			
479	<i>Acer platanoides</i>	Norway Maple	CH 1987
480	<i>Acer campestre</i>	Field Maple	BB 1988
481	<i>Acer pseudoplatanus</i>	Sycamore	CCB 1838
Simaroubaceæ			
482	<i>Ailanthus altissima</i>	Tree-of-Heaven	RW 1986
Oxalidaceæ			
483	<i>Oxalis corniculata</i>	Procumbent Yellow-sorrel	FLS 1956
484	<i>Oxalis corniculata</i> var. <i>villosa</i>	#	DMcC 1975
485	<i>Oxalis exilis</i>	Least Yellow Sorrel	CMR 1957
486	<i>Oxalis articulata</i>	Pink-sorrel	CMR 1957
487	<i>Oxalis debilis</i>	# Large-flowered Pink-sorrel	JM 1959
488	<i>Oxalis latifolia</i>	Garden Pink-sorrel	CMR 1954
489	<i>Oxalis latifolia</i> var. <i>cornubiensis</i>	Mexican Oxalis #	JP 1977
490	<i>Oxalis pes-caprae</i>	# Bermuda Buttercup	PHAS 1970
491	<i>Oxalis incarnata</i>	Pale Pink-sorrel	PHAS 1957
Geraniaceæ			
492	<i>Geranium endressii</i>	# French Crane's-bill	BB 1987
493	<i>Geranium x oxonianum</i>	# Druce's Crane's-bill	BB 1994
494	<i>Geranium endressii x oxonianum</i>		BB 1989
495	<i>Geranium versicolor</i>	# Pencilled Geranium	WH 1881
496	<i>Geranium rotundifolium</i>	Round-leaved Geranium	EDM 1901
497	<i>Geranium pratense</i>	# Meadow Crane's-bill	JM 1959
498	<i>Geranium sanguineum</i>	# Bloody Crane's-bill	EDM 1899
499	<i>Geranium columbinum</i>	# Long-stalked Crane's-bill	EDM 1901
500	<i>Geranium dissectum</i>	Cut-leaved Crane's-bill	CCB 1838
501	<i>Geranium submolle</i>	Alderney Crane's-bill	ABJ 1938
502	<i>Geranium x magnificum</i>	# Purple Crane's-bill	BB 1986
503	<i>Geranium pyrenaicum</i>	# Hedgerow Crane's-bill	RW 1977
504	<i>Geranium pusillum</i>	# Small-flowered	JDG 1937
505	<i>Geranium molle</i>	Dove's-foot Crane's-bill	CCB 1838
505	<i>Geranium lucidum</i>	# Shining Crane's-bill	PS 1998

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
r	4 to 6	3,8. Fontaine David to 13.5m. All specimens planted
1 site	5 to 6	1 young specimen in old churchyard, planted
a	4 to 6	Commonest tree in island. Self seeds freely
	6 to 7	8, Planted in gardens, Connaught Square & Coin des Ecailles,
vlf	1 to 12	7,8,12, Garden escape. Old walls in town. Roots at most nodes Only record
o	2 to 12	7,8. Smaller than <i>O. corniculata</i> , only roots at a few nodes
f	1 to 12	'Verges etc. Rhizomatous. Orange dots lf. margins. Occ. white
r	5 to 8	4,7. (As <i>O. corymbosa</i> , Det. by SMW). Bulbous Refound BB 1988
o	5 to 9	7. Bulbous with bulbils on stolons 8
vr	3 to 6	7,8,9. Found in Fontaine David Det. TGT & AOC
r	4 to 7	7,8,9. Along railway track at Newtown. Bulbils in leaf axils
vr	6 to 8	8. Garden escape
r	5 to 8	8
vr	6 to 7	7,8. Determined by PFY 8 Only record
o	3 to 8	3,4,5,7,10. Lvs 5-9 lobes. Gland/hairy. Petals 6mm pink, rounded tip
r	6 to 11	4,9. Large plant at Hammond Memorial. Petals 20 mm, blue or pink
vr	4 to 11	4,12. Petals 20mm bright purplish. Next 1932.
vr	4 to 7	3,4,7,8 Other records CMR 1959 then BB 1989
c	4 to 11	Annual. Lvs 7-lobes, cut almost to base, deep sublobes. fls 5mm pink
vlf	4 to 11	3,4,7,8,9. Peren. to 60cm. Lvs 5-7 deep lobes /sublobes. Fls. 5mm
r	6 to 8	7, Garden escape
vr	4 to 9	2,8. Perennial to 60cm. Lvs 7 deep lobes. Fls c.8mm pinkish-purple
vr	7 to 10	3,4,7,10. Annual to 40cm. Fls 3-4mm. Next 1966 Refound ML 1992
a	1 to 12	Variable annual, 5-40cm. Petals. 5mm, cleft, pink, mauve or whitish
vr	3 to 5	7. Walls nr. Cofil du Nord and at Robin Rock.

THE PLANT LIST

No.	Scientific name	English name	Finder/date
506	<i>Geranium robertianum</i> ssp. <i>robertianum</i>	Herb-Robert	EDM 1901
507	<i>Geranium robertianum</i> ssp. <i>maritimum</i>	Herb-Robert	CCB 1838
508	<i>Geranium purpureum</i> # *	Little Robin (RDBk)	DMcC 1957
509	<i>Erodium maritimum</i>	Sea Stork's-bill	CCB 1838
510	<i>Erodium moschatum</i> *	Musk Stork's-bill Scarce)	CCB 1838
511	<i>Erodium cicutarium</i>	Common Stork's-bill	EDM 1901
512	<i>Erodium cicutarium</i> ssp. <i>dunense</i>	Dune Stork's-bill	NVM 1956
512	<i>Erodium cicut.</i> var. <i>pipinellifolium</i>		EDM 1901
513	<i>Erodium lebellii</i>	Sticky Stork's-bill	PHAS 1970
Tropaeolaceæ			
514	<i>Tropaeolum majus</i>	Nasturtium	DMcC 1958
Araliaceæ			
515	<i>Hedera helix</i> ssp. <i>hibernica</i>	Atlantic Ivy	CCB 1838
516	<i>Aralia elata</i>	Japanese Angelica-tree	BB 1989
Apiaceæ		Formerly Umbelliferae	
517	<i>Hydrocotyle vulgaris</i> #	Marsh Pennywort	CCB 1838
518	<i>Eryngium maritimum</i>	Sea-Holly	CCB 1838
519	<i>Eryngium campestre</i> #	Field Eryngo (RDBk)	NVM 1979
520	<i>Chaerophyllum temulentum</i> #	Rough Chervil	CCB 1838
522	<i>Anthriscus caucalis</i> #	Bur Parsley	CCB 1838
523	<i>Scandix pecten-veneris</i> # *	Shepherd's-needleScarce)	EDM 1901
524	<i>Coriandrum sativum</i> #	Coriander	1977
525	<i>Smyrniolum olusatrum</i> #	Alexanders	EDM 1901
526	<i>Smyrniolum perfoliatum</i> #	(A Southern European sp.)	JO 1953
527	<i>Aegopodium podagraria</i> #	Ground-elder	NJ 1972
528	<i>Berula erecta</i> #	Lesser Water-parsnip	CCB 1838
529	<i>Crithmum maritimum</i>	Rock Samphire	CCB 1838
530	<i>Oenanthe crocata</i>	Hemlock	CCB 1838
531	<i>Aethusa cynapium</i> #	Fool's Parsley	ABJ 1932
532	<i>Foeniculum vulgare</i>	Fennel	CCB 1838
533	<i>Conium maculatum</i>	Hemlock	CCB 1838
534	<i>Bupleurum baldense</i>	Small Hare's-ear	CCB 1838
521	<i>Anthriscus sylvestris</i>	Cow Parsley	EDM 1901

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
c	1 to 12	Annual/biennial; to 50cm. Frequently with red stems and leaf edges
o	3 to 10	1,2,7,11. Procumbent, smaller always nr sea. anthers purple /orange
vr	5 to 9	4. Shingle, cliffs & rocky places nr sea, Anthers yellow
f	3 to 7	Dunes. Lvs simple lower toothed. Petals usually absent
f	3 to 11	4,5, Also occasional garden weed. Lvs pinnate, toothed leaflets.
c	1 to 12	Annual. Fls dark pink (occ with black spot at base upper 2 petals)
f	3 to 12	Sandy bays. Dwarf coastal type. Fls dark pink Now disregarded as a variety
?	4 to 9	Fls pale mauve, pink or white
o	4 to 12	Garden escape
a	9 to 12	Most Alderney plants are probably this ssp.
vr	7 to 8	8,10. Planted, Longis Road and some gardens in town
r	4 to 7	4,11,13. Baxter's Valley, many in dried up Longis pond bed 1992
lf	6 to 9	1,2,3,4
vr		8. below Butes Not seen since
r	3 to 9	5,10,12
vr	6	4,9.
vr	6	Garden escape
l site	3 to 6	8, Not seen 1986-93 Reported by DB 1994, refound 'Butes Hill, plentiful' Only record
No voucher specimen found. The only record		
c	6 to 10	Most shores and lower cliffs Extinct by 1860. But still put onto Mauvaises Herbes list in 1933
vr	8 to 9	8. 1932 'Blaye' Refound in churchyard BB 1992.
la	7 to 12	Dune areas and N. coast
lf	5 to 10	2,4,5,7,9,10,11,12,13,14
vlc	5 to 7	4,5,9,14
c	2 to 12	

THE PLANT LIST

No.	Scientific name	English name	Finder/date
535	<i>Apium graveolens</i> #	Wild Celery	EDM 1902
536	<i>Apium nodiflorum</i> #	Fool's Water-cress	CCB 1838
537	<i>Apium nodiflorum</i> var. <i>pseudorepens</i>	Fool's Water-cress #	CCB 1838
538	<i>Petroselinum crispum</i> #	Garden Parsley	CCB 1838
539	<i>Petroselinum segetum</i> # *	Corn Parsley (RDBk)	BB 1990
540	<i>Falcaria vulgaris</i> #	Longleaf	BHR 1956
541	<i>Pastinaca sativa</i> var. <i>sylvestris</i>	Wild Parsnip	CCB 1838
542	<i>Heracleum sphondylium</i>	Hogweed	CCB 1838
543	<i>Heracleum sphondylium</i> var. <i>angustifolium</i> #	Hogweed	1974
544	<i>Torilis japonica</i> #	Upright Hedge-parsley	CCB 1838
545	<i>Torilis nodosa</i>	Knotted Hedge-parsley	CCB 1838
546	<i>Daucus carota</i> ssp. <i>carota</i>	Wild Carrot	CCB 1838
547	<i>Daucus carota</i> ssp. <i>gummifer</i>	Sea Carrot	CCB 1838
548	<i>Daucus carota</i> ssp. <i>sativa</i> #	Carrot	BB 1995
Gentianaceae			
549	<i>Centaurium erythraea</i>	Common Centaury	CCB 1838
550	<i>Centaurium pulchellum</i> # *	Lesser Centaur (RDBook)	DMcC 1958
Apocynaceae			
551	<i>Vinca major</i>	Greater Periwinkle	EDM 1901
Solanaceae			
552	<i>Nicandra physalodes</i> #	Apple-of-Peru	KEB 1969
553	<i>Lycium barbarum</i>	Duke of Argyll's Teaplant	WH 1881
554	<i>Lycium chinense</i>	Chinese Teaplant	EDM 1908
555	<i>Hyocyanus niger</i> #	Henbane	CCB 1838
556	<i>Lycopersicon esculentum</i> #	Tomato	RSRF 1956
557	<i>Solanum nigrum</i>	Black Nightshade	EDM 1901
558	<i>Solanum nigrum</i> var. <i>atriplicifolium</i> #		ABJ 1932
559	<i>Solanum sarachoides</i> #	Leafy-fruited Nightshade	HCP 1969
560	<i>Solanum dulcamara</i>	Bittersweet	CCB 1838
561	<i>Solanum tuberosum</i> #	Potato	RSRF 1956
562	<i>Datura stramonium</i> #	Thorn-apple	GJ pre 1940
562a	<i>Datura stramonium</i> var. <i>tatula</i> #	Angel's Trumpets	BB 2001

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes	
r	5 to 7	2,4,5	Refound TD 1989
c	6 to 10	1,2,4,5,6,7,11,12,13	
vr	7 to 8	5	Refound BB 1992.
vr	5 to 11	2,4,8.	Garden escape and n.r old ruins 10+ by E Bunker Platte Saline
vr	6 to 8	5.	Path down to Houmet Herbé from road
vr	7 to 9	9,13.	¼ acre on Rond But 1988
lf	5 to 8	4,5,7,8,9.	Pinnate lvs large lflets. Petioles long, wavy. No sepals
a	1 to 12	On Mauvaises Herbes Law list	
vr		10,11.	Very large leaves, by entrance to German tunnel BB
vr		4,10.	Erect 1.2m Fruit spines curved Other records, 1906, 80, 95
o	5 to 8	4,5,9,10,11,	50cm Fls. white. Fruit spines straight. No bracts
f	4 to 8	4,5,6,7,9,10,13.	Umbel concave in fruit
c	1 to 12	Dunes, cliff tops. Umbel convex or flat in fruit	
	6 to 8	13. Relic of cultivation on Blaye. Large patch Only record	
c	6 to 11	Cliffs and commons. Biennial to 50cm	
1 site	5 to 8	5. Annual, usually c.5cm. Damp ground nr. sea	
o	1 to 12	2,3,4,7,8,9,10. Leaf & calyx margins pubescent	
vr	6 to 11	7.	1.10.88 On Platte Saline in quantity after soil tipped
o	4 to 12	5,7,8,9.	Thick clumps along N. coast . Lvs. elliptic. Corolla 7-12mm
o	3 to 12	2-4,6-9.	Leaves lanceolate/narrow ovate, fl. larger. Corolla 12-17mm
vr	6 to 8	2,3,4,7	Recent records. 1 plant; NVM 1979, Berry's Quarry BB 1990
vr	5 to 9	Occasionally on rubbish tips	
lf	6 to 12	3,4,5,7,8,9,10.	Frequent garden weed
		Specimen in BM. Only record	
f	5 to 10		
vr	5 to 9	Rubbish tips .	Refound BB 1988
vr	6 to 9	7,9. Disturbed old ground	
vr	8 to 9	8. By railway station old tipped soil. Persists	

THE PLANT LIST

No.	Scientific name	English name	Finder/date
563	<i>Nicotiana rustica</i>	# Wild Tobacco	OJ 1976
564	<i>Nicotiana glauca</i>	# Sweet Tobacco	BB 1988
Convolvulaceae			
565	<i>Convolvulus arvensis</i>	Field Bindweed	CCB 1838
566	<i>Calystegia soldanella</i>	Sea Bindweed	CCB 1838
567	<i>Calystegia sepium</i>	Hedge Bindweed	EDM 1901
568	<i>Calystegia silvatica</i>	Large Bindweed	DMcC 1953
Cuscutaceae			
569	<i>Cuscuta epithymum</i>	Dodder	CCB 1838
570	<i>Cuscuta campestris</i>	# Yellow Dodder	EDM1901
Menyanthaceae			
571	<i>Nymphoides peltata</i>	#(Scarce) Fringed Water-lily	BB 1986
Hydrophyllaceae			
572	<i>Phacelia tanacetifolia</i>	# Phacelia	BB1994
Boraginaceae			
573	<i>Lithospermum arvense</i>	# Field Gromwell	EDM 1900
574	<i>Echium vulgare</i>	Viper's-bugloss	CCB 1838
575	<i>Echium plantagineum</i>	# Purple Viper's-bugloss	CMR 1956
576	<i>Echium pininana</i>	# Giant Viper's-bugloss	DMcC 1973
577	<i>Symphytum officinale</i>	Common Comfrey	EDM 1901
578	<i>Symphytum x uplandicum</i>	Russian Comfrey	ABJ 1933
579	<i>Symphytum tuberosum</i>	Tuberous Comfrey	CMR 1954
580	<i>Anchusa arvensis</i>	Bugloss	CCB 1838
581	<i>Pentaglottis sempervirens</i>	Green Alkanet	BB 1987
582	<i>Borago officinalis</i>	Borage	CCB 1838
583	<i>Myosotis secunda</i>	Creeping Forget-me-not	CCB 1838
584	<i>Myosotis sylvatica</i>	# Wood Forget-me-not	DMcC 1957
585	<i>Myosotis arvensis</i>	Field Forget-me-not	CCB 1838
586	<i>Myosotis ramosissima</i>	Early Forget-me-not	EDM 1901
587	<i>Myosotis ramosissima</i> ssp. <i>lebelii</i>	Early Forget-me-not	EDM 1901
588	<i>Myosotis discolor</i>	Changing Forget-me-not	EDM 1901
589	<i>Cynoglossum officinale</i>	# Hound's-tongue	CCB 1838

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
		8. Probably garden escape
		9. Probably garden escape
		Only record
c	5 to 11	Found with deep pink fls. nr. Whitegates, occ. pure white
lc	6 to 10	1,2,3,4,5. Mainly N. coast dunes
lc	6 to 11	Bracteoles 10-20mm, don't overlap
vlc	6 to 10	Bracteoles 20-40mm, overlap strongly
lc	6 to 9	Here usually grows on ground on Thyme. Much rarer on Gorse (As <i>C. trifolii</i> , but description agrees with this sp.)
vr	6 to 8	7,10. (Tank on Essex Hill)
vr	6 to 7	12. Remnant of planted fodder crop
vr		2. Only other record 1971. Not seen since
lc	5 to 11	
vr	6 to 7	4,7. Doubtful 1st record, but 5 plants P/S BB 1996 Many T. Rd 2007
r	4 to 9	2,5,7,8,10. Frequently grown in gardens. Occasional escape
vlf	5 to 8	Stem lvs strongly decurrent >1 internode
lf	3 to 9	The more common species. Intermediate hybrids
vr	3 to 8	7,8,12. RSRF 1956. Refound in Churchyard BB 1995
f	2 to 12	4,5,11,12,13. Annual to 50cm. Hispid. Calyx divided nearly to base
o	2 to 8	3,4,5,11. Perennial to 1m. Hispid perennial. Calyx divided ¾ way
o	1 to 12	Well scattered
vlf	5 to 8	7,11. Perennial with stolons. Long pedicels & patent hairs below
	5 to 6	Possibly the only record
o	3 to 9	6,8,13. Plant greyish-green to 40cm. Corolla 3mm.
c	2 to 6	2,3,4,5,9,10,13,14. Calyx with hooked hairs. Fl blue 3mm
o	2 to ?	11,13. White fls. with yellow throat. Refound BB 29.4.1988
lc	3 to 8	1,2,3,4,5,7,9,10,14. Erect, corolla 2mm, yellow first, then blue
		2. Only other record, Crabby CMR 1955

THE PLANT LIST

No.	Scientific name	English name	Finder/date
Verbenaceæ			
590	<i>Verbena officinalis</i>	Vervain	CCB 1838
Lamiaceæ		Formerly Labiataæ	
591	<i>Stachys sylvatica</i>	Hedge Woundwort	EDM 1901
592	<i>Stachys x ambigua</i> #	Hybrid Woundwort	GCD 1932
593	<i>Stachys palustris</i>	Marsh Woundwort	CCB 1838
594	<i>Stachys arvensis</i>	Field Woundwort	CCB 1838
595	<i>Ballota nigra</i> #	Black Horehound	CCB 1838
596	<i>Lamium stramonium</i> ssp. <i>argentatum</i>	Yellow Archangel	JM1959
597	<i>Lamium purpureum</i>	Red Dead-nettle	EDM 1901
598	<i>Lamium hybridum</i>	Cut-leaved Dead-nettle	CCB 1838
599	<i>Lamium amplexicaule</i>	Henbit Dead-nettle	CCB 1838
600	<i>Galeopsis tetrahit</i> #	Common Hemp-nettle	GCD 1932
601	<i>Marrubium vulgare</i> #*(Scarce)	White Horehound	CCB 1838
602	<i>Teucrium scorodonia</i>	Wood Sage	CCB 1838
603	<i>Nepeta x faassenii</i>	Garden Cat-mint	JES 1972
604	<i>Glechoma hederacea</i>	Ground-ivy	CCB 1838
605	<i>Prunella vulgaris</i>	Selfheal	CCB 1838
606	<i>Prunella laciniata</i> #	Cut-leaved Selfheal	RAEB 1932
607	<i>Melissa officinalis</i> #	Balm	DMcC 1956
608	<i>Clinopodium vulgare</i> #	Wild Basil	CRBA 1900
609	<i>Origanum vulgare</i> #	Wild Marjoram	AS 1966
610	<i>Thymus polytrichus</i>	Wild Thyme	CCB 1838
611	<i>Mentha aquatica</i>	Water Mint	CCB 1838
612	<i>Mentha x piperita</i>	Peppermint	EDM 1901
613	<i>Mentha spicata</i> #	Spear Mint	EDM 1901
614	<i>Mentha x villosa</i> ssp. <i>nervata</i>	Sharp-toothed Mint	OJ 1980
615	<i>Mentha x villosa</i>	Apple Mint	DMcC1963
616	<i>Mentha x rotundifolia</i> #	Round-leaved Mint	GCD 1907
617	<i>Mentha longifolia</i> #	Horse Mint	GCD1907
618	<i>Salvia pratensis</i> #	Meadow Clary	JO 1953
619	<i>Salvia verbenaca</i>	Wild Clary	CCB 1838

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
vr	6 to 10	8,10
f	5 to 11	4,7,8,11,12,13. Usually damper spots. Perennial. Leaves petiolate.
?		In GCD 1932 'S'. In WFG Supplement DMcC
o	7 to 9	9,12,13,14. Perennial. Upper & middle stem lvs sessile
o	5 to 10	7,12,13. Annual. No glandular hairs
r	6 to 10	2,9. Waste ground
2 sites	4 to 7	7,8. Refound BB 1993, Churchyard & St. Vignalis' Garden
c	1 to 12	All lvs petiolate, serrate, teeth <2mm
o	1 to 12	5,8,12,13. As above, many teeth >2mm
o	4 to 8	7,8,10. Scattered. Middle & upper lf. subtending whorls sessile In GCD 1932 'S'. Refound JO 1953 Only other record AS1960
r	6 to 11	1,3,4,7,9
a	3 to 10	All round coast and many inland hedges etc.
o	5 to 9	3,4,5,7,8. Probably garden escape
v/c	3 to 9	6,7,9,10,11. Calyx 5 teeth Stolons c. 1m. with petiolate lvs.
f	6 to 9	5,6,7,8,9,10,11,13. Calyx 3 short teeth. 14, Upper lvs divided to midrib Only other record CMR 1951
vr	6 to 11	7,9
vr	7 to 11	5,9. Only other records EDM 1901, GCD "Alderney" 1932, AT '78
vr	7 to 9	7,11 Refound BB 1987. not seen since
c	5 to 11	
c	8 to 11	1,4,5,7,8,12,13. Wet places. Fertile
o	7 to 10	9,10. plant often red-tinged. Sterile
?	6 to 11	8,9. Frequent in gardens. Fertile. Last wild record 1980
vr	8 to 9	8, along railway line. Possibly a garden escape.
	6 to 9	9, Stranger's Cemetery. Sterile Marked 'S' in comital Flora
vr	7 to 9	8, behind Museum. Marked 'S'. Refound BB 1988 10, 'Nunnery' Bracts green/ violet-blue, shorter than fls Only record
lf	5 to 10	1,3,4,5,7,9. Long calyx hairs white. Lvs lobed

THE PLANT LIST

No.	Scientific name	English name	Finder/date
Hippuridaceæ			
620	<i>Hippuris vulgaris</i>	Mare's-tail	EDM 1901
Callitrichaceæ			
621	<i>Callitriche stagnalis</i>	# Common Water-starwort	GCD 1906
622	<i>Callitriche platycarpa</i>	# Various-leaved	CCB 1838
Plantaginaceæ			
623	<i>Plantago coronopus</i>	Buck's-horn Plantain	CCB 1838
624	<i>Plantago maritima</i>	# Sea Plantain	AS 1955
625	<i>Plantago major</i>	Greater Plantain	CCB 1838
626	<i>Plantago lanceolata</i>	Ribwort Plantain	CCB 1838
626a	<i>Plant lanceolata</i> var. <i>sphaerostachys</i>	#	CCB1838
Buddlejaceæ			
627	<i>Buddleja davidii</i>	Butterfly-bush	NVM 1967
628	<i>Buddleja x weyeriana</i>	Hybrid Buddleia	BB 1987
629	<i>Buddleja globosa</i>	Orange-ball-tree	BB 1987
Oleaceæ			
630	<i>Fraxinus excelsior</i>	Ash	CCB 1838
631	<i>Fraxinus angustifolia</i>	# Ash	CH 1987
632	<i>Syringa vulgaris</i>	Lilac	BB 1988
633	<i>Ligustrum vulgare</i>	Wild Privet	EDM 1901
634	<i>Ligustrum ovalifolium</i>	Garden Privet	DMcC 1957
Scrophulariaceæ			
635	<i>Verbascum thapsus</i>	Great Mullein	JDG 1937
636	<i>Verbascum nigrum</i>	Dark Mullein	CCB 1838
637	<i>Scrophularia auriculata</i>	Water Figwort	EDM 1901
638	<i>Antirrhinum majus</i>	# Snapdragon	EDM 1901
639	<i>Misopates orontium</i>	Weasel's-snout	JES 1970
640	<i>Cymbalaria muralis</i>	Ivy-leaved Toadflax	EDM 1901
641	<i>Kickxia elatine</i>	# Sharp-leaved Fluellen	EDM 1901
642	<i>Linaria vulgaris</i>	Common Toadflax	EDM 1901

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
vlf	5 to 10	5. Mannez quarry only. Earlier records, Longis pond 4, Longis pond 7, 'Mill pond'. Only other record EDM 1901
a	4 to 10 7	Very variable in size Also recorded on subsequent visits, 58, 60, 65, doubtful records
f	4 to 11	
a	3 to 11	No Longer considered a separate variety. Only other record GCD 1907
o	4 to 10	3,4,5,8,9,10. Scattered self-sown garden escape.
vr	7 to 10	8,10, Newtown, 10, Bottom Essex valley. Probably planted
r	5 to 8	10. Probably planted, bottom Essex Hill
vlf	4 to 6	7,8,9,10. Self-seeds readily
l site	4 to 5	7. With Brown Buds
vr	4 to 6	9. Rare garden escape
o	5 to 7	4,5,7,9,10,11,12,13. Scattered. Lvs narrow acute, semi-deciduous
r	5 to 10	4,7,8. Frequently planted hedging, some standard trees. Occ. escape
o	5 to 10	2,3,4,5,7,8,9,10
vlc	5 to 8	3,9,10
o	5 to 10	7,8,9,11,12,13. streamsides. Stem angles winged 8
r	7 to 10	2. Garden escape
f	1 to 12	Old walls 9
r	6 to 11	3,5,8,9,11

THE PLANT LIST

No.	Scientific name		English name	Finder/date
643	<i>Linaria alpina</i>	#	Alpine Toadflax	BB 1986
644	<i>Linaria purpurea</i>	#	Purple Toadflax	AT 1983
645	<i>Linaria maroccana</i>	#	Annual Toadflax	DMcC 1973
646	<i>Digitalis purpurea</i>		Foxglove	CCB 1838
647	<i>Veronica serpyllifolia</i>	#	Thyme-leaved Speedwell	CMR 1951
648	<i>Veronica officinalis</i>	#	Heath Speedwell	CCB 1838
649	<i>Veronica chamaedrys</i>		Germander Speedwell	CCB 1838
650	<i>Veronica beccabunga</i>	#	Brooklime	CCB 1838
651	<i>Veronica anagallis-aquatica</i>	#	Blue Water-speedwell	CCB 1838
652	<i>Veronica arvensis</i>		Wall Speedwell	CCB 1838
653	<i>Veronica agrestis</i>	#	Green Field-speedwell	CCB 1838
653a	<i>Veronica filiformis</i>	#	Slender Speedwell	BB 2003
654	<i>Veronica polita</i>	#	Grey Field-speedwell	CCB 1838
655	<i>Veronica persica</i>		Common Field-speedwell	EDM 1901
656	<i>Veronica hederifolia</i>		Ivy-leaved Speedwell	CCB 1838
657	<i>Veronica longifolia</i>		Garden Speedwell	BB 1987
658	<i>Hebe salicifolia</i>		Koromiko	DMcC 1963
659	<i>Hebe x franciscana</i>		Hedge Veronica	DMcC 1963
660	<i>Euphrasia</i> agg.		Eyebrights	CCB 1838
661	<i>Euphrasia rostkoviana</i>	#	An Eyebright	CCB 1838
662	<i>Euphrasia tetraqueta</i>		An Eyebright	EDM 1900
663	<i>Euphrasia nemorosa</i>		An Eyebright	EDM 1900
664	<i>Euphrasia confusa</i>		An Eyebright	DMcC 1963
665	<i>Euphrasia stricta</i>	#	An Eyebright	EDM 1901
666	<i>Euphrasia ostenfeldii</i>	#	An Eyebright	GCD 1906
667	<i>Odontites vernus</i>	#	Red Bartsia	CCB 1838
668	<i>Parentucellia viscosa</i>		Yellow Bartsia	CCB 1838
669	<i>Pedicularis sylvatica</i>		Lousewort	EDM 1901

Orobanchaceae

670	<i>Orobanche purpurea</i>		Yarrow Broomrape	CCB 1838
671	<i>Orobanche rapum-genistae</i>	*	Greater Broomrape (Scarce)	CCB 1838
672	<i>Orobanche hederae</i>	*	Ivy Broomrape (Scarce)	EDM 1901
673	<i>Orobanche minor</i>		Common Broomrape	CCB 1838
674	<i>Orobanche minor</i> var. <i>maritima</i>	*	Carrot Broomrape (RDBk)	EDM 1901

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
vr	7 to 8	2. Garden escape
r	5 to 12	7,8,9
		Only record
lc	5 to 9	Mainly on cliffs
r	3 to 6	7. Petite Blaye. Perennial, roots at nodes. Fls white/pale blue, 6-8mm
r	4 to 8	7,8,11. Fls. lilac, 5-9mm
f	3 to 9	Fls. bright blue 8-12mm
r	5 to 9	7,9. Wet places, fls, bright blue
	5 to 9	7,10,11, Trois Vaux. Fls. pale blue, dk veins.
f	3 to 7	4,5,7,8,9,11,12. Common garden weed. Fls blue 2-3mm
vr		Fls white/pale blue, axillary, solitary, 3-8mm. Lvs longer than wide
r	3 to 5	7. In lawn at The Twins. 1st island record 27.4.03
vr		As above but lvs wider than long
c	4 to 12	Fls bright blue lower lobe white, 8-12mm
c	1 to 11	Walls, verges, garden weed. Fls solitary in lf axil 2-9mm
vr	7 to 9	Garden escape. Fls in dense terminal raceme
vr	3 to 12	
f	1 to 12	2,3,5,7,8,13. Much used as hedging. Self seeds
lc	4 to 10	
		Only other record EDM 1901
		4,10. Euphrasia aggregate common in short turf all round
lc	4 to 10	4. Right to the sea edge. Have not been differentiated Determined by PFY
		Appears to be the only record
		(4, As Euphrasia curta var, glabrescens)
vr	6 to 8	4,8. Two single patches in verges for many years until 1993
lf	6 to 9	Quarries & E. coast. Damp valleys
o	4 to 7	11,12,13. Moist places above cliffs. Baxter's Valley
c	4 to 7	Even in lawns.
vlc	5 to 7	11,12,13. S. cliffs on Cytisus & Ulex
o	6 to 7	8. Confined to Connaught Sq & Le Vallée
c	4 to 8	Many hosts.
vlf	4 to 8	3,7. On Daucus, Plantago, Glaucium & Ononis

THE PLANT LIST

No.	Scientific name		English name	Finder/date
Acanthaceæ				
674a	<i>Acanthus mollis</i>	#	Bear's-breech	BB 1998
Campanulaceæ				
675	<i>Campanula portenschlagiana</i>		Adria Bellflower	BB 1987
676	<i>Campanula poscharskyana</i>		Trailing Bellflower	BB 1988
677	<i>Jasione montana</i>		Sheep's-bit	CCB 1838
Rubiaceæ				
678	<i>Sherardia arvensis</i>		Field Madder	CCB 1838
679	<i>Asperula cynanchica</i>	#	Squinancywort	EDM 1901
680	<i>Galium palustre</i>		Common	CCB 1838
681	<i>Galium verum</i>		Lady's Bedstraw	CCB 1838
682	<i>Galium mollugo</i> ssp. <i>mollugo</i>		Hedge Bedstraw	CCB 1838
683	<i>Galium mollugo</i> ssp. <i>erectum</i>		Hedge Bedstraw	ABJ 1932
684	<i>Galium saxatile</i>		Heath Bedstraw	EDM 1901
685	<i>Galium aparine</i>		Cleavers	CCB 1838
686	<i>Cruciata laevipes</i>	#	Crosswort	DMcC 1962
687	<i>Rubia peregrina</i>		Wild Madder	CCB 1838
Caprifoliaceæ				
688	<i>Sambucus nigra</i>		Elder	CCB 1838
689	<i>Lonicera periclymenum</i>		Honeysuckle	CCB 1838
690	<i>Lonicera periclym</i> var. <i>quercifolia</i>		Oak-leaved Honeysuckle	BB 1988
Valerianaceæ				
691	<i>Valerianella locusta</i>		Common Cornsalad	EDM 1901
692	<i>Valerianella carinata</i>		Keeled-fruited Cornsalad	CCB 1838
693	<i>Valerianella dentata</i>	#	Narrow-fruited	EDM 1900
694	<i>Valerianella eriocarpa</i>		Hairy-fruited Cornsalad	EDM 1901
695	<i>Centranthus rubra</i>		Red Valerian	EDM 1901

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
vr	6 to 7	11. Single plant on Giffoine along circular track to guns
l	1 to 12	8. On old walls. Corolla funnel-shaped
vlf	1 to 11	8. Old walls. Corolla star-shaped
c	4 to 12	Always near sea
a	3 to 10	Dry banks, cliffs, commons. White or pale 3,5. Only other record JCNW 1956
o	5 to 7	4,6,7,12,13. Perennial. Rough. Pondsides/ditches. Fls white
la	6 to 9	Perennial. Short turf. One old wall in town. Fls yellow
r	5 to 8	5,7,10,11,12,13,14. Perennial. Smooth, branches at >45°. Fls white
vr	6 to 8	5,7,13. On more calcareous soils. Branches <45°BB 1995
o	6 to 7	5,6,9,11,12,14. Acid soils. Perennial. Lvs .prickles on margins
c	4 to 8	Strong prickles on stems. fruits with hooked bristles.
vr	5 to 8	4,5,12,13. Evergreen. Prickles recurved. Fls pale yellow-green
f	4 to 10	Scattered all over
a	5 to 12	Colour varies from almost white to dark pink
r	6 to 9	Leaves oak-leaf shaped
vr	3 to 7	7,12 Calyx absent from fruit. Fr.just longer than wide
c	3 to 11	1,2,3,5,8. Walls & garden weed. Fruit much longer than wide
vr	6	8,9. Calyx distinct in fruit 2 teeth
vr		3,9,10. Calyx 6 teeth. Refound RR 1991, previous record JO 1953
c	1 to 12	Principally on old walls & quarry spoil. Red, white or pink forms

THE PLANT LIST

No.	Scientific name	English name	Finder/date
Dipsaceae			
696	<i>Dipsacus fullonum</i>	# Wild Teasel	CMR 1951
697	<i>Dipsacus sativus</i>	# Fuller's Teasel	DMcC 1958
698	<i>Knautia arvensis</i>	# Field Scabious	EDM 1901
Asteraceae		Formerly Compositae	
699	<i>Carlina vulgaris</i>	Carlina Thistle	CCB 1838
700	<i>Arctium minus</i> ssp. <i>minus</i>	Lesser Burdock	EDM 1901
701	<i>Arctium minus</i> ssp. <i>nemorosum</i>	#	EDM 1901
702	<i>Arctium lappa</i>	# Greater Burdock	CCB 1838
703	<i>Carduus tenuiflorus</i>	Slender Thistle	CCB 1838
704	<i>Carduus nutans</i>	Musk Thistle	CCB 1838
705	<i>Cirsium vulgare</i>	Spear Thistle	CCB 1838
706	<i>Cirsium acaule</i>	Dwarf Thistle	EDM 1901
707	<i>Cirsium palustre</i>	Marsh Thistle	EDM 1901
708	<i>Cirsium arvense</i>	Creeping Thistle	CCB 1838
709	<i>Onopordum acanthium</i>	Cotton Thistle	BB 1991
710	<i>Centaurea scabiosa</i>	Greater Knapweed	CCB 1838
711	<i>Centaurea cyanus</i>	# * Cornflower (Scarce)	CCB 1838
712	<i>Centaurea aspera</i>	Rough Star-thistle	EDM 1901
713	<i>Centaurea x moncktonii</i>	A hybrid Knapweed	ABJ 1932
714	<i>Centaurea nemoralis x jacea</i>	# A hybrid Knapweed	DMcC 1958
715	<i>Centaurea nigra</i> ssp. <i>nigra</i>	Common Knapweed	EDM 1901
716	<i>Centaurea nigra</i> ssp. <i>nemoralis</i>	Common Knapweed	EDM 1901
717	<i>Centaurea cineraria</i>	# (A western Mediterranean sp.)	MK 1995
718	<i>Cichorium intybus</i>	Chicory	EDM 1901
719	<i>Lapsana communis</i>	Nipplewort	EDM 1901
720	<i>Hypochaeris radicata</i>	Cat's-ear	CCB 1838
721	<i>Hypochaeris glabra</i>	* Smooth Cat's-ear(Scarce)	CCB 1838
722	<i>Leontodon autumnalis</i>	Autumn Hawkbit	CCB 1838
723	<i>Leontodon hispidus</i>	Rough Hawkbit	CCB 1838
724	<i>Leontodon saxatilis</i>	Lesser Hawkbit	CCB 1838
725	<i>Picris echioides</i>	Bristly Oxtongue	EDM 1901
726	<i>Picris hieracoides</i>	Hawkweed Oxtongue	EDM 1901

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
vr	7 to 8	2,7,8,13. Bracts on receptacle stiff flexible, straight, apical spine
	7 to 8	Apical spine rigid & recurved. Only record
r	7 to 9	3,9,10
o	6 to 10	3,4,5
o	6 to 10	Scattered individuals in most squares. Lower leaf stalks hollow 2. As above, but peduncles much shorter, 0-1cm Last record 1954 Leaf stalks solid Possibly an error for the above.
c	4 to 8	1,2,3,4,5,7,11
lf	6 to 10	2,4,5,7,9,10
c	5 to 10	All <i>Cirsium</i> spp are on Mauvaises Herbes list
vlf	7 to 10	3,4,5. Usually stemless, occasionally 8 -10cm. Rarely with white fls.
o	5 to 11	3,4,5,7,9,11,12. Wet places, usually in small numbers
c	5 to 11	Also common garden weed
vr	6 to 8	3,8,9
o	6 to 11	3,4,8,9,12,13,14 8,12 Only other records EDM 1901, CMR 1955
r	5 to 8	3, since 1890's in same spot. (Claimed Platte Saline FF 1967) 3,4,5,11,12,14. CMR 1950 determined by WBT 1958. Determined by WBT 1958 Only other record BB 1992
vr		
f	6 to 10	3,4,5,12,14
f	6 to 10	11
1 site	6 to 10	8. Base of Water Tower. Determined by ACL
vr	6 to 10	4,5,7, 8,9,11
vr	6 to 7	8,9
c	4 to 1	
lf	4 to 2	2,3,4,5,7. Sandy soils mostly near coast
c	1 to 12	Outermost ligules red on underside
vr	6 to 8	9. Stace says "Not in CI". Refound by BSBI 1995.
c	1 to 12	All round coast. Outermost ligules grey-violet
c	5 to 1	Common on Sandy turf & verges
f	5 to 12	

THE PLANT LIST

No.	Scientific name	English name	Finder/date
727	<i>Picris hieracoides</i> var. <i>incana</i>	Druce's Oxtongue	GCD 1906
728	<i>Tragopogon pratensis</i> ssp. <i>minor</i>	Goat's-beard	EDM 1901
729	<i>Tragopogon porrifolius</i> #	Salsify	EDM 1901
730	<i>Sonchus arvensis</i>	Perennial Sow-thistle	CCB 1838
731	<i>Sonchus oleraceus</i>	Smooth Sow-thistle	CCB 1838
732	<i>Sonchus asper</i>	Prickly Sow-thistle	EDM 1901
733	<i>Lactuca serriola</i>	Prickly Lettuce	BB1987
734	<i>Taraxacum</i> spp. agg.	Dandelions	CCB 1838
	Sect. 1. <i>Erythrosperma</i>	Lesser Dandelions	EDM 1901
	Sect. 2. <i>Obliqua</i>		
	Sect. 3. <i>Palustria</i>		
	Sect. 4. <i>Spectabilia</i>	Red-veined Dandelions	
	Sect. 7. <i>Celtica</i>		
	Sect. 8. <i>Hamata</i>		
	Sect. 9. <i>Ruderalia</i>	Common Dandelions	CCB 1838
739	<i>Taraxacum pseudohamatum</i> #	Sect. 8.	RC 1986
740	<i>Taraxacum lamprophyllum</i> *	Sect. 8.	RC 1986
741	<i>Taraxacum bracteatum</i> #	Sect. 8.	RC 1986
742	<i>Taraxacum boekmanii</i> #	Sect. 8.	RC 1986
743	<i>Taraxacum expallidiforme</i> #	Sect. 9.	RC 1986
744	<i>Taraxacum ekmanii</i> #	Sect. 9.	RC 1986
745	<i>Taraxacum cordatum</i> #	Sect. 9.	RC 1986
746	<i>Taraxacum ancistrolobum</i> #	Sect. 9.	RC 1986
747	<i>Taraxacum insigne</i>	Sect. 9.	RC 1986
748	<i>Taraxacum polyodon</i> #	Sect. 9.	RC 1986
749	<i>Taraxacum dahlstedtii</i>	Sect. 9.	MMcW
750	<i>Taraxacum uncosum</i>		DMcC 1984
751	<i>Taraxacum crispifolium</i> #		AB 1974
752	<i>Taraxacum privum</i> #		AB 1974
753	<i>Taraxacum simile</i> #		AB 1974
754	<i>Crepis biennis</i> #	Rough Hawk's-beard	CCB 1838
755	<i>Crepis capillaris</i>	Smooth Hawk's-beard	CCB 1838
756	<i>Crepis vesicaria</i> ssp. <i>taraxacifolia</i>	Beaked Hawk's-beard	FJH 1884
757	<i>Pilosella peleteriana</i> ssp.	Shaggy	CCB 1838

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
r	7 to 9	2.4 Refound BB 1988
r	3 to 11	3,7,8,9. Yellow flowers 2. Purple flowers Last records 1907, 1954
r	1, 6 to 11	4,9,11,12
c	1 to 12	
f	3 to 9	
r	7 to 10	7,8.. Walls & gutters,
overall c	1 to 12	Have not been fully determined, but at least 19 ssp. recorded
		Habitat; open sandy turf by the sea
r		Habitat; wet, base-rich meadows, marshy places
c		Habitat; damp or wet acidic grassy places and roadsides
		Habitat; wet places in lowland grassland & rocky ledges
		Habitat; grassland, damp & dry, roadsides
c		Habitat; Common weeds, the largest group
		Det . AJR/CCH. Butes Hill
		Det. AJR/CCH. Windy corner
		Det. AJR/CCH. Longis Road verge
		DEt. AJR/CCH. New Churchyard
		Det. AJR/CCH. Grass bank Les Butes
		Det. AJR/CCH. Gravestone edge, New Cemetery
		Det AJR/CCH. Roadside nr. Sea View Hotel
		Det. AJR/CCH. Allée ès Fées
		Det. C.C. Howarth. Rubbish tip La Tchue.
		Det. CCH. Ladysmith
f		Det. van Soerst. Conf AJR 1968 from Herb. Cantab. specimen
		Det. AJR. 3ssp. noted in WFG Supplement
		Det. AJR. 1st CI record
		Det AJR. 1st. CI record
		Det AJR
c	5 to 12	4-14. Stem lvs not clasping (var. griseola recorded JDG 1937)
f	4 to 11	3-14. Stem lvs clasping at base
o	5 to 9	8. Short grass, dunes

THE PLANT LIST

No.	Scientific name	English name	Finder/date
758	<i>Pilosella officinarum</i>	Mouse-ear-hawkweed	EDM 1901
759	<i>Hieracium umbellatum</i>	Umbellate Hawkweed	EDM 1901
760	<i>Filago vulgaris</i>	Common Cudweed	CCB 1838
761	<i>Filago minima</i>	Small Cudweed	CCB 1838
762	<i>Gnaphalium uliginosum</i>	Marsh Cudweed	EDM 1900
763	<i>Gnaphalium luteo-album</i>	Jersey Cudweed	ML 1974
764	<i>Gnaphalium undulatum</i>	Cape Cudweed	BHR 1956
765	<i>Helichrysum petiolare</i>	Everlasting Flower	BB 1993
765a	<i>Helichrysum angustifolium</i>	Curry Plant	BB/JP 2007
766	<i>Inula conyza</i>	Ploughman's-spikenard	EDM 1903
767	<i>Inula crithmoides</i> *	Golden-samphire (Scarce)	CCB 1838
768	<i>Pulicaria dysenterica</i>	Common Fleabane	CCB 1838
769	<i>Solidago canadensis</i>	Canadian Goldenrod	RR 1963
770	<i>Aster tripolium</i>	Sea Aster	WF 1956
771	<i>Erigeron glaucus</i>	Seaside Daisy	BB 1987
772	<i>Erigeron karvinskianus</i> #	Mexican Fleabane	ML 1970
773	<i>Erigeron acer</i> #	Blue Fleabane	CCB 1838
774	<i>Conyza canadensis</i>	Canadian Fleabane	CMR 1950
774a	<i>Conyza sumatrensis</i>	Guernsey Fleabane	ML 2000
775a	<i>Conyza bilbaoana</i>	Hispid Fleabane	BB2007
775	<i>Olearia paniculata</i>	Akirabo	BB 1987
776	<i>Olearia macrodonta</i>	New Zealand Holly	BB 1987
777	<i>Olearia traversii</i>	Ake-ake	BB 1987
778	<i>Bellis perennis</i>	Daisy	CCB 1838
779	<i>Tanacetum parthenium</i>	Feverfew	DMcC 1956
780	<i>Tanacetum vulgare</i>	Tansy	ABJ 1932
781	<i>Artemesia vulgaris</i>	Mugwort	EDM 1901
782	<i>Artemesia absinthum</i>	Wormwood	CCB 1838
783	<i>Otanthus maritimus</i> # *	Cottonweed (RDBk)	CCB 1838
784	<i>Achillea ptarmica</i>	Sneezewort	VKM 1956
785	<i>Achillea millefolium</i>	Yarrow	CCB 1838
786	<i>Chamaemelum nobile</i> # *	Chamomile (Scarce)	CCB 1838
787	<i>Anthemis arvensis</i> # *	Corn Chamomile	CCB 1838
788	<i>Anthemis cotula</i> # *	Stinking Chamomile	EDM 1899
789	<i>Chrysanthemum segetum</i> #	Corn Marigold	CCB 1838
790	<i>Leucanthemum vulgare</i>	Oxeye Daisy	CCB 1838

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
f	5 to 10	Ligules red-striped beneath Determined by CW
o	6 to 10	4,8,10,13. Leaves all sessile
r	7 to 9	3,4,5,8,10,13. Stems to 40cm. fl clusters 10-12mm
o	6 to 9	5,10,13. Stems to 20cm. fl clusters 2-5mm
r	7 to 11	1,4,5,7,8. Flowering clusters leafy. Phyllaries pale brown, dk. tips
vr	7 to 1	2,7,8. Leaves grey, woolly. Phyllaries straw-coloured
lf	6 to 10	3,4,5,7,8,9,11,13. Lvs green on underside. Phyllaries whitish
r	6 to 8	10. Well established Bluestone Hill, seeding in verge. Dozens
vr	5 to 8	10. 2 plants in verge on S facing bank Bluestone Hill. Garden escape
r	6 to 9	4,14. Fl heads 1cm, fls yellowish , outer bracts purple. Scrub etc.
o	6 to 9	2,3,4,6,10,11,12. Fl heads 2.5cm, golden-yellow. Nr. or on old forts
f	4 to 10	Wet or damp places
1 site	7 to 10	6. Probably planted many years ago 5. Single colony rock ledges nr. Houmet Herbé. Refound BB 1990 White fl.
o	1 to 12	2,4,7,8,10. Walls & banks. Fls single 3-4cm, ray florets lilac
vr	3 to 11	7, 8 Old walls. fls in loose cluster, ray florets white, purple below
vr	5 to 7	8,9,10. Annual/biennial. Fl heads 12-18mm, ray florets pale purple
lf	1 to 12	2,3,4,5,6,7,8,9,10. Bare sandy places
a	8 to 11	1,2,4,7,8,9,10. Many plants at each site. Previously overlooked
f	8 to 11	mixed with above sp. Confirmed by ML 8-9/07
vr	8 to 10	5,8. Planted hedging. Occasional seedlings, inconspicuous flowers
o	6 to 10	8,9. Planted. Churchyard and gardens. Showy white fls.
o	6 to 8	5,8. Hedging. Plant on E. coast to 7m. Fls greyish axillary pyramid
a	1 to 12	
r	6 to 9	5,8. Garden escape. Churchyard & Museum
vlf	4 to 9	2,3,5,7,8. First record, 'near Mannez Cottages'
o	6 to 10	3,4,6-10,12,14. Waste places. Lvs green above, reddish, angled stem
o	5 to 11	1,2,3,4,5,7,10. Often with 718. Lvs. silvery both sides 3 Only other record 1896, gone by 1899
vr	6 to 12	2,7,9. Garden escape
a	3 to 12	
lf	5 to 8	6,11,12. (Cliffs) Only other record. FF 1967 Trois Vaux 10 Only confirmed record
vr	5 to 9	11,12. Formerly abundant in cultivated fields. Now rare
la	2 to 11	Cliffs

THE PLANT LIST

No.	Scientific name	English name	Finder/date
791	<i>Leucanthemum x superbum</i> #	Shasta Daisy	DMcC 1961
792	<i>Matricaria recutita</i> #	Scented Mayweed	CMR 1956
793	<i>Matricaria discoidea</i>	Pineapple-weed	CMR 1954
794	<i>Tripleurospermum maritimum</i>	Sea Mayweed	CMR 1954
795	<i>Tripleurospermum inodorum</i>	Scentless Mayweed	CCB 1838
796	<i>Senecio cineraria</i>	Silver Ragwort	JP 1977
797	<i>Senecio jacobea</i>	Ragwort	CCB 1838
798	<i>Senecio vulgaris</i>	Groundsel	CCB 1838
799	<i>Senecio sylvaticus</i>	Heath Groundsel	CCB 1838
800	<i>Tussilago farfara</i> #	Colt's-foot	JDG 1937
801	<i>Petasites fragrans</i>	Winter Heliotrope	EDM 1901
802	<i>Calendula officinalis</i>	Pot Marigold	DMcC 1957
803	<i>Helianthus annuus</i>	Sunflower	BB 1988
804	<i>Helianthus tuberosus</i> #	Jerusalem Artichoke	BB 1989
805	<i>Eupatorium cannabinum</i> #	Hemp Agrimony	JP 1977
806	<i>Cynara scolymus</i> #	Globe Artichoke	PHAS 1970

2. Liliidæ

Hydrocharitaceæ

807	<i>Elodea canadensis</i> #	Canadian Waterweed	BB 1987
808	<i>Lagarosiphon major</i>	Curly Waterweed	BHR 1956

Juncaginaceæ

809	<i>Triglochin palustre</i> #	Marsh Arrowgrass	CCB 1838
-----	------------------------------	------------------	----------

Potamogetonaceæ

810	<i>Potamogeton natans</i>	Broad-leaved Pondweed	DMcC 1963
811	<i>Potamogeton berchtoldii</i> #	Small Pondweed	JES 1972

Zosteraceæ

812	<i>Zostera marina</i> *	Eelgrass (Scarce)	EDM 1901
-----	-------------------------	-------------------	----------

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
vr	1 to 12	Frequent in gardens, rare escape Only record
vr	3 to 11	Very strongly sweet-scented. Phyllaries pale brown
c	4 to 1	Bare places, sandy tracks. Pineapple scent. Phyllar, white
c	3 to 12	Stems prostrate/decumbent. Strand line & above all round coast
c	1 to 12	6,11-14. Cultivated & waste ground. Leaf segments acute with bristle tips
r	6 to 9	1,2,3,4. Braye Beach. Probably a garden escape
a	4 to 1	On Mauvaises Herbes list
f	1 to 12	
lc	4 to 11	1,2,4,9. Mainly on cliffs & in old quarries. Erect, larger than above 5,14 Only other record AS 1958
la	12 to 4	3,7,8
f	1 to 12	2,3,7,8. Frequent garden escape
r	8 to 10	7,9,12. Garden escape or remnants of planted seed
vr	8 to 10	7, Hedge in Le Petit Val Only record
vr	6 to 9	4 Near Saye Farm. Only record in the wild
r	7 to 10	7,9. Mainly relic of cultivated crops in 1950's. Now spreading

(Formerly Monocotyledonidæ)

vr	5 to 10	4. Dioecious. Lvs in whorls of 3, apices obtuse
vla	?	Huge plants in deep water, Corblets quarry . BB

12 **Possibly the only record**

These have not been fully investigated

vla	5 to 9	4,5. Floaing lvs ovate, knee joint at jct petiole & stem
1 mainsite	5 to 9	4. Det. JED. Refound ML 1988. a in 1995. Lvs linear all submerged

lf	6 to 9	4,5,10,13,14, all below HWM, plentiful at Longis & Saye and below HWM in several other places
----	--------	---

THE PLANT LIST

No.	Scientific name	English name	Finder/date
Araceæ			
813	<i>Zantedeschia aethiopica</i>	# Altar-lily	BB 1990
814	<i>Arum maculatum</i>	# Lord's-and-Ladies	CMR 1954
815	<i>Arum italicum</i> ssp. <i>italicum</i>	Italian Lord's-and-Ladies	ABJ 1934
816	<i>Arum italicum</i> ssp. <i>neglectum</i>	Italian Lord's-and-Ladies	ABJ 1934
817	<i>Dracunculus vulgaris</i>	# Dragon Arum	CMR 1960
		* (Both <i>A.Italicum</i> RDBk)	
Lemnaceæ			
818	<i>Lemna minor</i>	Common Duckweed	CCB 1838
819	<i>Lemna trisulca</i>	# Ivy-leaved Duckweed	DMcC 1964
820	<i>Lemna minuta</i>	Least Duckweed	BB 1993
Juncaceae			
821	<i>Juncus gerardii</i>	# Saltmarsh Rush	FWS 1958
822	<i>Juncus bufonius</i>	Toad Rush	CCB 1838
823	<i>Juncus capitatus</i>	# * Dwarf Rush (RDBk)	ABJ 1933
824	<i>Juncus articulatus</i>	Jointed Rush	CCB 1838
825	<i>Juncus acutiflorus</i>	Sharp-flowered Rush	EDM 1901
826	<i>Juncus bulbosus</i>	# Bulbous Rush	EDM 1899
827	<i>Juncus maritimus</i>	Sea Rush	CCB 1838
828	<i>Juncus acutus</i>	# * Sharp Rush (Scarce)	CCB 1838
829	<i>Juncus inflexus</i>	Hard Rush	CCB 1838
830	<i>Juncus effusus</i>	Soft Rush	EDM 1901
831	<i>Juncus conglomeratus</i>	# Compact Rush	CCB 1838
832	<i>Luzula campestris</i>	Field Woodrush	CCB 1838
833	<i>Luzula multiflora</i>	Heath Woodrush	CCB 1848
Cyperaceæ			
834	<i>Eleocharis palustris</i>	Common Spike-rush	CCB 1838
835	<i>Eleocharis multicaulis</i>	# Many-stalked Spike-rush	CCB 1838
836	<i>Eleocharis quinqueflora</i>	# Few-flowered Spike-rush	DMcC 1963
837	<i>Bolboschoenus maritimus</i>	# Sea Club-rush	EDM 1901
838	<i>Schoenoplectus tabernaemontani</i>	# Grey Club-rush	EDM 1901
839	<i>Isolepis setacea</i>	Bristle Club-rush	EDM 1901

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
vr	1 to 8	8. Planted and escapes downstream
vr	5	8. The Terrace 1 plant. Lvs/spathe spotted purple. Spike 3-4cm
lf	2 to 6	3,7-10,13. White veins on leaves Mrs. Rae 1976 record as "Pictum"
vlf	2 to 6	8,9,10,13. No white veins. Fruiting spike 10-15cm.
vr	5 to 8	8.9. Occasional spectacular "visitor". 4 or 5 sites
lf	6 to 7	6,7,8,11,12,13. Wet places and old water troughs
2 sites	5 to 7	Abundant in Mannez quarry in some years. Longis Pond BB 1996
vla	6 to 7	1,7. Clonque Quarry 1993. Abundant Platte Saline pond

Alderney's Rush species are fairly limited

Lowest bract shorter than infl. Refound FWS 1958

c	5 to 9	2,4,5,7,8,11,12,13 etc. Damp habitats
vr	4 to 6	6,10,11,14. Stem to 5cm. Habitat wet in winter
o	5 to 9	2,4,5,7,11,12,13. Feel the septa in the stem
f	7 to 9	4. The latest flowering rush
vr	5 to 7	6,7. Amphibious Other records 1934, 1987
vla	7 to 8	4,5,6,10. Abundant in Berry's Quarry. Lowest bract exceeds infl.
o	5 to 9	5. (round Mannez pond). Stem very rigid
lc	6 to 8	4,5,12. Abundant in Mannez quarry. Stem ridged, interrupted pith
f	6 to 10	4,5,7,11,13. Esp, round Mannez pond. Stem smooth, glossy
r	6 to 8	7 (Bonne Terre),11. Stems ridged, dull
a	3 to 6	Fls in 1 sessile & 2 stalked clusters 3-12 in each
o	4 to 7	4. Fls in several stalked clusters, 8-18 in each

The Alderney Sedges need further investigation

vla	5 to 6, 10	5,10,11,13. Upper stem lf-sheath truncated
r	6 to 7	5,7,11. Upper stem lf-sheath oblique at apex
r	6 to 7	As distinct from above 2 spp, lowest glume completely encircles base
r	5 to 7	Burhou only Refound FWS 1958, BB 1989
?		4 Possibly the only record
vlf	5 to 8	5,7,11,12. Bract much longer than inflorescence

THE PLANT LIST

No.	Scientific name	English name	Finder/date
840	<i>Isolepis cernua</i>	# Slender Club-rush	CCB 1838
841	<i>Cyperus longus</i>	* Galingale (Scarce)	CCB 1838
842	<i>Schoenus nigricans</i>	Black Bog-rush	CCB 1838
843	<i>Carex paniculata</i>	Greater Tussock-sedge	EDM 1901
844	<i>Carex muricata</i> ssp. <i>muricata</i>	Prickly Sedge	CCB 1838
845	<i>Carex divulsa</i> ssp. <i>divulsa</i>	Grey Sedge	EDM 1901
846	<i>Carex divulsa</i> ssp. <i>leersii</i>	Tall Spiked-sedge	ABJ 1933
847	<i>Carex arenaria</i>	Sand Sedge	CCB 1838
848	<i>Carex disticha</i>	Brown Sedge	ML 1996
849	<i>Carex ovalis</i>	# Oval Sedge	ABJ 1932
850	<i>Carex echinata</i>	# Star Sedge	EDM 1901
851	<i>Carex lachenalii</i>	# Hare's-foot Sedge	ABJ 1932
852	<i>Carex hirta</i>	Hairy Sedge	EDM 1901
853	<i>Carex pseudocyperus</i>	Cyperus Sedge	BB 1992
854	<i>Carex pendula</i>	Pendulous Sedge	DMcC 1963
855	<i>Carex flacca</i>	Glaucous Sedge	CCB 1838
856	<i>Carex panicea</i>	Carnation Sedge	CCB 1838
857	<i>Carex distans</i>	Distant Sedge	ABJ 1933
858	<i>Carex extensa</i>	Long-bracted Sedge	EDM 1901
859	<i>Carex flava</i>	# Large Yellow-sedge	CCB 1838
860	<i>Carex viridula</i>	Yellow-sedge	JO 1953
861	<i>Carex caryophylla</i>	Spring Sedge	CCB 1838
862	<i>Carex pulicaris</i>	Flea Sedge	CCB 1838

Poaceæ

Subfamily Bambusoideæ

863	<i>Yushaniana anceps</i>	Indian Fountain-bamboo	AHL 1968
864	<i>Pseudosasa japonica</i>	Arrow Bamboo	AHL 1968

Subfamily Pooideæ

865	<i>Nardus stricta</i>	# Mat-grass	AS 1960
866	<i>Festuca pratensis</i>	Meadow Fescue	CCB 1838
867	<i>Festuca arundinacea</i>	Tall Fescue	EDM 1901
868	<i>Festuca arenaria</i>	# Rush-leaved Fescue	ABJ 1933
869	<i>Festuca rubra</i> ssp. <i>rubra</i>	Red Fescue	CCB 1838
870	<i>Festuca rubra</i> ssp. <i>junceæ</i>	Red Fescue	EDM 1901

Formerly Graminæ

Alderney's

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
r	6 to 8	6,7,11. Bract barely longer than inflorescence
r	6 to 10	1,3,6,7
vlf	5 to 6	6,11. (Next species also found in Clonque Valley by JDG 1937)
lf	5 to 8	7. Old Mill leat in Bonne Terre only. Several large plants
o	6	5,6,13. Female glumes shorter & darker than utricles
o	6 to 7	7,8. Verge Route de Picaterre. Ripe spikes greenish-brown
r	6 to 7	7,8,14. Requires calcareous soil. Ripe spikes reddish-brown
la	3 to 9	Sandy shores, dunes, etc
vla	6 to 8	Mannez Quarry nr well head. Det. CJ. from BB specimen 'Mannez Quarry, one tuft' Only record 'Clanque cliffs' Only record (Mannez Quarry as <i>C. leporina</i>) Doubtful record because of habitat
lc	4 to 8	Cliff valleys. Stem 3-sided. Lowest bract with pubescent
1 site	5 to 8	4. Longis pond
vlf	4 to 8	7,8, Line of LeVallée stream. Seeds vigorously
f	4 to 8	2,3,4,5,9,11. Stems in loose glaucous tufts. Wet or dry grassland
o	5 to 6	12,13,14. Cliff valleys. Prefers wet acid soils
r	5 to 7	3,4,13. Wet rocky places near sea. (With HKAS)
r	6 to 7	5,13. Lowest bract much longer than infl. bent at rt angle EDM 1901 'C in Cliff valleys'. Possibly the only records
		3
vr	4 to 5	10. Habitat short grassland
vr	5 to 6	11
Alderney's Grasses		need more investigation. There are probably species not yet recorded
		Planted
		Planted
		Mannez Quarry. Doubtful record
r	5 to 8	Quite plentiful in 8,9,11. 5. Specimen in BM Only record
c	5 to 8	3,4,6,7,8,10 Possibly the only record

THE PLANT LIST

No.	Scientific name	English name	Finder/date
871	<i>Festuca ovina</i>	Sheep's-fescue	CCB 1838
872	<i>Festuca filiformis</i>	Fine-leaved Sheep's-fescue	DMcC 1973
873	<i>Festuca lemanii</i> #	Confused Fescue	CCB 1838
874	<i>Festuca longifolia</i> #	Blue Fescue	CCB 1838
875	<i>Festuca huonii</i>	Houn's Fescue	BB 1987
876	<i>Festuca maritima</i> #	Sea Fescue	GCD 1907
877	X <i>Festulpia hubbardii</i>	A hybrid Fescue	ABJ 1933
878	<i>Lolium perenne</i>	Perennial Rye-grass	CCB 1838
879	<i>Lolium x boucheanum</i>	A hybrid Rye-grass	PL 1976
880	<i>Lolium multiflorum</i>	Italian Rye-grass	EDM 1901
881	<i>Lolium temulentum</i> #	Darnel	CCB 1838
882	<i>Vulpia fasciculata</i> * (Scarce)	Dune Fescue	CCB 1838
883	<i>Vulpia bromoides</i>	Squirreltail Fescue	CCB 1838
884	<i>Vulpia myuros</i>	Rat's-tail Fescue	CCB 1838
885	<i>Vulpia ciliata</i> ssp. <i>ambigua</i> #	Bearded Fescue	GCD 1906
886	<i>Cynosurus cristatus</i>	Crested Dog's-tail	CCB 1838
887	<i>Cynosurus echinatus</i> #	Rough Dog's-tail	BB 1995
888	<i>Briza media</i> #	Quaking-grass	EDM 1901
889	<i>Briza minor</i> #	Lesser Quaking-grass	BB 1986
890	<i>Briza maxima</i>	Greater Quaking-grass	WFS 1986
891	<i>Poa infirma</i> * (RDBk)	Early Meadow-grass	MMcW
892	<i>Poa annua</i>	Annual Meadow-grass	CCB 1838
893	<i>Poa annua</i> var. <i>reptans</i> #		MH 1978
894	<i>Poa trivialis</i>	Rough Meadow-grass	CCB 1838
895	<i>Poa humilis</i>	Spreading Meadow-grass	CCB 1838
896	<i>Poa pratensis</i>	Smooth Meadow-grass	CCB 1838
897	<i>Poa angustifolia</i>	Narrow-leaved	TR 1990
898	<i>Poa chaixii</i> #	Broad-leaved	EDM 1901
899	<i>Poa bulbosa</i> * (Scarce)	Bulbous Meadow-grass	
900	<i>Dactylis glomerata</i>	Cock's-foot	CCB 1838
901	<i>Catabrosa aquatica</i> #	Whorl-grass	JES 1970
902	<i>Catapodium rigidum</i> .	Fern-grass	EDM 1901
903	<i>Catapodium marinum</i>	Sea Fern-grass	CCB 1838
904	<i>Parapholis strigosa</i> #	Hard-grass	JES 1969
905	<i>Parapholis incurva</i> #	Curved Hard-grass	HJMB 1957

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes	0
c	5 to 9	The many subspecies have not been separated	
o	5 to 6	1,7	
		Possibly the only record	
		Possibly the only record	
r		4. Rocky scree Saye Bay (As <i>Vulpia membranacea</i>)	Only record
		Other Festuca hybrids may exist	
c	5 to 12	Spikelets with 4-14 florets	
r	5 to 9	7. Refound BB 1987	
vlf	5 to 9	Spikelets with 11-22 florets	
		St. Anne'	Only record
f	5 to 7	3,4,5,7,8,10. Spikelets 10-18mm	
c	4 to 7	3,7,8. Spikelets 6-11mm	
vr	5 to 7	10 Refound above Impôt BB 8.6.89	
		Braye & Corblets	Only record
c	4 to 8	All ex 1,8.	
vr	6 to 7	4,5	
r	6 to 7	3,6,9	
vr	7	9	
lf	1 to 12	'2,7	
vlf	3 to 5	Culms to 10cm (and on Burhou)	
c	1 to 12	Culms to 20cm	
			Only record Burhou
f	3 to 7	6,7. Culms to 70cm	
f	5 to 7	7,9. Culms to 30cm	
c	5 to 7	9. Culms to 75cm	
1 site	5 to 7	8. Culms to 70cm	
		10	Only record
	4 to 5		Doubtful record
a	1 to 12		
vr	6 to 8	Wet, sandy places near sea.	Possibly the only record
o	4 to 7	3,4,5,7. Walls and bare ground. Prefers alkaline soil. Very rigid	
c	4 to 9	1,2,4,7. Very shiny. Later flowering than above. Often reddish	
			Only record
			Only record

THE PLANT LIST

No.	Scientific name	English name	Finder/date
906	<i>Glyceria fluitans</i>	Floating Sweet-grass	CCB 1838
907	<i>Glyceria x pedicellata</i>	Hybrid Sweet-grass	JO 1953
908	<i>Glyceria declinata</i> #	Small Sweet-grass	JES 1969
909	<i>Glyceria notata</i>	Plicate Sweet-grass	EDM 1901
910	<i>Helictotrichon pubescens</i>	Downy Oat-grass	EDM 1901
911	<i>Helictotrichon pratense</i> #	Meadow Oat-grass	GCD 1906
912	<i>Arrhenatherum elatius</i>	False Oat-grass	CCB 1838
913	<i>Arrhenatherum elatius</i> ssp. <i>bulbosum</i>	False Oat-grass	EDM 1901
914	<i>Avena strigosa</i> #	Bristle Oat	EDM 1901
915	<i>Avena fatua</i> #	Wild-oat	CCB 1838
916	<i>Avena sterilis</i> ssp. <i>ludoviciana</i>	Winter Wild-oat	# DMcC 1963
917	<i>Avena sativa</i>	Oat	DMcC 1958
918	<i>Gaudinia fragilis</i> #	French Oat-grass	ABJ 1933
919	<i>Trisetum flavescens</i>	Yellow Oat-grass	CCB 1838
920	<i>Koeleria macrantha</i>	Crested Hair-grass	EDM 1901
921	<i>Holcus lanatus</i>	Yorkshire-fog	CCB 1838
922	<i>Holcus mollis</i> #	Creeping Soft-grass	EDM 1901
923	<i>Aira caryophyllea</i>	Silver Hair-grass	CCB 1838
924	<i>Aira praecox</i>	Early Hair-grass	CCB 1838
925	<i>Anthoxanthum odoratum</i>	Sweet Vernal-grass	CCB 1838
926	<i>Phalaris aquatica</i> #	Bulbous Canary-grass	JP 1977
927	<i>Phalaris canariensis</i> #	Canary-grass	WH 1881
928	<i>Phalaris minor</i>	Lesser Canary-grass	CRPA 1889
929	<i>Agrostis capillaris</i>	Common Bent-grass	CCB 1838
930	<i>Agrostis gigantea</i>	Black Bent	ABJ 1932
931	<i>Agrostis stolonifera</i>	Creeping Bent	CCB 1838
932	<i>Agrostis canina</i> #	Velvet Bent	EDM 1901
933	<i>Ammophila arenaria</i>	Marram	CCB 1838
934	<i>Lagurus ovatus</i>	Hare's-tail	RM 1951
934	<i>Apera spica-venti</i> * (Scarce)	Loose Silky-bent	JES 1966
936	<i>Polypogon viridis</i>	Water Bent	GCD 1906
937	<i>Alopecurus pratensis</i>	Meadow Foxtail	EDM 1901
938	<i>Alopecurus geniculatus</i>	Marsh Foxtail	EDM 1901
939	<i>Phleum pratense</i>	Timothy	EDM 1901
940	<i>Phleum bertolinii</i>	Smaller Cat's-tail	GM 1969

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
vlf	5 to 8	7,12,13
o	6 to 7	5
vr	6 to 7	12
o	5 to 7	7,9,10,13
o	6 to 7	4,9,12
		Braye'
		Only record
c	5 to 11	1,2,3,6,7,8,9,10,11
c	5 to 11	1,2,7. Bulbils at base of stems
r	6 to 7	9,12
		Refound BB 1987
r	6 to 9	4,10,12,13
		Determined by AM.
		Only record
o	5 to 8	4,12,13. Relic of cultivation
r	5 to 7	4,8. Whitegates. (With HKAS) Roadside nr. Dairy DMcC 1963
o	6 to 8	4,8,9
f	6 to 7	4,5
c	4 to 8	
r	7 to 9	7
c	4 to 7	Stems to 25cm
c	4 to 5	Stems to 10cm
c	4 to 9	
		10
		Only record
r	6 to 7	7,9,10. Probably a casual from bird seed. Panicle ovoid
lf	6 to 11	1,7,8,12,13,14. Panicle longer & slenderer
c	5 to 9	
?	6 to 8	
		Refound FH
c	3 to 11	
r	5 to 8	7
la	6 to 9	3,4,5,6. Dunes & commons.
r	5 to 9	2,4,5,7,9
		Refound JES 1966, BB
		Only record ?
o	6 to 7	3,8. AS 1956 det. FLS No site. Refound in a gutter in town MK 1995
r	4 to 6	8,12
r	4 to 8	11,12
		Refound ABJ 1932
r	6 to 8	8,10. Culms erect to 1.5m
vr		Culms to 50cm Cliff path BB 1989, not conf. ? the only records

THE PLANT LIST

No.	Scientific name	English name	Finder/date
941	<i>Phleum arenarium</i>	Sand Cat's-tail	CCB 1838
942	<i>Bromus commutatus</i> #	Meadow Brome	GCD 1906
943	<i>Bromus hordeaceus</i> ssp. <i>hordeaceus</i>	Soft Brome	CCB 1838
944	<i>Bromus hordeaceus</i> ssp. <i>ferronii</i>	Least Soft-brome	ABJ 1933
945	<i>Bromus hordeaceus</i> ssp. <i>thominei</i>	Least Soft-brome	EDM 1901
946	<i>Bromus</i> x <i>pseudothominei</i>	Lesser Soft-brome	GCD 1906
947	<i>Anisantha diandra</i>	Great Brome	EDM 1901
948	<i>Anisantha rigida</i>	Ripgut Brome	EDM 1901
949	<i>Anisantha sterilis</i>	Barren Brome	CCB 1838
950	<i>Anisantha madritensis</i>	Compact Brome	DMcC 1956
951	<i>Ceratochloa cathartica</i>	Rescue Brome	ABJ 1933
952	<i>Ceratochloa carinata</i> #	California Brome	JP 1979
953	<i>Brachypodium pinnatum</i> #	Tor-grass	DMcC 1973
954	<i>Brachypodium sylvaticum</i>	False Brome	CCB 1838
955	<i>Elymus caninus</i>	Bearded Couch	BB 1987
956	<i>Elytrigia repens</i>	Common Couch	CCB 1838
957	<i>Elytrigia repens</i> var. <i>aristatum</i> #		GCD 1932
958	<i>Elytrigia atherica</i>	Sea Couch	EDM 1901
959	<i>Elytrigia juncea</i>	Sand Couch	CCB 1838
960	<i>Elytrigia</i> x <i>obtusiuscula</i> #	A hybrid Couch	GCD 1907
961	<i>Elytrigia</i> x <i>laxa</i>	A hybrid Couch	FWS 1958
962	<i>Leymus arenarius</i> #	Lyme-grass	BB 1993
963	<i>Hordeum vulgare</i>	Six-rowed Barley	?
964	<i>Hordeum distichon</i>	Two-rowed Barley	NJ 1974
965	<i>Hordeum murinum</i>	Wall Barley	CCB 1838
	<i>Subfamily Arundinae</i>		
966	<i>Danthonia decumbens</i> #	Heath-grass	CCB 1838
967	<i>Cortaderia selloana</i>	Pampas-grass	BB 1988
968	<i>Phragmites australis</i>	Common Reed	CCB 1838
	<i>Subfamily Chloridoideae</i>		
969	<i>Cynodon dactylon</i> *	Bermuda-grass (RDBk)	AS 1955
	<i>Subfamily Paniceae</i>		
970	<i>Panicum capillare</i> #	Witch-grass	OJ 1983
971	<i>Panicum miliaceum</i> #	Common Millet	JP 1977
972	<i>Echinochloa crusgalli</i> #	Cockspur	EDM 1900

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
lc	5 to 7	1,3,4,5. Sandy coast and commons. Rarely above 10-15cm Only record
c	4 to 9	Culms to 80cm
f	5 to 7	1,4, 7, Culms 10-15cm
lf	5 to 7	4, 5, 9, 10-14, Culms 5-8cm (Marquand as <i>B. nanus</i>)
o	5 to 8	8,9,10,11. Culms erect, to 80cm Refound FLS 1984
a	5 to 8	2,7,8,9,12,13
lc	5 to 8	Braye
a	4 to 10	
r	5 to 7	
r	5 to 10	9,10. Near former Devereux House Hotel 4. 1st Bailiwick record. Only record
vr	6 to 7	2,4,9. First Bailiwick record Longis. Has extensive rhizomes
r	7 to 8	7,8,13. Few short rhizomes
o	7	7 GCD 1932 Marked 'S'
c	5 to 10	
vr	7 to 8	GCD 1932 Marked 'S'. Refound PL 1976 Det. CEH
f	6 to 10	1,2,3,4,5,11. Older dune margins, especially if damp
c	5 to 11	1,2,3,4,5,7,14. Mostly on dunes
r	6 to 9	Last confirmed record, AS 1956. Determined FLS
vr	6 to 8	Determined by DMcC. later doubtful
r	6 to 7	Relic of cultivation
o	5 to 8	2,11,13 Previous record 1956 from Burhou
lf	5 to 9	Increasing considerably last 5 years
o	7	Heaths & cliffs
vr	6 to 12	7. Self seeded from planted specimens
vla	5 to 10	4
vlc	5 to 10	4,5. Spreading considerably behind sea wall in last ten years
		8 Only record, probably from birdseed
vr	3,4,9,	Refound BB 6/1989
vr	4,7.	Refound Saye Farm, JP 1977

THE PLANT LIST

No.	Scientific name	English name	Finder/date
973	<i>Setaria pumila</i>	Yellow Bristle-grass	ML 1992
973a	<i>Setaria viridis</i> #	Green Bristle-grass	BB 2000
974	<i>Sorghum halapense</i> #	Johnson-grass	OJ 1982
975	<i>Sorghum bicolor</i> #	Great Millet	DMcC 1982
975a	<i>Digitaria sanguinalis</i>	Hairy Finger-grass	LP2007
976	<i>Zea mays</i>	Maize	BB 1993
Sparganiaceæ			
977	<i>Sparganium erectum</i> #	Branched Bur-reed	EDM 1901
Typhaceæ			
978	<i>Typha latifolia</i>	Bulrush	DMcC 1961
979	<i>Typha angustifolia</i>	Lesser Bulrush	BB 1987
Liliaceæ			
Now includes the forms r Amaryllid			
980	<i>Kniphofia praecox</i>	Greater Red-hot-poker	BB 1987
981	<i>Kniphofia tuckii</i>	A Red-hot-poker	BB 1989
982	<i>Tulipa gesneriana</i> #	Garden Tulip	DB 1994
983	<i>Convallaria majalis</i> #	Lily-of-the-valley	DMcC 1963
983	<i>Polygonatum multiflorum</i> #	Solomon's Seal	BB 1993
984	<i>Ornithogallum angustifolium</i>	Star-of-Bethlehem	CCB 1838
985	<i>Scilla peruviana</i> #	Portuguese Squill	TWA 1958
986	<i>Scilla autumnalis</i>	Autumn Squill	CCB 1838
987	<i>Hyacinthoides non-scripta</i>	Bluebell	CCB 1838
988	<i>Hyacinthoides non-scripta</i> x <i>hispanica</i>	A hybrid Bluebell	DMcC 1973
989	<i>Hyacinthoides hispanica</i>	Spanish Bluebell	CMR 1950
990	<i>Muscari comosum</i> #	Tassel Hyacinth	CH /RW
990a	<i>Muscari armeniacum</i> #	Grape Hyacinth	BB 1997
991	<i>Allium schoenoprasum</i> #	Chives	NVM 1967
992	<i>Allium roseum</i> ssp. <i>bulbiferum</i>	Rosy Garlic	NVM 1982
993	<i>Allium neapolitanum</i> #	Neapolitan Garlic	BB1988
994	<i>Allium subhirsutum</i> #	Hairy Garlic	FLI 1962
995	<i>Allium triquetrum</i>	Three-cornered Garlic	EDM 1901
996	<i>Allium sphaerocephalon</i> #	Round-headed Leek	BB 1987
997	<i>Allium vineale</i>	Wild Onion	DMcC 1967
998	<i>Tristagma uniflorum</i>	Spring Starflower	BB 1988

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
vr	7 to 8	10. Gravel at Old Barn
o	8 to 10	7. Platte Saline. 2 plants Confirmed ML. 1st island record 20.8.00
vr	7 to 8	7,8. Refound above edge of old sandpit Platte Saline, BB 1992
vr	7 to 8	3,4. By seats Braye Bay. Refound BB 1989
vr	9 to 10	8. 8 plants wall base Chemin du Meunier Det . BB 1st record
vr	7 to 8	7, Relic of old crop in field in Le Petit Val
r	6 to 8	4,7 Refound FF 1967, ML1996 in Longis pond
1 site	7 to 9	10. Tank on Essex Hill, probably planted originally
1 site	6 to 9	5. Mannez pond. Male & female flowers separated by stem
aceae		
o	1 to 12	5,7,10. Scattered garden escape
vr	1 to 12	5,9
	4	2, Garden escape or throwout Only record
vr	2 to 7	9. Garden escape. Refound BB 1988
vr	4 to 6	7. St Vignalis Garden, Pont Martin. Possibly planted
r	2 to 6	4,5,8,9
1 site	6 to 7	4. Single plant persisting many years since 15 were dug up c. 1980
la	7 to 10	E. & S. Coasts, Corblets Quarry & Mannez Garenne
c	3 to 6	Less robust than hisp. Tip of raceme bent over. Also with white fls.
f	3 to 5	11. Probably frequent, characters intermediate difficult to determine
la	3 to 5/6	Tip of raceme straight. Occasionally with white flowers
1 site	5 to 6	8. above quarry below Butes. Still there 1994
1 site	3 to 5	9. Parking area to W of Impôt, probably escaped from tip
	4 to 8	Garden escape Only record
	4 to 6	Only record
vr	5 to 7	8,9. 1962-82 records error for following sp. Spathe 1 large, broad
r	3 to 6	4, 5,6,7,8,9,10. Spathe 3, small
a	9 to 7	Single patch in 1900. On Mauvaises Herbes list 1952-2003
vr	6 to 8	7. Infl has fls only, no bulbils
r	5 to 8	3,4,5. Infl. has fls & bulbils or bulbils only
2 sites	2 to 4	4,10. Garden escape. Fort Essex & Route des Mielles in asphalte

THE PLANT LIST

No.	Scientific name	English name	Finder/date
999	<i>Amaryllis belladonna</i>	# Jersey Lily	JP 1985
1000	<i>Leucojum aestivum</i>	# Summer Snowflake	MMcW
1001	<i>Narcissus</i> spp. ag.	Mostly relics of cultivation in the 1948	
1002	<i>Narcissus x medioluteus</i>	Primrose-peerless	KF 1957
1003	<i>Narcissus poeticus</i>	Pheasant's-eye Daffodil	CMR 1954
1004	<i>Narcissus pseudonarcissus</i>	Daffodil	CMR 1954
1005	<i>Narcissus</i> x ?	Scilly White	JR 1957
1006	<i>Asparagus officinalis</i> ssp. <i>officinalis</i>	Garden Asparagus #	DMcC 1957
1007	<i>Ruscus aculeatus</i>	Butcher's Broom	CCB 1838
1008	<i>Crinum powelli</i>	# Crinum	BB 1994

Iridaceae

1009	<i>Iris germanica</i>	# Bearded Iris	BB 1989
1010	<i>Iris pseudacorus</i>	Yellow Iris	CCB 1838
1011	<i>Iris foetidissima</i>	Stinking Iris	EDM 1901
1011a	<i>Iris x hollandica</i>	# Dutch Iris	BB 1989
1012	<i>Romulea columnae</i>	* Sand Crocus (RDBk)	CCB 1838
1013	<i>Gladiolus communis</i>	Eastern Gladiolus	CMR 1954
1013a	<i>Ixia paniculata</i>	# Corn Lily	BB 2000
1014	<i>Crocsmia x crocosmiflora</i>	Montbretia	CMR 1950

Agavaceae

1015	<i>Agave americana</i>	# Century Plant	BB 1986
1016	<i>Cordyline australis</i>	Cabbage Palm	BB 1986
1017	<i>Phormium tenax</i>	# New Zealand Flax	BB 1987
1018	<i>Trachycarpus fortunei</i>	# Chusan Palm	RW 1987

Orchidaceae

1019	<i>Spiranthes spiralis</i>	Autumn Lady's-tresses	EDM 1901
1020	<i>Anacamptis pyramidalis</i>	Pyramidal Orchid	EDM 1901
1021	<i>Dactylorhiza fuchsii</i>	# Common Spotted-orchid	EDM 1901
1022	<i>Dactylorhiza maculata</i>	# Heath Spotted Orchid	CMR 1954
1023	<i>Dactylorhiza incarnata</i>	# Early Marsh-orchid	CMR 1954
1024	<i>Dactylorhiza praetermissa</i>	# Southern Marsh-orchid	EDM 1900

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
1 site	7 to 11	8, Roadside bank near school
2 sites	2 to 5	8,9 Found in The Terrace BB
-1960s pe ri od. The re are several unidentified species in hedges and fields		
o	3 to 5	
f	11 to 5	4,5,6,7,10,12
1 site	3 to 5	4,7. Field beside Fort Corblets. Probably planted
o	1 to 5	
vr	5 to 7	9. Garden escape
o	12 to 4	5,9,10,12,13,14. Cliffs
1 site	5 to 10	3. Large clump in dune behind Saye Bay, W. end
vr	7 to 8	5. verge near Lighthouse for many years
vlc	4 to 7	1,4,7,10,12. Now only round Longis & Platte Saline
f	6 to 7	Abundant in Sq.4. Occasional in other damp spots
r	3 to 5	7. Under hedge in Le Petit Val for many years. About 6 +
lf	3 to 5	Cliffs, rim of Berry's quarry and E. coast
o	4 to 7	Scattered round coasts
vr	5 to 6	9. Roadside bank N. side Bluestone Hill 5 plants
lf	7 to 9	Scattered, cliffs & quarries
vr	?	5, 7. Planted many years ago, has suckered
lf	5 to 8	3,7,8,10. Planted in the 1930's, and in many gardens.
r	6 to 8	7,8. Garden escape. Field on cliffs sq, 7 ?planted
vr	5 to 6	8,13. Planted in several gardens
c	7 to 10	Scattered in short turf on dunes and cliffs
lc	5 to 10	1,2,3,4,5,7,8,9,10. Short sandy turf
vr	6 to 8	7. Lvs with dark elongated spots
vr		3, 10. single plant Essex Hill 1989. Lvs with dark round
		7. CMR 1954, JM 1960, both in Clonque Quarry
r	5 to 8	7,11. A few specimens, St. Vignalis & old Mill leat. No spots

THE PLANT LIST

No.	Scientific name	English name	Finder/date
1025	<i>Orchis morio</i>	Green-winged Orchid	EDM 1902
1026	<i>Orchis mascula</i>	# Early-purple Orchid	CMR 1954
1027	<i>Ophrys apifera</i>	Bee Orchid	SH 1949

THE PLANT LIST

Frequency	Flowering period	Squares where found and notes
vlf	4 to 6	4,5. Mannez Garenne & E. coast
vr	5	4, Longis Common, large patch (?confused with above) VKM 1957
2 sites	6 to 7	4. A single colony, 1-7 plants. None seen 1983-1991 Recovered 1992 onwards with variable numbers. Despite States mowing and some very poor years 25 plants flowered in May/June 2007
1,052		Total number of species recorded to November 2007 Additions may be made on the following 2 pages. New records will be found on my website the following year, Please visit flora.org.gg

