

The Illustrated Wild Flowers of Alderney, a revised check list with notes

THE PLANT LIST

Text & Illustrations © Brian Bonnard 2012

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 1

It should be noted that in .pdf format the hypertext links do not function

PTERIDOPHYTA

Notes; >>>> Click on sequence number blue links for a distribution map at 1km square and on plant name link for a photo
Distribution maps are not always drawn for abundant or common species See A4 map of Alderney above on this site for square numbers,
or use a copy printed from file Alderne3.jpg Note; Individual squares are not generally shown for abundant, common or frequent plants

Lycopodiopsida

Selaginellaceae

1 **2/1.2** [Selaginella kraussiana](#) Kraus's clubmoss JP 1993 1 site 8, Large patch 10 x 1.5m
along base of church wall NW corner

Isoetaceae

2 **3/1.3** [Isoetes histrix](#) Land Quillwort (RDBk) EDM 1902 vr 1 to 6 5,10,14. "Abundant E. La
Quoire 1902" 2 new sites E coast 6/1995, JB, MK

Equisetopsida

Equisetaceae

3 **4/1.5** [Equisetum arvense](#) Field Horsetail EDM 1901 lc 3 to 7 Generally distributed in damp places.
Separate fruiting stem

4 **4/1.8** [Equisetum palustre](#) Marsh Horsetail CCB 1838 lf 4 to 8 4,7,8,9,10,13. Cones on vegetative
shoots.

5 **4/1.9** [Equisetum telmateia](#) Great Horsetail CRPA 1900 2 sites in Le Grand Val. Only natural site in CIs
3 to 5 12, Veg stem to 1.5m. Fruiting stems 25cm, brown, appear first

Pteropsida

Ophioglossaceae

6 **5/1.1** [Ophioglossum vulgatum](#) Adder's-tongue JES 1969 1 site in Mannez quarry 5 to 8 5
7 **5/1.3** [Ophioglossum lusitanicum](#) Least Adder's-tongue (RDBk) EDM 1906 Only
record, probably an error

[27](#) [19/1.1](#) [Azolla filiculoides](#) Water Fern JES 1972 4 sites in Mannez & Corblets quarries & Platte Saline ponds.
6 to 9 1,4,5,7 Found with spores 22.8.88 and other, later, years

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 4

Gymnospermæ

Pinopsida

Mostly planted

Pinaceæ

28	20/4.2	Picea abies	Norway Spruce	BB 1987	vr	5 to 6	
29	20/5.1	Larix decidua	European Larch	CH 1987	r	3 to 6	8. 3 specimens to 13m. others smaller
30	20/6.1	Cedrus deodora	Deodar	CH 1987	vr	12 to 1	9
31		Cedrus libani	Cedar-of-Lebanon	BB1987	vr	10 to 12	
32	20/7.1	Pinus sylvestris	Scots Pine (Scarce)	CH 1987	vr	3 to 7	3,5,7,8,13. Pollen found in ancient peat at Longis
33	20/7.2a	Pinus nigra subsp. nigra	Austrian Pine	CH 1987	r	4 to 7	3,8
34	20/7.2b	Pinus nigra subsp. laricio	Corsican Pine	BB 1986	r	4 to 6	3,8. Specimen to 16.5.m
35	20/7.4	Pinus pinaster	Maritime Pine	BB 1986	o	4 to 7	3,7,8,10,13. Plantation Newtown Road
36	20/7.5	Pinus radiata	Monterey Pine	DMcC 1975	c	2 to 7	Specimens to 14.5m. Most frequent planting
37		Pinus attenuata	Knobcone Pine	BB 1987	1 site	5 to 6	8. One stand in Newtown Road
38		Tsuga heterophylla	Western Hemlock	RW 1987	1 site, Island Hall garden	4 to 5	8. Planted

Cupressaceæ

39	21/2.1	Chamaecyparis lawsoniana	Lawson's Cypress	BB 1987	o	3 to 4	3,7,9,13. 'Mostly in gardens'
40	22/2.1	X Cupressocyparis leylandii	Leyland Cypress	BB 1987	o	5 to 6	Much used in hedging
41	21/1.1	Cupressus macrocarpa	Monterey Cypress	BB 1986	r	5 to 6	3,7,8,9. Specimens to 20.5m badly damaged in hurricane 10/1987
42		Thuja occidentalis	White Cedar	CH 1987	vr	8	
43	21/3.1	Thuja plicata	Western Red-cedar	BB 1985	1 site, Island Hall garden	3	8.

Araucariaceæ

44	22/1.1	Araucaria araucana	Monkey-puzzle	BB 1986	r	6	8. Several in Gardens
----	------------------------	------------------------------------	---------------	---------	---	---	-----------------------

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 5

Taxaceæ

45	23/1.1	Taxus baccata	Yew	BB 1986	vr	2	8. Only in gardens, churchyard and public places
----	------------------------	-------------------------------	-----	---------	----	---	--

MAGNOLIOPSIDA

Angiospermae

1. Magnoliidæ

(Formerly Dicotyledonidæ)

Lauraceæ

46	24/1.1	Laurus nobilis	Bay	BB 1987	r	3 to 6	4, 8, Large specimens, Vicarage, St. Vignalis, Berry's Quarry & gardens
----	------------------------	--------------------------------	-----	---------	---	--------	---

Nymphaeaceæ

47	26/1.1a	Nymphaea alba	White Water-lily	BB 1986	2 sites	5 to 9	4,10
48	26/2.1	Nuphar lutea	Yellow Water-lily	BB 1989	6 to 7	10. Not refund, possible error for <i>Nymphoides peltata</i> . Only record	

Ceratophyllaceæ

49	27/1.2	Ceratophyllum submersum	Soft Hornwort	BB 1987	1 site, Corblets pond	7 to 9	4. Leaves. light green, segments scarcely toothed
----	--------	---	---------------	---------	-----------------------	--------	---

Ranunculaceæ

50	28/1.1	Caltha palustris	Marsh Marigold	AJR 1966	vr	4 to 7	7. Now seems to be confined to domestic ponds
50a	28/3.1	Helleborus foetidus	Stinking Hellebore	BB 2011	o/vlf	1 to 12	8. Frequent garden weed, occasional escapes. Previously omitted from lists
51	28/7.1	Consolida ajacis	Larkspur	EDM 1901	vr	6 to 7	8. Occasional garden escape. Only other record 1957
52	28/12.1	Clematis vitalba	Traveller's-joy	CMR 1950	vlf	5 to 10	2,3,8,9, 12 Newtown, Rochers, Braye Dunes
53	28/13.1	Ranunculus acris	Meadow Buttercup	CCB 1838	c	12 to 9	
54	28/13.2	Ranunculus repens	Creeping Buttercup	EDM 1900	c	1 to 12	
55	28/13.3	Ranunculus bulbosus	Bulbous Buttercup	CCB 1838	c	12 to 8	

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 6

56	28/13.4	Ranunculus sardous	Hairy Buttercup	BB 1987	vr	3 to 4	7
57	28/13.7	Ranunculus parviflorus	Small-flowered Buttercup	EDM 1901	o	4 to 6	4,5,7,9, 10,14
58	28/13.12	Ranunculus lingua	Greater Spearwort	CCB 1838	vr, Platte Saline pond since 1991. ? garden escape	6 to 7 4,7,12,13 Refound BB 1991. Previous GCD 1906, CMR 1954	
59	28/13.13	Ranunculus flammula	Lesser Spearwort	EDM 1901	o	4 to 8	7,11,13. In damp valleys
60	28/13.17	Ranunculus ficaria agg.	Lesser Celandine	EDM 1901	la	1 to 6	
61	28/13.17a	<i>Ranunculus ficaria</i> subsp. <i>ficaria</i>	Lesser Celandine	BB 1996			10. Small flowers, 25mm, long stems, leaves spreading, no bulbils
62		<i>Ranunculus ficaria</i> subsp. <i>calthifolius</i>	Lesser Celandine	BB 1996			7. Small flowers short stems, congested leaves, no bulbils
64	28/13.17d	<i>Ranunculus ficaria</i> subsp. <i>chrysocephalus</i>	Lesser Celandine	BB 1996			7. Large fls35mm. No bulbils
65	28/13.18	<i>Ranunculus hederaceus</i>	Ivy-leaved Crowfoot	CCB 1838	vr	6 to 9	6,7, Vau Pommier. "Still there" AB/AKJ 1934. Only records
66	28/13.22	Ranunculus trichophyllus	Three-leaved Water-crowfoot	EDM 1901	vlc	5 to 6	4, 5. Mannez & Longis ponds. Berry's Quarry pool
67	28/17.4	Thalictrum minus	Lesser Meadow-rue	MM 1985	vr	6 to 8	3,5

Berberidaceæ

68	29/1.1	Berberis vulgaris	Barberry	BB 1988	vr	1 to 8	10, Longis Bay Determined by DMcC 6/88
69	29/1.9	Berberis darwinii	Darwin's Barberry	BB1989	vr	10 to 6	4, Corblets Bay in scrub

Papaveraceæ

70		<i>Roemeria hybrida</i>	Violet Horned Poppy	VKM 1956		Now considered extinct in Britain		10. Nr. Essex Castle. Only record
71	30/1.3	Papaver somniferum	Opium Poppy	EDM 1901	o	5 to 11	2,5,7,8, 11-14	
72	30/1.4	Papaver rhoeas	Common Poppy	CCB 1838	f	5 to 12	1,2,3,7,9	
73	30/1.5	Papaver dubium	Long-headed Poppy	EDM 1901	lf	6 to 9	6,10,11,12	
74	30/1.6	Papaver hybridum	Rough Poppy	CCB 1838	o	5 to 8	7,8,10,14	
75	30/4.1	Glaucium flavum	Yellow Horned-poppy	CCB 1838	lc	5 to 11	1,2,3,7,12	

Juglandaceae Planted
 97 37/1.1 [Juglans regia](#) Walnut NVM 1966 1 site 5 8 Single specimen in Island Hall garden

Fagaceae Mostly planted
 98 39/1.1 [Fagus sylvatica](#) Beech DMcC 1957 o 4 to 5 7,8 Including copper beeches
 99 39/3.1 [Castanea sativa](#) Sweet Chestnut EDM 1901 o 5 to 7 7,8,10. Scattered
 100 39/4.2 [Quercus cerris](#) Turkey Oak DMcC 1957 o 4 to 7 8, plus scattered small specimens
 101 39/4.3 [Quercus ilex](#) Evergreen Oak CH 1987 f 4 to 7 Frequent planting, self-seeds
 102 39/4.5 [Quercus petraea](#) Sessile Oak BB 1987 vr 4 to 7 7,8,9.
 103 39/4.6 [Quercus robur](#) Pedunculate Oak CCB 1838 r 4 to 7 2,7,8,9,10,13.

Betulaceae Mostly planted
 104 40/1.1 [Betula pendula](#) Silver Birch EDM 1901 o 3 to 6 7,8,9,10, 13.
 105 40/2.1 [Alnus glutinosa](#) Alder DMcC 1975 o 1 to 5 6,7,8,9,12,13. Watered valleys, self-seeds
 106 40/3.1 [Carpinus betulus](#) Hornbeam RW 1986 vr 4 to 5 8. Churchyard
 107 40/4.1 [Corylus avellana](#) Hazel RW 1987 vr 1 to 4 4, 10.

Nyctaginaceae
 108 43a [Mirabilis jalapa](#) Marvel-of-Peru BB 1987 vr 7 to 10 Occasional persisting garden weed or escape

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
 Flowering period Squares where found and notes

Page 9

Aizoaceae
 109 42/1.1 [Aptenia cordifolia](#) Heart-leaf Ice-plant BB 1987 vr 3 to 12 2,8. On old walls
 109a 42/5.1 [Disphyma crassifolia](#) Purple Dewplant Hampshire bots (PB) o 1st record 6 to 11 8, 11 Previously overlooked as Aptenia
 110 42/3.2 [Lampranthus roseus](#) Rosy Dew-plant MR 1978 o 1 to 12 6,10,13. (S. Cliffs, first noted July 1978)
 111 42/8.1 [Carpobrotus acinaciformis](#) Sally-my-Handsome DMcC 1959 o 3 to 7 4,5,7,12,13. Flowers 8-10cm. pink/ purple. Leaves curved, thickest near top
 112 42/8.2 [Carpobrotus edulis](#) Hottentot-fig JO 1953 f 1 to 12 Old forts and cliffs. Flowers 4-9cm pink/purple or yellow. Leaves equal thickness
 113 42/8.3 [Carpobrotus glaucescens](#) Angular Sea-fig BB 1988 lf 3 to 10 2,4,5,6,7. Flowers 3.5-6cm, pink/purple, petals white at base
 113a [Tetragonia tetragonioides](#) New Zealand Spinach JG 2006 r Garden esxcape 1sy record 6-10
 5, 8

Chenopodiaceae
 114 43/1.8 [Chenopodium polyspermum](#) Many-seeded Goosefoot FS/CR 1958 vr 6 to 9 7. Refound BB 1995 Platte Saline
 115 43/1.9 [Chenopodium vulvaria](#) Stinking Goosefoot (RDBk) ? 7 to 9 **Marked as ? errors in WFG.**
 Recorded in Atlas 1962
 116 43/1.12 [Chenopodium murale](#) Nettle-leaved Goosefoot CCB 1838 f 5 to 10 Waste places
 117 43/1.13 [Chenopodium ficifolium](#) Fig-leaved Goosefoot EDM 1901 r 6 to 9 7,8,14. 1 plant Butes Hill 1901. GCD 1906
 118 43/1.14 [Chenopodium opulifolium](#) Grey Goosefoot GCD 1906 **No site. Only record**
 119 43/1.15 [Chenopodium album](#) Fat-hen CCB 1838 c 6 to 10 Gardens & cultivated ground
 120 43/3.2 [Atriplex prostrata](#) Spear-leaved Orache EDM 1901 f 7 to 9 1,2,3,5,8,11. Most shores
 121 43/3.2 x 3 [Atriplex prostrata](#) x [A. glabriuscula](#) EMJ 1967 ? **"On beaches". Only record**
 122 43/3.3 [Atriplex glabriuscula](#) Babington's Orache EDM 1901 o 6 to 10 1,2,3,4,6,7. Sandy shores
 123 43/3.6 [Atriplex littoralis](#) Grass-leaved Orache CCB 1838 7 to 8 **Possibly the only record**

124	43/3.7	Atriplex patula	Common Orache	CCB 1838	f	6 to 10	
125	43/3.8	Atriplex laciniata	Frosted Orache	EDM 1901	f	7 to 10	1,2,3,4,5. Beaches, just about HWM
126	43/3.10	Atriplex portulacoides	Sea-purslane	WC 1838	r	6 to 7	12,13. Almost never at beach level here, mostly on cliff faces
127	43/4.1a	Beta vulgaris subsp. maritima	Sea-beet	CCB 1838	c	5 to 9	
128	43/8.1	Salsola kali	Prickly Saltwort	CCB 1838	o	6 to 9	2,3,4,5. At top of sandy beaches

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 10

Amaranthaceae

129	44/1.1	Amaranthus retroflexus	Common Pigweed	DMcC 1957	r	8 to 9	2,8,9,10,13. and garden weed
---------------------	------------------------	--	----------------	-----------	---	--------	------------------------------

Portulacaceae

130	45/1.1	Portulaca oleracea	Common Purslane	ML2001	vr	8	10, First record, in gravel at The Old Barn, Longis. 27/8/2001
131	45/2.2	Claytonia sibirica	Pink Purslane	BB 1987	vr	4 to 7	7. Occasionally persisting garden weed
132	45/3.1a	Montia fontana subsp. fontana	Blinks	EDM 1901	vlf	4 to 10	5,8,10,11,13. Seeds smooth & very shiny
133	45/3.1d	Montia fontana subsp. chondrosperma		AB/AKJ 1934			Det by SMW 1963as subsp chondrosperma. Seeds with tubercles, dull
134	45/3.1b	Montia fontana subsp. variabilis		EDM 1901			11,13. (as <i>M. f. var. rivularis</i>) Trois Vaux & Val du Sud. Only records
135	45/3.1c	Montia fontana subsp. amporitana		1962			Det. by SMW 1963 as subsp. intermedia. Seeds 3 rows pointed tubercles

Caryophyllaceae

136	46/1.1a	Arenaria serpyllifolia subsp. serpyllifolia	Thyme-leaved Sandwort	CCB 1838	f	1 to 11	4,7,10. Common garden weed in sandy areas. Inflorescence diffuse at fruiting
137	46/1.1b	Arenaria serpyllifolia subsp. lloydii		EDM 1901			1997. Now grouped with 46/1.1a 4. More robust inflorescence at fruiting. "Longis Bay". Only record. Now included under subsp. serpyllifolia
138	46/1.1c	Arenaria serpyllifolia subsp. leptoclados	Lesser Thyme-leaved Sandwort	EDM 1901	o		5 to 8 5,7,8,11. Slender, inflorescence diffuse at fruiting
139	46/3.1	Honckenya peploides	Sea Sandwort	EDM 1901	lc	4 to 8	1,2,3,4,5,13. Sand dunes
140	46/5.2	Stellaria media	Common Chickweed	CCB 1838	c	1 to 12	All
141	46/5.3	Stellaria pallida	Lesser Chickweed	ABJ 1934		3 to 5	5. (As <i>S. apetalla</i> var. <i>glabella</i>).
142	46/5.7	Stellaria graminea	Lesser Stichwort	JO 1953	vr	5 to 8	5, 6. Only other records FLOWERS 1/9/57 (voucher) FF 1967
143	46/7.3	Cerastium tomentosum	Snow-in-Summer	CMR 1950	o	4 to 9	4,7,8,9. Garden escape
144	46/7.7a	Cerastium fontanum subsp. vulgare	Common Mouse-ear	CCB 1838	c	3 to 6	Petals (5) roughly equal sepals. Hairs sparse to dense, not glandular

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 11

145	46/7.7b	Cerastium fontanum subsp. holosteoides	Common Mouse-ear	FJH 1884	c	4 to 9	1,2,3,4,5. Less hairy than above subsp. Leaves narrower
146	46/7.8	Cerastium glomeratum	Sticky Mouse-ear	EDM 1901	f	1 to 12	Petals (5) equal sepals. Hairs abundant some glandular. Sticky
147	46/7.10	Cerastium diffusum	Sea Mouse-ear	CCB 1838	c	1 to 12	Dark-green. Petals (4) shorter than sepals. Abundant glandular hair

148	46/7.12	Cerastium semidecandrum	Little Mouse-ear	CCB 1838	c	3 to 5	Petals (5) shorter than sepals.
Abundant hairs. Dries up by June							
149	46/9.1	Moenchia erecta	Upright Chickweed	EDM 1901	vr	4 to 6	5, 10. Mannez & La Tchue. 4-merous, capsule has 8 teeth
150	46/10.1	Sagina nodosa	Knotted Pearlwort	CCB 1838	o	4 to 9	3,4,5,6,7,8. Petals twice as long as sepals (5). Perennial
151	46/10.3	Sagina subulata	Heath Pearlwort	CCB 1838	c	1 to 10	Petals slightly shorter than sepals (5). Perennial
152	46/10.5	Sagina procumbens	Procumbent Pearlwort	CCB 1838	f	5 to 9	2,5,7,8,9,12. Petals minute or absent, sepals 4 or 5.. Perennial
153	46/10.7a	Sagina apetala subsp. apetala	Annual Pearlwort	EDM 1901	f	3 to 10	2,4,5,7,8. Petals minute or absent, sepals 4, erect in fruit., seed >.3mm
154	46/10.7b	Sagina apetala subsp. erecta	Common Pearlwort	CCB 1838	lf	3 to 10	7,8. As 152, sepals patent in fruit, seeds<3mm. Lf point .1-.4mm
155	46/10.8	Sagina maritima	Sea Pearlwort	CCB 1838	o	5 to 9	4,5,6,7,8,10. Similar to above 2 spp, but leaf point <.1mm
156	46/11.2	Scleranthus annuus	Annual Knawel	CCB 1838	vr	4 to 6	9,10,12,14. (Sq. 12, 1908, 1951. Refound Sq. 9 BB 1996)
157	46/13.1	Herniaria glabra	Smooth Rupturewort	CCB 1838			3,4,5,14. Records ? next species. Bibette Head JDG 1935
158	46/13.2a	Herniaria ciliolata subsp. ciliolata	Fringed Rupturewort (RDBk)	EDM 1901			7 to 8 10. Evergreen perennial, stems hairy on one side. Last seen 1956
159	46/15.1	Polycarpon tetraphyllum	Four-leaved Allseed (RDBk)	CCB 1838	lf	3 to 11	1,2,3,5,6,7,9,11,12,13, 14. (var. diphyllum recorded JDG 1937. Only record)
160	46/16.1	Spergula arvensis	Corn Spurrey	CCB 1838	lc	3 to 12	7,11,12,13. Cultivated ground. (var. nana found GCD 1907 Only record)
161	46/17.1	Spergularia rupicola	Rock Sea-spurrey	EDM 1901	c	5 to 10	2,3,6,7. Perennial. Flowers >8mm, stems glandular hairy
162	46/17.4	Spergularia rubra	Sand-spurrey	CCB 1838	lc	5 to 10	2,3,6,7,11,13,14. Old walls and sandy tracks. Ann/bien, upper bracts = leaves
163	46/17.5	Spergularia bocconei	Greek Sea-spurrey	PR 1979			11. Trois Vaux. Upper bracts much shorter than leaves
164	46/18.2	Lychnis flos-cuculi	Ragged-robin	CCB 1838	o	4 to 10	7,11,12, 13.
165	46/19.1	Agrostemma githago	Corncockle	CCB 1838	vr	5 to 8	2,7,8,12. Refound BB 1987 Platte Saline
166	46/20.2	Silene nutans	Nottingham Catchfly (Scarce)	EDM 1901	r	5 to 6	5,7, 12. Petal limb with conspicuous acute scales at base
167	46/20.4	Silene vulgaris	Bladder Campion	EDM 1901	lf	5 to 6	Scattered in many squares

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 12

168	46/20.5	Silene uniflora	Sea Campion	CCB 1838	la	1 to 12	
169	46/20.7	Silene armeria	Sweet William Catchfly	CJ 1993	vr	7 to 8	13. Garden escape or casual.
Only record							
170	46/20.9	Silene latifolia	White Campion	CCB 1838	lf	2 to 12	Flowers dioecious, calyx 10-veined male, 20-veined female
171	46/20.9 x 10	Silene x hampeana	Pink Campion	JDG 1937	c	1 to 12	Flower colour very variable
172	46/20.10	Silene dioica	Red Campion	CCB 1838	lc	1 to 12	Most coastal squares
173	46/20.11	Silene gallica	Small-flowered Catchfly (Scarce)	CCB 1838	f	4 to 8,11	4,5,9,10,12,14. Petals yellowish-white
174		Silene gallica var. quinquevulnera	Variegated Catchfly	SM 1981	o	6 to 8	1, 10,11, 12. Petals with red blotches
175	46/12	Silene conica	Sand Catchfly	EDM 1901	vr	5 to 7	4,5,7,8. Very rare since late 1960s. 2 plants refound ML 1991

176	46/22.1	Saponaria officinalis	Soapwort	BHR 1956				No site. Only record
177	46/24.1	Petrohragia nanteuillii	Childing Pink	OJ 1985				No site. Only record
178	46/25.7	Dianthus armeria	Deptford Pink (Scarce)	EDM 1900				13. Only other record ABJ

1932 (photo taken in Jersey)

Polygonaceæ

179	47/1.9	Persicaria amphibia	Amphibious Bistort	CCB 1838	lc	6 to	1,2,4,5,7. Wet places and ponds
180	47/1.11	Persicaria maculosa	Redshank	CCB 1838	lc	6 to 11	Cultivated ground. Common garden weed
181	47/1.12	Persicaria lapathifolia	Pale Persicaria	EDM 1901		o	7 to 11 7,10,11,12. Cultivated ground
182	47/1.14	Persicaria hydropiper	Water-pepper	EDM 1901		7 to 9	7. Possibly the only record
183	47/3.1	Fagopyrum esculentum	Buckwheat	(photo taken in Jersey)		EDM 1901	Rochers

Hill & Reuter's valley. Only records

184	47/4.2	Polygonum oxyspermum	Ray's Knotgrass	WH 1881			4,7. Platte Saline & Corblets.
-----	--------	--------------------------------------	-----------------	---------	--	--	--------------------------------

Only records

185	47/4.3	Polygonum arenastrum	Equal-leaved Knotgrass	ABJ 1932		vlc	6 to 10 6,10,13,14.
-----	--------	--------------------------------------	------------------------	----------	--	-----	---------------------

(Possibly others). Forming dense mat to 30cm across, common on sandy tracks.

186	47/4.4	Polygonum aviculare	Knotgrass	CCB 1838	c	6 to 11	Robust, spreading to 2m. Sandy areas. Frequent garden weed
187	47/5.1	Fallopia japonica	Japanese Knotweed	DMcC 1957	r	6 to 10	3,7,8. Garden escape. Refound BB 1987
188	47/5.3	Fallopia baldschuanica	Russian Vine	BB 1987	r	6 to 11	7,8,9. Persisting garden escape. Old buildings
189	47/5.4	Fallopia convolvulus	Black Bindweed	CCB 1838	o	6 to 11	7,9,11,12. Sandy areas, cultivated land and gardens
190	47/7	Rheum x hybridum	Rhubarb	DMcC 1963			Probably the only record

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 13

191	47/8.1	Rumex acetosella	Sheep's Sorrel	CCB 1838	la	4 to 11	
192	47/8.3	Rumex acetosa	Common Sorrel	CCB 1838	c	4 to 9	
193	47/8.10	Rumex hydrolapathum	Water Dock	BB 1987	1 site	7 to 9	4. Only record
194	47/8.13	Rumex crispus	Curled Dock	CCB 1838	c	5 to 7	
195	47/8.13 x 18	Rumex x pseudopulcher	A hybrid Dock	DMcC 1953			Only record
196	47/8.14	Rumex conglomeratus	Clustered Dock	EDM 1901	o	3 to 8	4,7. Moist places
197	47/8.15a	Rumex sanguineus var. viridis	Wood Dock	EDM 1901	f	5 to 9	4,7,8,11,12,13.

Green-veined. Woods and hedges, damp places

198	47/8.15b	Rumex sanguineus var. sanguineus	Wood Dock	BB 1987	o	5 to 9	4,5,7,8,11,12,13.
-----	----------	--	-----------	---------	---	--------	-------------------

Red-veined. not previously separated from var. viridis

199	47/8.16	Rumex rupestris	Shore Dock (RDBk)	DMcC 1958		4-8, 11	Only record
200	47/8.18	Rumex pulcher	Fiddle Dock	EDM 1901	f	4-8, 11	All squares. Widespread in small numbers
201	47/8.19	Rumex obtusifolius	Broad-leaved Dock	CCB 1838	c	4 to 8	

Plumbaginaceæ

202	48/1.1	Limonium vulgare	Common Sea-lavender	AS 1965	vr	7 to 8	8. Garden escape. Refound BB 1987
203	48/1.4	(Limonium auriculæ-ursifolium)	Broad-leaved Sea-lavender				(as <i>S. lychnidifolia</i>)

Alderney plants now all referred to next sp.

203	48/1.5	Limonium normanicum	Alderney Sea-lavender	CRPA 1900	vr	5 to 8	5. 5 to 7 leaf veins.
-----	--------	-------------------------------------	-----------------------	-----------	----	--------	-----------------------

One colony, variable numbers, spreading in area 1996-99

204	48/1.7	Limonium binervosum	Rock Sea-lavender	WMR 1842	r	5 to 8	5,6,10. 1 to 3 leaf veins.
-----	--------	-------------------------------------	-------------------	----------	---	--------	----------------------------

(Finder Rev W.M.Rogers)

205	48/2.1	Armeria maritima	Thrift	CCB 1838	la	3 to 1	Occasionally found <u>with white flowers</u>
-----	--------	----------------------------------	--------	----------	----	--------	--

206.1 48/2.2 [Armeria arenaria](#) Jersey Thrift (photo taken in Jersey) FLOWERS 1984

FLOWERS was an expert on Jersey flora, but probably an error. Only record

Clusiaceae Formerly Hypericaceae

[207](#) 51/1.3 [Hypericum androsæmum](#) Tutsan NJ 1974 vr 6 to 11 7,8,9,11. Fontaine David 9.9.74
[208](#) 51/1.9 [Hypericum tetrapterum](#) Square-stalked St. John's-wort CCB 1838 lf 4 to 10
6,7,8,11,12,13. Wet places
[209](#) 51/1.10 [Hypericum humifusum](#) Trailing St. John's-wort CCB 1838 o 5 to 9
5,6,7,10,11,12,13,14

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 14

[210](#) 51/1.11 [Hypericum linariifolium](#) Toadflax-leaved St. John's-wort (RDBk) CRPA 1899 1 site 6 to 8
10,11,12,13,14. The single remaining colony thriving. 56 plants in 1995
[211](#) 51/1.12 [Hypericum pulchrum](#) Slender St. John's-wort CCB 1838 vr 5 to 9 6,7,10,12
[212](#) 51/1.16 [Hypericum elodes](#) Marsh St. John's-wort CCB 1838 11. Last record 1901 (photo
taken in Jersey)

Tiliaceae

213 52/1.1 [Tilia platyphyllos](#) Large-leaved Lime (Scarce) CH 1987 vr 6 to 7 Planted. 1 near stream above
St. Vignalis' Garden
214 52/1.1 x 2 [Tilia x europea](#) Lime CH 1987 vr 6 to 7 Planted
215 52/1.2 [Tilia cordata](#) Small-leaved Lime RW/CH 1987 r 6 to 7 Planted
215a [Tilia heterophylla](#) (?) RW/CH 1987 vr 6 to 7 Planted, possible species for a specimen in the Churchyard or
perhaps a mistake in reporting for 213 at same site and time

Malvaceae

216 53/1.1 [Malva moschata](#) Musk-mallow EDM 1901 vr 7,9,10. Last record, 2 plants Blaye
AB/AKJ 1932
[217](#) 53/1.3 [Malva sylvestris](#) Common Mallow CCB 1838 c 5 to 12
[218](#) 53/1.7 [Malva neglecta](#) Dwarf Mallow CCB 1838 vr 6 to 9 4,5,8,10. Refound JDG 1937
[219](#) 53/2.1 [Lavatera arborea](#) Tree-mallow CCB 1838 o 5 to 11 1,2,3,4,5,7,8,10. Specimen to 94" high, Bray
Hill 1989

Cistaceae

[220](#) 56/1.1 [Tuberaria guttata](#) Spotted Rock-rose CCB 1838 2 sites 4 to 7 12,13. 1995 over c. 200 yds of cliffs.
Drops petals by noon

Violaceae

221 57/1.1 [Viola odorata](#) Sweet Violet RW/BB 1987 vr 9 to 6 9. In the Ash wood in Water Lanes
[222](#) 57/1.4 [Viola riviniana](#) Common Dog-violet EDM 1901 a 1 to 12
223 57/1.6 [Viola canina](#) Heath Dog-violet CCB 1838 Possibly in error for previous species. Only
record

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 15

[224](#) 57/1.13 [Viola arvensis](#) Field Pansy EDM 1901 o/vla 5 to 10 7,8,11,12,13. Cultivated ground

Tamaricaceae

[225](#) 58/1.1 [Tamarix gallica](#) Tamarisk EDM 1901 1 to 12 2,3,4,5,7,8. Scattered in old hedges. Some over
100 yrs old

Frankeniaceæ

226 *Frankenia laevis* Sea Heath (Scarce) JNCW 1956 7 to 8 **Only record**

Cucurbitaceæ

227 **62/5.1 (Stace)** *Cucurbita pepo* Marrow NJ 1974 **Waste heap at Banquage. Probably the only record**

227a **60/2.1** [Ecballium elaterium](#) Squirting Cucumber BB 2010 vr 8-9 5 Self-sown in a garden along the E. coast path

Salicaceæ

228 61/1.1 [Populus alba](#) White Poplar EDM 1901 f 2 to 3 Scattered. Suckers freely. Much planted by States 1990-95

229 61/1.1 x 2 [Populus x canescens](#) Grey Poplar ABJ 1933 o 3 4,5,9,13. Up to 12.5m in Val du Saue

230 61/1.3 [Populus nigra](#) Black Poplar EDM 1901 o 2 to 5 8,10. Up to 22m. 11 pollarded specimens by Old Barn

231 **61/1.3 x del** [Populus x canadensis](#) Hybrid Black-poplar ? vr 2 to 5 7,8,10. Another record, Churchyard BB 1989

232 **61/1.5** [Populus candicans](#) Balm of Gilead OJ 1987 o 3 7,8,10. Up to 9m. Some seedlings found. Suckers freely

233 **61/2.1** *Salix pentandra* Bay Willow BB 1987 vr 6 to 7 5. Twigs red-brown, hairless, glossy

234 61/2.2 [Salix fragilis](#) Crack Willow CCB 1838 o 4 to 5 6,7,8,10,11,13. Twigs very brittle

235 **61/2.3** [Salix alba](#) White Willow BB 1987 vr 4 to 7 4,5,7,9,10,13. Twigs usually yellowish-reddish, downy when young

236 **61/2.5** [Salix purpurea](#) Purple Willow EDM 1901 vr 3 to 5 10. Twigs hairless, usually purplish-brown

237 **61/2.6** [Salix daphnoides](#) Violet Willow MM 1986 vr 2 to 5 10. Twigs violet-brown. glossy with dense whitish bloom

238 61/2.9 [Salix viminalis](#) Osier EDM 1901 r 2 to 4 4,5,7,10. Twigs yellow-brown

239 **61/2.9 x 10 x 12** *Salix x stipularis* Eared Osier CCB 1838 **"Assisted by W. Borer". No site. Only record**

240 **61/2.9 x 10** *Salix x sericans* Broad-leaved Willow EDM 1901 7. Leaves densely grey pubescent underneath

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 16

241 **61/2.9 x 11** *Salix x smithiana* Silky-leaved Osier EDM 1901 vr 3 to 4 5,7,8,9. Refound RW 1987.ML near Cat's Bay 8/2001

242 **61/2.10** *Salix caprea* Goat Willow EDM 1901 **7,8,12. Mouriaux & Reuter's Valley. Only other record Vau Renier CMR**

243 **61/2.11** *Salix cinerea* Grey Willow EDM 1901 3 to 4 **Only record, probably in mistake for next species**

244 61/2.11b [Salix cinerea subsp. oleifolia](#) Rusty Sallow (**photo in seed**) CMR 1949 lc 3 to 4
Commonest sp. in Alderney. Leaves. with stiff rust-coloured hairs below

245 **61/2.12** *Salix aurita* Eared Willow CCB 1838 **12. "Not present in other C.Is". in 1838. Last record 1901**

246 *Salix babylonica* Weeping Willow CH 1987 o 3 to 4 8. Probably hybrid with *S. alba*. Planted in gardens

247 [Salix matsudana](#) Contorted Willow RW 1986 vr 3 to 6 7,13. Planted Val du Saue & Platte Saline

Brassicaceæ Formerly Cruciferæ

247a [Cleome sesquiorialis](#) Spider Plant BB 2009 1st record 8-9 5 About 20 plants in disturbed quarry spoil

248 **62/1.6** [Sisymbrium orientale](#) Eastern Rocket CMR 1953 6 to 8 **13. (As *Erysimum orientale*.) a few plants Val du Saue. Only record**

250	62/1.7	Sisymbrium officinale	Hedge Mustard	CCB 1838	f	4 to 12	3,7,9,10,13.
251	62/3.1	Alliaria petiolata	Garlic Mustard	CCB 1838	o	3 to 8	3,7,8,9,10,13.
252	62/4.1	Arabidopsis thaliana	Thale Cress	EDM1902	r	4 to 6	3,7,8,14.
253	62/7.2	Erysimum cheiri	Wallflower	FLOWERS 1960	r	1 to 12	5,8. Top of old walls in town
255	62/8.1	Hesperis matronalis	Dame's-violet	AS 1965	vr	5 to 7	Mostly a garden weed. Atlas 1962. Only records
256	62/9.1	Malcolmia maritima	Virginia Stock	DMcC 1973	vr	5 to 7	Garden escape
257	62/10.1	Matthiola incana	Hoary Stock (RDBk)	EV 1929	lc	1 to 12	1,2,3,4,5,7,10. <u>With white flowers</u> only, at Corblets Quarry
259	62/11.3	Barbarea intermedia	Medium-flowered Winter-cress	DMcC 1963		5 to 8	9
260	62/11.4	Barbarea verna	American Winter-cress	EDM 1901	vr	1 to 12	7,8,9,12. Nowadays only a garden escape
261	62/12.1	Rorippa nasturtium-aquaticum	Water-cress	CCB 1838	lf	5 to 12	5,7,8,11,12,13.
262	62/12.1 x 2	Rorippa x sterilis	Hybrid Water-cress	DMcC 1958	r	5 to 9	7
263	62/12.2	Rorippa microphylla	Narrow-fruited Water-cress	DMcC 1958	r	5 to 8	7. (Refound BB 1993. Clonque Quarry)
264	62/13.1	Armoracia rusticana	Horse-radish	CCB 1838	o	5 to 10	4,7,10.

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 17

266.1	62/14.5	Cardamine pratensis	Lady's Smock, Cuckooflower			BB 2002	1 site	4	7. In gravel in wet stream bed at Ladysmith. Patch c. 3 x 5m. 100+ plants
265	62/14.7	Cardamine flexuosa	Wavy Bitter-cress	EDM 1901					7,8,9,10,13. Claimed refound FH 1981, no voucher. Refound BB 15.8.09
266	62/14.8	Cardamine hirsuta	Hairy Bitter-cress	EDM 1901	c		2 to 8		
267	62/15.6	Arabis hirsuta	Hairy Rock-cress	CCB 1838	f		2 to 8	4,5,8,9.	
268	62/16.1	Aubrieta deltoidea	Aubretia	BB 1987	r		1 to 8	4,10.	Garden escape
269	62/17.1	Lunaria annua	Honesty	DMcC 1957			3 to 6	8,9,10.	
270	62/19.1	Bertoroa incana	Hoary Alison	FJH 1884					Braye Bay. Possibly the only record, Det. TGC 25.11.1988
271	62/20.1	Lobularia maritima	Sweet Alison	WH 1881	a		1 to 12		All squares (but rare on cliffs)
272	62/22.2	Erophila verna	Common Whitlow-grass	EDM 1901	lf		3 to 4	4,5,7,8.	
273	62/23.3	Cochlearia officinalis	Common Scurvy-grass	EDM 1901	r		3 to 6	2,3,4,5,6,7,12,13.	Rare in each sq.
274	62/23.5	Cochlearia danica	Danish Scurvy-grass	CCB 1838	c		2 to 7		Coastal only
275	62/25.1	Capsella bursa-pastoris	Shepherd's-purse	CCB 1838	o		1 to 12		Scattered plants
276	62/27.1	Teesdalia nudicaulis	Shepherd's-purse	EDM 1901	lc		2 to 4	4,5,10,11,12,13.	
277	62/28.1	Thlaspi arvense	Shepherd's-purse	BB 1987	vr		5 to 7	12,13	
278	62.30.1	Lepidium sativum	Shepherd's-purse	EDM 1903					Butes Hill. Only record
279	62/30.3	Lepidium heterophyllum	Shepherd's-purse	EDM 1901					Fort Tourgis Only other records AB/AKJ 1933 'increased'. 1 plant 1958
281	62/30.5	Lepidium ruderae	Narrow-leaved Water Pepper			EDM 1901			2, Crabby Bay 20 plants 1899 & 1900. Only records
280	62/30.8a	Lepidium draba subsp. draba	Hoary Cress	EDM 1901	r		3 to 6	4,7.	By Hammond Memorial & Platte Saline jct Picaterre
283	62/31.1	Coronopus squamatus	Swine-cress	CCB 1838	o		5 to 7	7,8.	Stem glabrous. Petals longer than sepals
284	62/31.2	Coronopus didymus	Lesser Swine-cress	EDM 1901	c		1 to 12		Vigorous garden weed. Strong smell when crushed. Petals shorter or absent
285	62/33.1	Diplotaxis tenuifolia	Perennial Wall-rocket	CCB 1838	la		4 to 1	1,2,3,4,7.	Branched leafy stem
286	62/33.2	Diplotaxis muralis	Annual Wall-rocket	EDM 1901	vo		6 to 10	3,7,8.	Stem only leafy near base

308	69/6.1	Anagallis tenella	Bog Pimpernel	CCB 1838	vr	5 to 8	11,12. Over 50m. stretch in Baxter's Valley 1996
309	69/6.2a	Anagallis arvensis subsp. arvensis	Scarlet Pimpernel (<u>photo blue form</u>)	CCB 1838	c	1 to 12	Several colour forms, <u>photos pale, flesh, red</u> , and blue. Flowers close in afternoon
310	69/6.2b	Anagallis arvensis subsp. foemina	Blue Pimpernel	EDM 1901	vr	12. Recorded as var. azure. Others CMR 1950, AS 1955, AT 1981	
311	69/6.3	Anagallis minima	Chaffweed	EDM 1901		6 to 7	7,12. ABJ 1934, CMR 1954. Only other records
312	69/7.1	Glaux maritima	Sea-milkwort	EDM 1901	r	6 to 10	5. Rock ledges nr. Limonium (and on Burhou)
313	69/8.1	Samolus valerandi	Brookweed	CCB 1838	o	4 to 8	4,5,11,12,13.

Pittosporaceae

313a	70/1.1	Pittosporum crassifolium	Karo	BB, JG	Probably planted . 2 specimens about ½-mile apart	vr	8.,9. In Memorial Gardens Victoria St.& by railway track near school
------	--------	--	------	--------	---	----	--

Hydrangeaceae

314	71/3	Hydrangea macrophylla	Hydrangea	BB 1987	vr	5 to 2	8,13. Garden escape
-----	------	---------------------------------------	-----------	---------	----	--------	---------------------

Grossulariaceae

315	72/1.1	Escallonia macrantha	Escallonia	JCNW 1956	o	1 to 12	2,4,5,7,8,10. Much used as hedging. self-seeds regularly
316	72/2.3	Ribes nigrum	Black Currant	BB 1988	vr	3 to 5	4,9, Longis Road among pines. Berry's quarry
317	72/2.4	Ribes sanguineum	Flowering Currant	BB 1988	vr	3 to 6	6,10. Bastion of Fort Clonque. Longis Bay. Also self-seeds in gardens
318	72/2.7	Ribes uva-crispa	Gooseberry	MMcW1962	vr	2 to 5	8. Ash wood in Water Lanes BB 1987

Crassulaceae

319	73/1.3	Crassula helmsii	New Zealand Pigmy-weed	ML 1983	vlf	3 to 9	4,5,10. Tank on Essex, Mannez & Corblets. New Bailiwick record
320	73/1.5	Crassula pubescens	Jersey Pigmy-weed	BB 1991		5 to 8	5. Only record, possibly an error
321	73/2.1	Umbilicus rupestris	Navelwort	CCB 1838	a	4 to 9	Up to 90cm at Val du Saue
322	75/4 (Stace)	Aeonium x velutinum		BB 1991	vr	5 to 9	5, Determined DG, Kew 1996

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 20

323	73/5.4	Sedum spectabile	Butterfly Stonecrop	MM 1987	vr	8 to 10	7. Garden escape
324	73/5.5	Sedum telephium	Orpine	EDM 1901			4. Only record
325	73/5.?	Sedum spurium	Caucasian-stonecrop	DMcC 1963			Only record
326	73/5.12	Sedum acre	Biting Stonecrop	CCB 1838	c	5 to 9	Seems to prefer coastal alkaline sandy areas
327	73/5.14	Sedum album	White Stonecrop	DMcC 1953	o	5 to 8	2,4,5,6,7,8,9,10. Usually in small quantity at site
328	73/5.16	Sedum anglicum	English Stonecrop	CCB 1838	c	5 to 8	2-9,13. Seems to prefer slightly acid soils

Saxifragaceae

329	74/3.1	Bergenia crassifolia	Elephant-ears	BB 1987	vr	11 to 8	4. Garden escape
330	74/5.8 x 9	Saxifraga x urbium	Londonpride	DMcC 1963	vr	5 to 7	Garden escape. Only record
331	74/5.19	Saxifraga tridactylites	Rue-leaved Saxifrage	EDM 1901	lf	3 to 5	4,5,8.

Rosaceae

332	75/6.1	Filipendula vulgaris	Dropwort	DMcC 1963		6	Only record
333	75/8.7	Rubus idaeus	Raspberry	EDM 1901		10.	In GCD Comital Flora 1932 'Alderney'

334	75/8.321	Rubus caesius	Dewberry	EDM 1901	c	4 to 8	All leaves in 3s. Flowers white, earliest. Few large drupelets in fruit
335		Rubus fruticosus agg.	Blackberry. (Many subspecies)	CCB 1838	a	5 to 11	Most leaves in 5s. flowers pinkish, later. Fruits more smaller drupelets
336	77/8.14 (Stace)	Rubus series rhamnifolia	Blackberry	CCB 1838			
337		Rubus leucostachys	Blackberry	GCD 1906			
338	77/8.24 (Stace)	Rubus corylifolii	Blackberry	GCD 1906			
339		Rubus rusticanus	Blackberry	GCD 1907			
340	75/9.1	Potentilla fruticosa	Shrubby Cinquefoil	BB 1987	vr	6 to 7	Garden escape
341	75/9.3	Potentilla anserina	Silverweed	CCB 1838	f	5 to 10	7,11,12. Some plants silver both sides of leaf
343	75/9.13	Potentilla erecta	Tormentil	CCB 1838	c	5 to 11	2,4,5,6,7,10,11,12,13 Mainly cliffs and quarries
344	75/9.14 x 15	Potentilla x mixta	Hybrid Cinquefoil	CMR 49-63	f	6 to 9	7,11,12.
345	75/9.15	Potentilla reptans	Creeping Cinquefoil	CCB 1838	c	6 to 10	Fixed dune areas

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 21

346	75/9.16	Potentilla sterilis	Barren Strawberry	EDM 1901	o	2 to 5	7, 8,12. Next CMR 1954. Refound JB 1992
348	75/11.1	Fragaria vesca	Wild Strawberry	BB 1997	vr	6 to 7	8, New churchyard in wall at N end in flower & fruit 13.7.97
347	75/11.3	Fragaria ananassa	Garden Strawberry	DMcC 1973	vr	2, 5 -12	Garden escape or throwout
349	75/13.2	Geum urbanum	Herb Bennet	EDM 1901	o	2 to 8	2,7,8,9. 'Damp places'
350	75/15.1	Agrimonia eupatoria	Agrimony	EDM 1901	vr	5 to 9	4, 8, 9. E. rim Battery Quarry. E. side of Les Rochers JR 1/9/57 Voucher
351	75/17.3a	Sanguisorba minor subsp. minor	Salad Burnet	JM 1959		5 to 8	2,4,9. Garden escape
352	75/20.1	Aphanes arvensis	Parsley-piert	CCB 1838	r	3 to 6	5,7,11.
353	75/20.2	Aphanes australis	Lesser Parsley-piert	JO 1953	a	2 to 5	All squares
354	75/21.1	Rosa multiflora	Many-flowered Rose	DMcC 1953	1 site	6 to 8	10. Refound BB 1989 at The Old Barn
357.1	75/21.4 x 10	Rosa x alba	White Rose of York	BB 1996	1 only	7 to 8	11, Single long established specimen on hedgebank (La Costière) on Grande Blaye above Trois Vaux
355	75/21.5	Rosa pimpinellifolia	Burnet Rose	CCB 1838	r	5 to 9	3,4,12,13, on walls bounding Blaye. (Braye Common, BB1988 only).
356	75/21.5 x 16	Rosa x involuta	(R. pimpinell. x R. sherardii)	EDM 1901			10. Only record
357	75/21.6	Rosa rugosa	Japanese Rose	CMR 1950	r	4 to 11	5,7,8,10. Much used as hedging. Suckers and seeds
358	75/21.11	Rosa stylosa	Short-styled Field-rose	EDM 1901	o	6 to 8	A few each in 2-14. Refound JES 1971, BB 1992
359	75/21.12	Rosa canina	Dog-rose	CCB 1838	o	6 to 8	2-14, widely scattered, few in numbers
360	75/21.15	Rosa tomentosa	Harsh Downy-rose	EDM 1901			7. Only record
361	75/21.17	Rosa mollis	Soft Downy-rose	FF 1967			7 Only record
362	75/21.18	Rosa rubiginosa	Sweet-briar	EDM 1901	o	6 to 7	4,10,11,12,13,14.
363	75/21.19	Rosa micrantha	Small-flowered Sweet-briar	EDM 1901	o	6 to 8	3,4,5,7,12,13,14. Refound BB 1992
364		Rosa wicheriana x Shirley Hibberd	Alberic Barbier Rose	NVM 1967	1 site	1 to 12	7. On rim of sand pit for about 25 years. Habitat destroyed 1991. Small area recovered 1998 on fence of Tennis Courts erected over site
365		Rosa wicheriana x Gabriel Luiget	Dorothy Perkins Rose	BB 1988	4 sites	7 to 9	6,7,11,13. Vau Pommier, S. cliffs. Probably planted 30+ years ago
366		Rosa wicheriana x R. setigera	American Pillar Rose	BB 1986	2 sites	6 to 10	2,6. Bank by "Pilot's View" Crabby for 40+ years. Probably planted

391 77/8.3	Lotus pedunculatus	Greater Bird's-foot-trefoil	CCB 1838	o	5 to 10	3,4,5,7,9,11,12,13. Mostly in damp places
392 77/8.4	Lotus subbiflorus	Hairy Bird's-foot-trefoil;	CCB 1838	la	4 to 9	Cliffs and low coasts
393 77/8.5	Lotus angustissimus	Slender Bird's-foot-trefoil	CCB 1838	o	6 to 8	5,10,11,12,13,14. Refound OJ 1982. Was rare, now occasional
395 77/10.3	Ornithopus perpusillus	Bird's-foot	CCB 1838	vla	4 to 7	4,5,9,10,11,12,13,14
396 77/10.4	Ornithopus pinnatus	Orange Bird's-foot	MD 1881	lf	5 to 7	4,5,9,10,11,12,13,14
397 77/14.2	Vicia cracca	Tufted Vetch	EDM 190	vr	5 to 8	4,7,10,11,12
398 77/14.5	Vicia villosa	Fodder Vetch	EDM 1900			9. Only other records CPH 1902, ABJ 1932
399 77/14.6	Vicia hirsuta	Hairy Tare	CCB 1838	a	4 to 7	2-7fls. 2-seeded pods
400 77/14.8	Vicia tetrasperma	Smooth Tare	EDM 1901	vr	6 to 7	5,7,12,13. 1-2fls, 4-seeded pods
401 77/14.11a	Vicia sativa subsp. nigra	Narrow-leaved Vetch	EDM 1901	c	4 to 10	Plant very slender often procumbent
402a 77/14.11b	Vicia sativa subsp. segetalis	? (No details)				5. Old record found by BB 12/5/1997
402 77/14.11c	Vicia sativa subsp. sativa	Common Vetch	CCB 1838	c	1 to 11	Robust plant. Many white flowered plants at Crabby Bay 1979
403 77/14.12	Vicia lathyroides	Spring Vetch	CMR 1952	r	4 to 6	5,10,11,12. Annual to 20cm. Mannez quarry, BB June 1995
404 77/14.14	Vicia bithynica	Bithynian Vetch (Scarce)	DMcC 1963		2 sites	4 to 8 2,4. Robust plant, flowers purple & white. Present site along low cliff edge, Crabby Bay
405	Vicia faba	Broad Bean	DMcC 1963	vr		7. Garden escape
406 77/15.4	Lathyrus pratensis	Meadow Vetchling	EDM 1901	vr	6 to 8	4,9,10. Perennial. Flowers yellow, stem angled but not winged
407 77/15.7	Lathyrus grandiflorus	Two-flowered Everlasting-pea	BB 1988	vr	6 to 9	2,3,5,8. Flowers >25mm, pinkish/purple
408 77/15.9	Lathyrus latifolius	Broad-leaved Everlasting-pea	PHAS 1969	vr	6 to 9	5,8,9. Flowers 15-30mm, pink, brighter than previous sp.
409 77/15.13	Lathyrus aphaca	Yellow Vetchling (Scarce)	JES 1970	vr		Annual. Mature leaves reduced to tendrils. Refound JP & Soc G. 1986
410 77/16.2	Ononis reclinata	Small Restharrow (RDBk)	CCB 1828	vr	5 to 6	4,5. Not seen 1988-92 Refound coast MM 1993, Man. hill MK 1995, both sites BB 1996, 97, coast only 98, 99
411	Ononis repens f. albiflorus	White Small-restharrow (RDBk)	OJ 1979	vr	5 to 6	5. Refound OJ 1979, BB 29.6.1988 photo
412 77/16.4	Ononis repens	Common Restharrow	CCB 1838	c	4 to 10	Occasionally with white flowers
413 77/17.1	Melilotus altissimus	Tall Melilot	JDG 1937	r	5 to 8	2,3,4,8. Biennial/perennial. Flowers standard, wings & keel all equal

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 24

414 77/17.2	Melilotus albus	White Melilot	GCD 1906	o	6 to 9	2,3,7. Refound NVM, Crabby/Platte Saline, 1978, still there 1999
415 77.17.3	Melilotus officinalis	Ribbed Melilot	EDM 1903	lf	5 to 9	2,3,7,8. Annual, Flowers keel shorter
416 77/17.4	Melilotus indicus	Small Melilot	EDM 1899	o	5 to 10	2,3,4,5,7,8,12. Flowers standard longer
417 77/18.1	Medicago lupulina	Black Medick	CCB 1838	c	4 to 10	Annual or SLperennial. Flowers 10-50. Fruit single turn, net black when ripe
418 77/18.2c	Medicago sativa subsp. sativa	Lucerne	EDM 1901	f	4 to 10	Sandy parts. Flower colour varies from v. dark purple to white
419 77/18.4	Medicago polymorpha	Toothed Medick	CCB 1838	r	5 to 8	3,4,8,11,12. Flowers 1-5. Fruit flat, 1½-6 turns, spiny
420 77/18.5	Medicago arabica	Spotted Medick	CCB 1838	a	3 to 9	Flowers 1-4. Fruit globose spiral, 3-7 turns, spines grooved ½way

421 77/19.1	Trifolium ornithopodioides	Bird's-foot Clover	EDM 1901	o	4 to 8	4,5,6,7,10,11,12,13,14.
Flowers 1-4 in raceme, pedicels >1mm						
422 77/19.2	Trifolium repens	White Clover	CCB 1838	a	1 to 11	Flowers globose, peduncles up to 20cm, about a month later than next sp
423 77/19.3	Trifolium occidentale	Western Clover (Scarce)	RWW 1962	lc	3 to 8	Smaller, peduncles short. Cliff and dune edges usually within 100m from sea
424 77/19.4	Trifolium hybridum	Alsike Clover	BHR 1956			4,7. Flowers usually pink & white. Only other record FF 1967, Longis
425 77/19.5	Trifolium glomeratum	Clustered Clover (Scarce)	EDM 1901	o	6 to 8	4,5,13,14. Cliffs.
Flowers 4-5mm purple. 2-seeded fruit enclosed by calyx						
426 77/19.6	Trifolium suffocatum	Suffocated Clover	EDM 1901	o	3 to 6	4,5,10-14. Plant 5cm across, whitish 3mm flowers mostly near root
427 77/19.8a	Trifolium fragiferum subsp. fragiferum	Strawberry Clover	EDM 1901	vr	6 to 7	4,7,11,13. Marshy places. Pink, flowers 5-7mm, Few records
428 77/19.8b	Trifolium fragiferum subsp. bonannii	Strawberry Clover (Southern)				JH 1932
Status now considered doubtful (Stace). Only record						
429 77/19.11	Trifolium campestre	Hop Trefoil	CCB 1838	vf	1, 4 to 8	Bare places. Yellow racemes globose 10-15mm with 20-40 flowers
430 77/19.12	Trifolium dubium	Lesser Trefoil	CCB 1838	c	4 to 10	Racemes 6-8mm with 5-20 flowers
431 77/19.13	Trifolium micranthum	Slender Trefoil	CCB 1838	o	4 to 8	All except. 1.
Racemes c.5mm with 2-6 flowers						
432 77/19.14	Trifolium pratense	Red Clover	CCB 1838	c	4 to 12	Colour varies considerably pinkish-purple, pale pink, occ. cream
433 77/19.15	Trifolium medium	Zigzag Clover	JO 1953	vr	4 to 8	7,8. Rhizomatous, flowers terminal in globose red-purple raceme
434 77/19.18	Trifolium stellatum	Starry Clover	ABJ/HKAS			11. Trois Vaux 1932. Only record
435 77/19.19a	Trifolium incarnatum subsp. incarnatum	Crimson Clover	EDM 1901			
4,10, Above Hanging Rock. Only other record JDG 1937						
436 77/19.20	Trifolium striatum	Knotted Clover	CCB 1838	o	5 to 7	5,7,9,10,11,12,13,14.
Coastal. Pink racemes sessile, axillary & terminal						

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 25						
437 77/19.21	Trifolium bocconeii	Twin-headed Clover (RDBk)	DMcC 1957		5 to 6	Trois Vaux. Only record
438 77/19.22	Trifolium scabrum	Rough Clover	CCB 1838	f	5 to 8	Cliff tops, dune edges, short turf. Pubescent annual, flowers white/pale pink
439 77/19.23	Trifolium arvense	Hare's-foot Clover	CCB 1838	lf	3 to 6	Mainly cliffs & bare ground. Whitish terminal racemes, cylindrical
440 77/19.25	Trifolium subterraneum	Subterranean Clover	EDM 1901	f	3 to 8	1,5,6, & Cliffs.
Raceme axillary 2-5 whitish fertile flowers, plus sterile calices						
442 77/21.1	Lupinus arboreus	Tree Lupin	DMcC 1957	o	3 to 9	Usually in sandy places close to sea. Occasionally with white flowers
443 77/22.1	Laburnum anagyroides	Laburnum	BB 1987	o	5 to 6	8,9. Planted in gardens. 2 seedlings pavement wall bottom in QEII St.
444 77/23.4a	Cytisus scoparius subsp. scoparius	Broom	CCB 1838	r	4-7, 12	4,11,13,14. Mainly on cliff tops
445 77/23.4b	Cytisus scoparius subsp. maritimus	Prostrate Broom	M'Nab 1850	lc	3-6, 9	11,12,13,14. Cliffs SW end of island
446 77/24.1	Spartium junceum	Spanish Broom	BB 1989	vr	6 to 11	9, Derelict Glebe land on Les Rochers. Several large bushes, still there

464	Epilobium tetragonum subsp. lamyi	Square-stalked Willowherb	BB 1987	r	6 to 9	5,7. As previous
ssp but leaves with short petioles						
465 84/1.6	Epilobium obscurum	Short-fruited Willowherb	DMcC 1958	r	6, 8, 11	9,11. Refound Trois Vaux, BB 7.1989
467 84/1.8	Epilobium ciliatum	American Willowherb	BB 1987	o	6 to 9	4,5,7,8,13. Flowers 5mm pinkish-purple. Leaves rounded at base
468 84/1.9	Epilobium palustre	Marsh Willowherb	BB 1992	vr	6 to 8	5,7. Flowers 4-5mm pale pink. Leaves narrow at base
469 84/2.1	Chamerion angustifolium	Rosebay Willowherb	DMcC 1953	o	7 to 9	3,6,7,8,9,10. Small patches only
471 84/4.1	Oenothera glazioviana	Large-flowered Evening-primrose	JO 1953	vlf	6 to 11	3,8,9. Green parts of stems & fruits with red bulbous-based hairs
471a 84/4.4	Oenothera cambrica	Small-flowered Evening-primrose	ML 2005	vlf	6 to 10	1,4,7. Platte Saline on tipped soil. Sepals green. Red hairs ±
472 84/4.5	Oenothera stricta	Fragrant Evening-primrose (RDBk)	EDM 1901	r	5 to 12	1,7,8,9. Petals turn reddish on withering. No red bulbous-based hairs
472b	Gaura lindheimeri		ML 1975		6 to 9	7, near the House Holly-fern. Persisted to 1978/9
473 84/5.1	Fuchsia magellanica	Fuchsia	PHAS 1970	o	1 to 12	2,5,6,7,8,10. Vau Pommier & scattered. Det. TGT & AOC
474 84/6.1	Circaea lutetiana	Enchanter's-nightshade	EDM 1901	o	6 to 8	7,8,9,13. Petals 2, white/pale pink. Stamens 2

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 27

Santalaceae

475 86/1.1	Thesium humifusum	Bastard-toadflax	CCB 1838	o	5 to 11	1,4,5,7,8,9. Seems to be increasing again. Leaves & stem olive-green. Large area on Mannez Hill noted, BB 1996-99
------------	-----------------------------------	------------------	----------	---	---------	---

Celastraceae

476 88/1.3	Euonymus japonicus	Evergreen Spindle	FLOWERS 1956	f	4 to 6	2,4,5,7,8,9,10. Much planted as hedging, seedlings occasionally found
------------	------------------------------------	-------------------	--------------	---	--------	---

Aquifoliaceae

477 89/1.1	Ilex aquifolium	Holly	BB 1987	r	3 to 5	6,7,8,13. Self seeds in gardens. Also planted by States
------------	---------------------------------	-------	---------	---	--------	---

Euphorbiaceae

477a	Ricinus communis	Castor-oil-plant	TD 1.9.11	vr	1st record	8. Waste land in St. anne
478 91/1.2	Mercurialis annua	Annual Mercury	CCB 1838	c	1 to 12	Dioecious
479 91/2.1	Euphorbia peplis	Purple Spurge (RDBk)	JS 1838	extinct		1,2. Habitat destroyed by storm 1977. Not seen since. Possibly now extinct
480 91/2.8	Euphorbia helioscopia	Sun Spurge	CCB 1838	lf	1 to 12	Annual to 50cm.
481 91/2.9	Euphorbia lathyris	Caper Spurge	FLOWERS 1977	r	6 to 7	5,7,8. Probably garden escape. Biennial 1-2m. Pops up here and there most years.
482 91/2.11	Euphorbia peplus	Petty Spurge (close up)	CCB 1838	c	1 to 12	Annual to 30cm.
483 91/2.12	Euphorbia portlandica	Portland Spurge (Scarce)	CCB 1838	c	2 to 10	All round coast. Perennial. Stems usually reddish, leaves yellow-green
484 91/2.13	Euphorbia paralias	Sea Spurge	CCB 1838	lf	4 to 9	1,2,3,4. Dune edges. Perennial to 60cm. Leaves narrow, succulent
485 91/2.15	Euphorbia cyparissias	Cypress Spurge	BB 1987	r	1 to 12	2,7,8. 'Garden Escape'. Rhizomatous perennial. To 60cm, leaves linear.

Linaceae

486 94/1.1	Linum bienne	Pale Flax	CCB 1838	lf	3 to 11	2,3,4,5,7,9,13. Short sandy turf. Flowers 10mm, pale blue
------------	------------------------------	-----------	----------	----	---------	---

Tropaeolaceae

533 **104/1.1** [Tropaeolum majus](#) Nasturtium DMcC 1958 o 4 to 12 Garden escape

Araliaceae

534 **106/1.2b** [Hedera helix subsp. hibernica](#) Atlantic Ivy CCB 1838 a 9 to 12 Most, if not all,
Alderney plants are probably this subsp.
535 **106/2.2** [Aralia elata](#) Japanese Angelica-tree BB 1989 vr 7 to 8 8,10. Planted, Longis Road and some
gardens in town

Apiaceae Formerly Umbelliferæ

536 **107/1.1** [Hydrocotyle vulgaris](#) Marsh Pennywort CCB 1838 r 4 to 7 4,11,13. Baxter's Valley,
many plants in dried up Longis pond bed 1992
537 **107/4.3** [Eryngium maritimum](#) Sea-Holly CCB 1838 lf 6 to 9 1,2,3,4
538 **107/4.4** [Eryngium campestre](#) Field Eryngo (RDBk) NVM 1979 vr 8, below Butes Not seen
since 1986/7
539 **107/5.3** [Chaerophyllum temulum](#) Rough Chervil CCB 1838 **Only record**
540 **107/6.1** [Anthriscus sylvestris](#) Cow Parsley EDM 1901 c 2 to 12
541 **107/6.2** [Anthriscus caucalis](#) Bur Parsley CCB 1838 vr 3 to 9 5,10,12. Only records 1838 1901, 1992

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 31
542 **107/7.1** [Scandix pecten-veneris](#) Shepherd's-needle (Scarce) EDM 1901 vr 6 **4,9. Only records**
543 **107/9.1** [Coriandrum sativum](#) Coriander 1,977 vr 6 **Garden escape. Only record**
544 **107/10.1** [Smyrniolum olusatrum](#) Alexanders EDM 1901 1 site 3 to 6 8, Not seen 1986-93 Reported by DB
1994, refound BB 1996
545 **107/10.2** [Smyrniolum perfoliatum](#) (A Southern European sp.) JO 1953 'Butes
Hill, plentiful' Only record, probably error for previous species.
546 **107/14.1** [Aegopodium podagraria](#) Ground-elder NJ 1972 **Only record**
547 **107/16.1** [Berula erecta](#) Lesser Water-parsnip CCB 1838 **Only record**
548 **107/17.1** [Crithmum maritimum](#) Rock Samphire CCB 1838 c 6 to 10 5-10,12,13,14.
Most shores and lower cliffs
549 **107/19.5** [Oenanthe crocata](#) Hemlock Water-dropwort (Photo taken in Jersey) CCB 1838
Reported as "Extinct by 1860". But still put onto Mauvaises Herbes list in 1933
550 **107/20.1** [Aethusa cynapium](#) Fool's Parsley ABJ 1932 vr 8 to 9 8. 1932 'Blaye'. Refound in
churchyard BB 1992. Det. DMcC
551 **107/21.1** [Foeniculum vulgare](#) Fennel CCB 1838 la 7 to 12 Dune areas, cliff paths and N. coast
552 **107/25.1** [Conium maculatum](#) Hemlock CCB 1838 lf 5 to 10 2,4,5,7,9,10,11,12,13,14. Commons &
cliff paths
553 **107/26.4** [Bupleurum baldense](#) Small Hare's-ear CCB 1838 vlc 5 to 7 45,914
554 **107/28.1** [Apium graveolens](#) Wild Celery EDM 1903 vr 5 to 7 2,4,5,7. Refound TD 1989
555 **107/28.2** [Apium nodiflorum](#) Fool's Water-cress CCB 1838 c 6 to 10 1,2,4,5,6,7,11,12,13
556 [Apium nodiflorum var. pseudorepens](#) Fool's Water-cress (prostrate form) CCB 1838 Single site
7 to 8 5. Refound outlet Mannez pond BB 1992. Det. D.Meikle
557 **107/29.1** [Petroselinum crispum](#) Garden Parsley CCB 1838 vr 5 to 11 4,8. Garden escape
and near old ruins 5. Path down to Houmet Herbé from road
558 **107/29.2** [Petroselinum segetum](#) Corn Parsley (RDBk) BB 1990 vr 6 to 8
559 **107/33.1** [Falcaria vulgaris](#) Longleaf BHR 1956 vr 7 to 9 9,13. Refound TD, ¼ acre on Rond But
1988
560 **107/40.1** [Pastinaca sativa var. sylvestris](#) Wild Parsnip CCB 1838 lf 5 to 8 4,5,7,8,9. Pinnate
leaves large flflets. Petioles long, wavy. No sepals
561 **107/41.1** [Heracleum sphondylium](#) Hogweed CCB 1838 a 1 to 12 On Mauvaises Herbes Law
list. Spreading greatly

562	Heracleum sphondylium var. angustifolium	Hogweed	DMcC 1974	vr				10,11. Very large leaves, by entrance to German tunnel BB 1993, conf. DMcC. Giffoine 1997. Only records
563 107/43.1	Torilis japonica	Upright Hedge-parsley	CCB 1838	vr			4,10. Erect 1.2m	Fruit spines curved Other records, 1906, 1980,95
564 107/43.3	Torilis nodosa	Knotted Hedge-parsley	CCB 1838	o	5 to 8		4,5,9,10,11.	Procumbent 50cm Flowers white. Fruit spines straight. No bracts
564a	Torilis arvensis	Spreading Hedge-parsley		vr 1st record	5 to 8		Verge in Val Fontaine ? garden escape	

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 32

565 107/44.1a	Daucus carota subsp. carota	Wild Carrot	CCB 1838	f	4 to 8		4,5,6,7,9,10,13. Umbel	concave in fruit
567 107/44.1b	Daucus carota subsp. sativus	Carrot	BB 1995		6 to 8		13. Relic of	cultivation on Blaye. Large patch. Only record
566 107/44.1c	Daucus carota subsp. gummifer	Sea Carrot	CCB 1838	c	1 to 12		1,8,9,12,14. Dunes,	cliff tops. Umbel convex or flat in fruit

Gentianaceæ

568 108/3.2	Centaurium erythraea	Common Centaury	CCB 1838	c	6 to 11		Cliffs and commons. Biennial	to 50cm
569 108/3.4	Centaurium pulchellum	Lesser Centaury (RDBk)	DMcC 1958		1 site	5 to 8	5. Annual, usually	c.5cm. Damp ground nr. sea

Apocynaceæ

570 109/1.3	Vinca major	Greater Periwinkle	EDM 1901	r	1 to 12		2,3,4,5,7,8,9,10. Leaf & calyx margins pubescent	
---	-----------------------------	--------------------	----------	---	---------	--	--	--

Solanaceæ

571 110/1.1	Nicandra physalodes	Apple-of-Peru	KEB 1969	BB 2008	vr		Masses in Mannez Quarry spoil heap 24.10.08	& 09 6 to 11 5, 7. Refound 1.10.88 BB. On Platte Saline in quantity after soil tipped. Occasionally in same area since.
572 110/2.1	Lycium barbarum	Duke of Argyll's Teaplant	WH 1881		o	4 to 12	5,7,8,9. Thick clumps along	N. coast . Leaves. elliptic. Corolla 7-12mm
573 110/2.2	Lycium chinense	Chinese Teaplant	EDM 1908		o	3 to 12	2-4,6-9. Leaves lanceolate/narrow	ovate, fl. larger. Corolla 12-17mm
574 110/4.1	Hyocyanus niger	Henbane	CCB 1838	vr	6 to 8		2,3,4,7 Recent records. 1 plant; NVM 1979,	Berry's Quarry BB 1990
575 110/7.1	Lycopersicon esculentum	Tomato	RSRF 1956	vr	5 to 9		Occasionally on rubbish tips	
576 110/8.1	Solanum nigrum	Black Nightshade	EDM 1901	lf	6 to 12		3,4,5,7,8,9,10. Frequent garden weed	
577	Solanum nigrum var. atriplicifolium		ABJ 1932				Specimen in BM. Only record	
578 110/8.4	Solanum sarachoides	Leafy-fruited Nightshade		HCP 1969			Only other record	1973
579 110/8.6	Solanum dulcamara	Bittersweet	CCB 1838	f	5 to 10		Usually in hedges	
580 110/8.7	Solanum tuberosum	Potato	RSRF 1956	vr	5 to 9		Rubbish tips. Refound BB Platte Saline 1988 &	1993
581 110/9.1	Datura stramonium	Thorn-apple	GJ pre 1940	BB 2008, 2009			6 to 9 7,9. Disturbed old ground. Refound BB 1993 and 2009	vr Masses in Mannez Quarry spoil heap 24.10.08
581a	Datura stramonium var. tatula	A Thorn-apple		BB 2001		Single site	6 to 9 9. Spread soil by	Railway Station. More in 2002

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

582	Nicotiana rustica	Wild Tobacco	OJ 1976					8. Probably garden escape. Only record
583	Nicotiana glauca	Sweet Tobacco	BB 1988					9. Probably garden escape. Only record

Convolvulaceæ

584	111/1.1	Convolvulus arvensis	Field Bindweed	CCB 1838	c	5 to 11	Found with deep pink flowers nr. Whitegates, occ. pure white
585	111/3.1	Calystegia soldanella	Sea Bindweed	CCB 1838	lc	6 to 10	1,2,3,4,5. Mainly N. coast dunes
586	111/3.2	Calystegia sepium	Hedge Bindweed	EDM 1901	lc	6 to 11	1,4,8,12. Bracteoles 10-20mm, don't overlap
587	111/4.1	Calystegia silvatica	Large Bindweed	DMcC 1953	vlc	6 to 10	Bracteoles 20-40mm, overlap strongly

Cuscutaceæ

589	112/1.1	<i>Cuscuta campestris</i>	Yellow Dodder	EDM 1901			(As <i>C. trifolii</i> , but his description agrees with this sp.) Also CMR 1950. No other records
588	112/1.3	Cuscuta epithymum	Dodder (<u>Photo yellow form</u>) see <i>C. campestris</i>	CCB 1838	lc	6 to 9	Here usually grows on ground on Thyme in 1-6,8,9. Much rarer on Gorse in 12,13,14. Patch of Yellow form of <i>C. epithymum</i> on shore nr. Houmet Herbé BB 8.02. Determined CAS

Menyanthaceæ

590	113/2.1	Nymphoides peltata	Fringed Water-lily (Scarce)	BB 1986	vr	6 to 8	7,10, (Tank on Essex Hill, gone in 1996).
-----	---------	------------------------------------	-----------------------------	---------	----	--------	---

Hydrophyllaceæ

591	115/1.1	Phacelia tanacetifolia	Phacelia	BB 1994	vr	6 to 7	12, (Remnant of planted fodder crop) 7, (1998) in Bracken on Les Rochers
-----	---------	--	----------	---------	----	--------	--

Boraginaceæ

592	116/1.3	<i>Lithospermum arvense</i>	Field Gromwell	EDM 1900	vr		2. Only other record 1971. Not seen since
593	116/2.1	Echium vulgare	Viper's-bugloss	CCB 1838	lc	5 to 11	1,2,5,6,8,11,12,13. Spreading recently
594	116/2.2	Echium plantagineum	Purple Viper's-bugloss	CMR 1956	vr	6 to 7	4,7. DMcC doubtful of 1st record, but 5 plants P/Saline BB 1996
595	116/2.4	Echium pininana	Giant Viper's-bugloss	DMcC 1973	r	4 to 9	2,5,7,8,10. Frequently grown in gardens. Occasional escape

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

596	116/4.1	Symphytum officinale	Common Comfrey	EDM 1901	vlf	5 to 8	1,2,6,14. Stem leaves strongly decurrent >1 internode
597	116/4.1 x 2	Symphytum x uplandicum	Russian Comfrey	ABJ 1933	lf	3 to 9	1,2,5,6,7,9,14. The more common species. More bristly. Intermediate hybrids likely. Leaves clasping or slightly decurrent
598	116/4.3	Symphytum tuberosum	Tuberous Comfrey	CMR 1954	vr	3 to 8	7,8,12. RSRF 1956. Refound in Churchyard BB 1995
598a		Symphytum grandiflorum	Creeping Comfrey	BB 9.5.2010	vr	1st record	5 to 7 Verge in Val Fontaine
599	116/6.4	Anchusa arvensis	Bugloss	CCB 1838	f	2 to 12	4,5,11,12,13. Annual to 50cm. Hispid. Calyx divided nearly to base
600	116/8.1	Pentaglottis sempervirens	Green Alkanet	BB 1987	vr	2 to 8	3,4,5,11. Perennial to 1m. Hispid perennial. Calyx divided ¾ way
601	116/9.1	Borago officinalis	Borage	CCB 1838	o	1 to 12	1,3,4,6,10,12. Well scattered

602 116/15.2	Myosotis secunda	Creeping Forget-me-not	CCB 1838	vlf	5 to 8	7,11. Perennial
with stolons. Long pedicels & patent hairs below						
603 116/15.4a	<i>Myosotis laxa</i> subsp. <i>caespitosa</i>		BB 1988	r	7	11, Trois Vaux in scree at jct. Baxter's Valley 5.7.88
604 116/15.7	<i>Myosotis sylvatica</i>	Wood Forget-me-not	DMcC 1957		5 to 6	Possibly the only record
605 116/15.8	Myosotis arvensis	Field Forget-me-not	CCB 1838	o	3 to 9	6,8,13. Plant greyish-green to 40cm. Corolla 3mm.
606 116/15.9	Myosotis ramosissima	Early Forget-me-not	EDM 1901	c	2 to 6	All squares ex 8. Calyx with hooked hairs. Fl blue 3mm Inflorescence elongates greatly after fertilisation
607 116/15.9	Myosotis ramosissima subsp. lebelii	Early Forget-me-not	EDM 1901	o	2 to ?	11,13. White flowers with yellow throat. Refound BB 29.4.1988
608 116/15.10	Myosotis discolor	Changing Forget-me-not	EDM 1901	lc	3 to 8	All squares ex 8. Erect c.10cm. corolla 2mm, yellow first, then blue. Inflorescence elongates after fertilisation.
609 116/17.1	Cynoglossum officinale	Hound's-tongue (photo taken in Jersey)	CCB 1838			
2. Only other record, Crabby CMR 1955						

Verbenaceae

610 117/1.1	Verbena officinalis	Vervain	CCB 1838	vr	6 to 10	8,10
---	-------------------------------------	---------	----------	----	---------	------

Lamiaceae Formerly Labiatae

611 118/1.5	Stachys sylvatica	Hedge Woundwort	EDM 1901	f	5 to 11	4,7,8,11,12,13. Usually in damper spots. Perennial. Leaves petiolate.
612 118/1.5 x 6	<i>Stachys x ambigua</i>	Hybrid Woundwort	GCD 1932	?		In GCD 1932 'S'. In WFG Supplement DMcC 1983
613 118/1.6	Stachys palustris	Marsh Woundwort	CCB 1838	o	7 to 9	9,12,13,14. Perennial. Upper & middle stem leaves sessile

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 35

614 118/1.8	<i>Stachys arvensis</i>	Field Woundwort	CCB 1838	o	5 to 10	7,12,13. Annual. No glandular hairs
615 118/2.1	Ballota nigra	Black Horehound	CCB 1838	r	6 to 10	2,8,9. Waste ground
616 118/4.1c	Lamium purpureum subsp. argenteum	Yellow Archangel	JM 1959		2 sites	4 to 7 7,8. Refound BB 1993, Churchyard & St. Vignalis' Garden
617 118/5.3	Lamium purpureum	Red Dead-nettle	EDM 1901	c	1 to 12	All leaves petiolate, serrate, teeth <2mm
618 118/5.4	Lamium hybridum	Cut-leaved Dead-nettle	CCB 1838	o	1 to 12	5,8,12,13. As above, many teeth >2mm
619 118/5.6	<i>Lamium amplexicaule</i>	Henbit Dead-nettle	CCB 1838	r	4 to 8	7,8,10. Scattered. Middle & upper lf. subtending whorls sessile
620 118/6.4	Galeopsis tetrahit (Photo taken in Guernsey)	Common Hemp-nettle	EDM 1901	vr		8. "1 plant 8/1901". Refound JO 1953. Only other records FLOWERS 1957, (voucher) AS 1960
621 118/9.1	Marrubium vulgare	White Horehound (Scarce)	CCB 1838	r	6 to 11	13,479
622 118/11.1	Teucrium scorodonia	Wood Sage	CCB 1838	a	3 to 10	All round coast and many inland hedges etc.
623 118/13rac x nep	Nepeta x faassenii	Garden Cat-mint	JES 1972	o	5 to 9	3,4,5,7,8. Probably garden escape
624 118/14.1	Glechoma hederacea	Ground-ivy	CCB 1838	vlf	3 to 9	6,7,9,10,11. Calyx 5 teeth Stolons c. 1m. with petiolate leaves.
625 118/15.1	Prunella vulgaris	Selfheal	CCB 1838	f	6 to 9	5,6,7,8,9,10,11,13. Calyx 3 short teeth.
626 118/15.2	<i>Prunella laciniata</i>	Cut-leaved Selfheal	RAEB 1932			14, Upper leaves divided to midrib. Only other record CMR 1951

627	118/16.1	Melissa officinalis	Balm	DMcC 1956	vr	6 to 11	7,9	Only other record BB 1987
628	118/18.4	Clinopodium vulgare	Wild Basil	CRPA 1900	vr	7 to 11	5,9,10.	Only other records EDM 1901, GCD "Alderney" 1932, AT 1978
629	118/20.1	Origanum vulgare	Wild Marjoram	AS 1966	vr	7 to 9	7,11	Refound BB 1987. not seen since 88
630	118/21.3	Thymus polytrichus	Wild Thyme	CCB 1838	c	5 to 11		
631	118/23.2	Mentha aquatica	Water Mint	CCB 1838	c	8 to 11	1,4,5,7,8,12,13.	Wet places. Fertile
632	118/23.2 x 3	<i>Mentha x piperita</i>	Peppermint	EDM 1901	o	7 to 10	9,10.	plant often red-tinged. Sterile
633	118/23.3	<i>Mentha spicata</i>	Spear Mint	EDM 1901	?	6 to 11	8,9.	Frequent in gardens. Fertile. Last wild record 1980
634	118/3 x lon	Mentha x villosonervata	Sharp-toothed Mint	OJ 1980	vr	8 to 9	8,	along railway line. Possibly a garden escape. Sterile. Det. RMH
635	118/23.3 x 4	<i>Mentha x villosa</i>	Apple Mint	DMcC1963		6 to 9	9,	Stranger's Cemetery. Sterile
636	118/23.4 x lon	<i>Mentha x rotundifolia</i>	Round-leaved Mint	GCD 1907				Marked 'S' in Comital Flora Possibly the only record
637	118/23.?	<i>Mentha longifolia</i>	Horse Mint	GCD1907	vr	7 to 9	8,	behind Museum. Marked 'S'. Refound BB 1988 Determined DMcC

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 36

638	118/25.3	<i>Salvia pratensis</i>	Meadow Clary	JO 1953			10,	'Nunnery' Bracts green/violet-blue, shorter than flowers Only record
639	118/25.4	Salvia verbenaca	Wild Clary	CCB 1838	lf	5 to 10	1,3,4,5,7,9.	Long calyx hairs white. Leaves lobed

Hippuridaceæ

640	119/1.1	Hippuris vulgaris	Mare's-tail	EDM 1901	vlf	5 to 10	4,5,7.	Mannez quarry only now. Earlier records, Longis pond Bonne Terre
-----	---------	-----------------------------------	-------------	----------	-----	---------	--------	--

Callitrichaceæ

641	120/1.3	Callitriche stagnalis	Common Water-starwort	GCD 1906			4,	Longis pond Only record
		(Photo taken in Guernsey)	Various-leaved Water-starwort	CCB 1838				7, 'Mill pond'. Only other record EDM 1901
642	120/1.4	<i>Callitriche platycarpa</i>						

Plantaginaceæ

		<i>Buck's-horn Plantain</i>	CCB 1838	a	4 to 10			Very variable in size
643	121/1.1	Plantago coronopus	Sea Plantain	AS 1955		7		Also recorded on subsequent visits, 58, 60, 65, but doubtful records
644	121/1.2	Plantago maritima	Greater Plantain	CCB 1838	f	4 to 11		Seen in small numbers in most squares
645	121/1.3	Plantago major	Ribwort Plantain	CCB 1838	a	3 to 11		Fasciated forms found occasionally
646	121/1.5	Plantago lanceolata						

Buddlejaceæ

648	122/1.2	Buddleja davidii	Butterfly-bush	NVM 1967	o	4 to 10	3,4,5,8,9,10.	Scattered self-sown garden escape. Also on old walls
649	122/1.2 x 3	Buddleja x weyeriana	Hybrid between 648 and 650	BB 1987	vr	7 to 10		8,10, Newtown, Bottom Essex valley. Probably planted
650	122/1.3	Buddleja globosa	Orange-ball-tree	BB 1987	r	5 to 8	10.	Probably planted, bottom Essex Hill

Oleaceæ

651	123/2.1	Fraxinus excelsior	Ash	CCB 1838	vlf	4 to 6	7,8,9,10.	Self-seeds readily
652		Fraxinus angustifolia	Ash	CH 1987	1 site	4 to 5	7.	With Brown buds, planted. Still surviving 1999 Only record

679	124/17.1	Hebe salicifolia	Koromiko	DMcC 1963	vr	3 to 12	Probably planted. Only record outside gardens
680	124/17 ell x spec	Hebe x franciscana	Hedge Veronica	DMcC 1963	f	1 to 12	2,3,5,7,8,13. Much used as hedging. Self seeds outside gardens
681	124/20	Euphrasia officinalis agg.	Eyebrights	CCB 1838	lc	4 to 10	1,2,4,5,6,7,9,10. Most of the Eyebrights have not been differentiated recently
682	124/20.1 x 3	Euphrasia rostkoviana	An Eyebright	CCB 1838)		Only other record EDM 1901
683	124/20.6	Euphrasia tetraqueta	An Eyebright	EDM 1900)		4,10. Euphrasia aggregate common in short turf all round the island
684	124/20.7	Euphrasia nemorosa	An Eyebright	EDM 1900	lc	4 to 10	4. Right to the sea edge.
685	124/20.9	Euphrasia confusa	An Eyebright	DMcC 1963)		Determined by PFY
686	124/20.10	Euphrasia stricta	An Eyebright	EDM 1901) lc		Appears to be the only record
687	124/20.14	Euphrasia ostenfeldii	An Eyebright	GCD 1906)		(4, As Euphrasia curta var. glabrescens) Only record
688	124/21.2	Odontites vernus	Red Bartsia	CCB 1838	vr	6 to 8	4,8. Two single patches in verges for many years until mowed off 1993. Recovered 1996 but not seen 2000-2002
689	124/23/1	Parentucellia viscosa	Yellow Bartsia	CCB 1838	lf	6 to 9	1,2,3,5,6,8,14 Quarries & E. coast. Damp valleys
690	124/25.2	Pedicularis sylvatica	Lousewort	EDM 1901	o	4 to 7	11,12,13. Moist places above cliffs. Baxter's Valley, abundant in Vallée des Gaudulons

Orobanchaceae

691	125/2.1	Orobanche purpurea	Yarrow Broomrape	CCB 1838	c	4 to 7	Even in lawns.
692	125/2.2	Orobanche rapum-genistae	Greater Broomrape (Scarce)	CCB 1838	vlc	5 to 7	11,12,13. S. cliffs on Cytisus & Ulex
693	125/2.8	Orobanche hederæ	Ivy Broomrape (Scarce)	EDM 1901	o	6 to 7	8. Confined to Connaught Sq & Le Vallée
695	125/2.10	Orobanche minor	Common Broomrape	CCB 1838	c	4 to 8	Many hosts.
696		Orobanche minor var. maritima	Carrot Broomrape (RDBk)	EDM 1901	vlf	4 to 8	3,7. On Daucus, Plantago, Glaucium & Ononis repens

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 39

Acanthaceae

697	127/1.1	Acanthus mollis	Bear's-breech	BB 1998	r	7 to 9	7, 11. One plant by track on Giffoine 1998, 1 on Platte Saline 2002
-----	---------	---------------------------------	---------------	---------	---	--------	---

Campanulaceae

698	129/1.9	Campanula portenschlagiana	Adria Bellflower	BB 1987	l	1 to 12	8. On old walls. Corolla funnel-shaped
699	129/1.10	Campanula poscharskyana	Trailing Bellflower	BB 1988	vlf	1 to 11	8. Old walls. Corolla star-shaped
700	129/6.1	Jasione montana	Sheep's-bit	CCB 1838	c	4 to 12	Always near sea

Rubiaceae

701	130/3.1	Sherardia arvensis	Field Madder	CCB 1838	a	3 to 10	Dry banks, cliffs, commons. White or pale pink/mauve flowers
702		Asperula cynanchica	Squinancywort	EDM 1901		3,5.	Only other record JCNW 1956
703	130/6.5	Galium palustre	Common Marsh-bedstraw	CCB 1838	o	5 to 7	4,6,7,12,13. Perennial. Rough. Scrambling, pondsides/ditches. Flowers white
704	130/6.6	Galium verum	Lady's Bedstraw	CCB 1838	la	6 to 9	Perennial. Short turf. One old wall in town. Flowers yellow

705 130/6.7a	Galium mollugo subsp. mollugo	Hedge Bedstraw	CCB 1838	r	5 to 8	
5,7,10,11,12,13,14. Perennial. Smooth, branches at >45°. Flowers white						
706 130/6.7b	Galium mollugo subsp. erectum	Hedge Bedstraw	ABJ 1932	vr	6 to 8	5,7,13.
On more calcareous soils. Branches <45°BB 1995						
707 130/6.10	Galium saxatile	Heath Bedstraw	EDM 1901	o	6 to 7	5,6,9,11,12,14. Acid soils.
Perennial. Leaves prickles on margins						
708 130/6.11	Galium aparine	Cleavers	CCB 1838	c	4 to 8	Annual. strong prickles on stems. fruits with hooked bristles. Flowers white
709 130/7.1	Cruciata laevipes	Crosswort	DMcC 1962			Only record
710 130/8.1	Rubia peregrina	Wild Madder	CCB 1838	vr	5 to 8	4,5,12,13. Evergreen. Prickles recurved. Flowers pale yellow-green. Only 1 plant seen since 1986, BB, by Vau Renier stream on cliff path

Caprifoliaceæ

711 131/1.2	Sambucus nigra	Elder	CCB 1838	f	4 to 10	Scattered all over
712 131/6.7	Lonicera periclymenum	Honeysuckle	CCB 1838	a	5 to 12	Colour varies from almost white to dark pink
713	Lonicera periclym var. quercifolia	Honeysuckle	BB 1988	r	6 to 9	7. Leaves oak-leaf shaped

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period

Page 40

Valerianaceæ

714 133/1.1	Valerianella locusta	Common Cornsalad	EDM 1901	vr	3 to 7	7,12 Calyx absent from fruit. Fr. just longer than wide
715 133/1.2	Valerianella carinata	Keeled-fruited Cornsalad	CCB 1838	c	3 to 11	1,2,3,5,8. Walls & garden weed. Fruit much longer than wide
716 133/1.4	Valerianella dentata	Narrow-fruited Cornsalad	EDM 1900	vr	6	8,9. Calyx distinct in fruit 2 teeth Refound DD 1995
717 133/1.5	Valerianella eriocarpa	Hairy-fruited Cornsalad (RDBk)	EDM 1901	vr		3,9,10. Calyx 6 teeth. Refound RR 1991, previous record JO 1953
718 133/3.1	Centranthus ruber	Red Valerian	EDM 1901	c	1 to 12	Principally on old walls & quarry spoil. Red, white or pink forms

Dipsaceæ

720 134/1.1	Dipsacus fullonum	Wild Teasel	CMR 1951	vr	7 to 8	2,7,8,13. Bracts on receptacle have stiff flexible, straight apical spine
721 134/1.2	Dipsacus sativus	Fuller's Teasel	DMcC 1958		7 to 8	Apical spine rigid & recurved. Only record
722 134/3.1	Knautia arvensis	Field Scabious	EDM 1901	r	7 to 9	3,9,10. E. side Les Rochers FLOWERS 7.57 Voucher

Asteraceæ Formerly Compositæ

723 135/2.1	Carlina vulgaris	Carlina Thistle	CCB 1838	o	6 to 10	3,4,5
723a	Osteospermum ? jocundrum	Cape Daisy	BB 18.6.11	vr	1st record	5 to 7 4. 5yds inland from longis sea wall 4 plants
727 135/3.1	Arctium lappa	Greater Burdock	CCB 1838			Leaf stalks solid. Possibly an error for above Only record
725 135/3.2c	Arctium minus subsp. minus	Lesser Burdock	EDM 1901	o	6 to 10	Scattered individuals in most squares. Lower leaf stalks hollow
726 135/3.2b	Arctium minus subsp. nemorosum		EDM 1901			2. As above, but peduncles much shorter, 0-1cm. Last record 1954
728 135/5.1	Carduus tenuiflorus	Slender Thistle	CCB 1838	c	4 to 8	1,2,3,4,5,7,11
729 135/5.4	Carduus nutans	Musk Thistle	CCB 1838	lf	6 to 10	2,4,5,7,9,10
731 135/6.2	Cirsium vulgare	Spear Thistle	CCB 1838	c	5 to 10	All Cirsium spp. are on Mauvaises Herbes list

732 135/6.8	Cirsium acaule	Dwarf Thistle	EDM 1901	vlf	7 to 10	3,4,5. Usually stemless, occasionally 8
-10cm. Rarely with white flowers						
733 135/6.9	Cirsium palustre	Marsh Thistle	EDM 1901	o	5 to 11	3,4,5,7,9,11,12. Wet places, usually in small numbers
734 135/6.10	Cirsium arvense	Creeping Thistle	CCB 1838	c	5 to 11	Also common garden weed
735 135/7.1	Onopordum acanthium	Cotton Thistle	BB 1991	vr	6 to 8	3,8,9
735a 139/8.1 (Stace)	Cynara cardunculus var. scolymus	Globe Artichoke (<u>close up</u>)	PHAS 1970	r		
7 to 10 7,9. Mainly relic of cultivated crops in 1950's. Now spreading						
736 135/11.1	Centaurea scabiosa	Greater Knapweed	CCB 1838	o	6 to 11	3,4,8,9,12,13,14

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 41

737 135/11.3	Centaurea cyanus	Cornflower (Scarce)	CCB 1838	vr		8,12 Only other records EDM 1901, CMR 1955, FLOWERS 30.5.97 (Voucher)
738 135/11.5	Centaurea aspera	Rough Star-thistle	EDM 1901	r	5 to 8	3, since 1890's in same spot. (Claimed Platte Saline FF 1967)
739 135/11.7 x jac	Centaurea x moncktonii	A hybrid Knapweed	ABJ 1932			3,4,5,11,12,14. CMR 1950 determined by WBT 1958.
741 135/11.7 x ?	Centaurea nemoralis x jacea	A hybrid Knapweed	DMcC 1958	vr		Determined by WBT 1958 Only other record BB 1992
742 135/11.7	Centaurea nigra var. nigra	Common Knapweed	EDM 1901	f	6 to 10	3,4,5,12,14
743 135/11.7	Centaurea nigra var. nemoralis	Common Knapweed	EDM 1901	f	6 to 10	11
744 135/11. ?	Centaurea cineraria	(A western Mediterranean sp.)	MK 1995		1 site	6 to 10 8. Base of Water Tower. Spreading 1997-99 Determined by ACL 11.9.95
745 135/12.1	Cichorium intybus	Chicory	EDM 1901	TD1.9.11	vr	6 to 10 4,5,7, 8,9,11
746 135/14.1	Lapsana communis	Nipplewort	EDM 1901	vr	6 to 7	8,9
747 135/15.1	Hypochaeris radicata	Cat's-ear	CCB 1838	c	4 to 1	
748 135/15.2	Hypochaeris glabra	Smooth Cat's-ear (Scarce)	CCB 1838	lf	4 to 2	2,3,4,5,7. Sandy soils mostly near coast
749 135/16.1	Leontodon autumnalis	Autumn Hawkbit	CCB 1838	c	1 to 12	Outermost ligules red on underside
750 135/16.2	Leontodon hispidus	Rough Hawkbit	CCB 1838	vr	6 to 8	9. Stace says "Not in CI". Refound by BSBI 1995. Previous 1900
751 135/16.3	Leontodon saxatilis	Lesser Hawkbit	CCB 1838	c	1 to 12	All round coast. Outermost ligules grey-violet underside
752 135/17.1	Picris echioides	Bristly Oxtongue	EDM 1901	c	5 to 1	Common on Sandy turf & verges
753 135/17.2	Picris hieracioides	Hawkweed Oxtongue	EDM 1901	f	5 to 12	
754	Picris hieracioides var. incana	Druce's Oxtongue	GCD 1906	r	7 to 9	2.4 Refound BB 1988. Locally in same area as Druce, obviously different jizz but not recognised as a separate var.
755 135/19.1b	Tragopogon pratensis subsp. minor	Goat's-beard	EDM 1901	r	3 to 11	3,7,8,9. Yellow flowers
756 135/19.2	Tragopogon porrifolius	Salsify	EDM 1901			2. Purple flowers. Last records 1907, 1954
757 135/21.2	Sonchus arvensis	Perennial Sow-thistle	CCB 1838	r	1, 6-11	4,9,11,12
758 135/21.3	Sonchus oleraceus	Smooth Sow-thistle	CCB 1838	c	1 to 12	
759 135/21.4	Sonchus asper	Prickly Sow-thistle	EDM 1901	f	3 to 9	
760 135/22.1	Lactuca serriola	Prickly Lettuce	ML1985	r	7 to 10	7,8.. Walls & gutters,
761 135/25	Taraxacum spp. agg.	Dandelions	CCB 1838	c	1 to 12	Have not been fully determined, but at least 19 subsp. recorded
	Sect. 1. Erythrosperma	Lesser Dandelions	EDM 1901			

785 135/27.1	Pilosella peleteriana	Shaggy Mouse-ear-hawkweed	CCB 1838		o	5 to 9	8. Short grass, dunes
Pilosella & Hieracium spp. need more investigation							
786 135/27.2	Pilosella officinarum	Mouse-ear-hawkweed	EDM 1901		f	5 to 10	Ligules red-striped beneath
Determined by CW							
787 135/28.6	Hieracium umbellatum	Umbellate Hawkweed	EDM 1901		o	6 to 10	4,8,10,13.
Leaves all sessile							
788 135/30.1	Filago vulgaris	Common Cudweed	CCB 1838		r	7 to 9	3,4,5,8,10,13. Stems to 40cm. fl clusters 10-12mm
789 135/30.4	Filago minima	Small Cudweed	CCB 1838		o	6 to 9	5,10,13. Stems to 20cm. fl clusters 2-5mm
790 135/33.5	Gnaphalium uliginosum	Marsh Cudweed	EDM 1900		r	7 to 11	1,4,5,7,8.
Flowering clusters leafy. Phyllaries pale brown with dk. tips							
791 135/33.6	Gnaphalium luteo-album	Jersey Cudweed	ML 1974		vr	7 to 1	2,7,8. Leaves grey, woolly. Phyllaries straw-coloured
792 135/33.7	Gnaphalium undulatum	Cape Cudweed	BHR 1956		lf	6 to 10	3,4,5,7,8,9,11,13.
To 80cm. Leaves green on underside. Phyllaries whitish							
793 135/34	Helichrysum petiolare	Everlasting Flower	BB 1993		r	6 to 8	10. Well established
Bluestone Hill, seeding in verge. Originally a garden escape							
794 135/35.3	Inula conyzae	Ploughman's-spikenard	EDM 1903		r	6 to 9	4,14. Fl heads 1cm, flowers yellowish, outer bracts purple. Scrub etc.
795 135/35.4	Inula crithmoides	Golden-samphire (Scarce)	CCB 1838		o	6 to 9	2,3,4,6,10,11,12. Fl heads 2.5cm, golden-yellow. Near or on old forts
796 135/37.1	Pulicaria dysenterica	Common Fleabane	CCB 1838		f	4 to 10	Wet or damp places
797 135/40.3	Solidago canadensis	Canadian Goldenrod	RR 1963		1 site	7 to 10	6. Probably planted many years ago
798 135/41.6	Aster tripolium	Sea Aster	WF 1956		r	8 to 10	5. 2nd record BB 1.9.97, c 30 plants nr. Houmet Herbé
799 135/43.1	Erigeron glaucus	Seaside Daisy	BB 1987		o	1 to 12	2,4,7,8,10. Walls & banks. Flowers single 3-4cm, ray florets lilac
800 135/43.4	Erigeron karvinskianus	Mexican Fleabane	ML 1970		vr	3 to 11	7, 8 Old walls. flowers in loose cluster, 1.5cm, ray florets white, purple below
801 135/43.6	Erigeron acer	Blue Fleabane	CCB 1838		vr	5 to 7	8,9,10. Annual/biennial. Fl heads 12-18mm, ray florets pale purple

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 44

802 135/44.1	Conyza canadensis	Canadian Fleabane	CMR 1950		lf	1 to 12	2,3,4,5,6,7,8,9,10. Bare sandy places
802a 135/44.2	Conyza sumatrensis	Guernsey Fleabane	ML 2000		o	6 to 10	7,8. Platte Saline, Braye
Common & gardens. Not previously separated from C. canadensis							
802b 135/44.3	Conyza bilbaoana	Hybrid Fleabane	BB 10.7.08		lf	1st. record	6 to 10 7,8,9. Becoming frequent
803 135/45.1	Olearia paniculata	Akirabo	BB 1987		vr	8 to 10	5,8. Planted hedging. Occasional seedlings, inconspicuous flowers
804 135/45.2	Olearia macrodonta	New Zealand Holly	BB 1987		o	6 to 10	8,9. Planted. Churchyard and gardens. Showy white flowers.
805 135/45.3	Olearia traversii	Ake-ake	BB 1987		o	6 to 8	5,8. Hedging. Plant on E. coast to 7m. Flowers greyish axillary pyramid
806 135/47.1	Bellis perennis	Daisy	CCB 1838		a	1 to 12	
807 135/48.1	Tanacetum parthenium	Feverfew	DMcC 1956		r	6 to 9	5,8. Garden escape. Churchyard & Museum
808 135/48.3	Tanacetum vulgare	Tansy	ABJ 1932		vlf	4 to 9	2,3,5,7,8. First record, 'near Mannez Cottages'

810 135/50.1	Artemisia vulgaris	Mugwort	EDM 1901	o	6 to 10	3,4,6-10,12,14. Waste places. Leaves green above, reddish, angled stem
811 135/50.3	Artemisia absinthium	Wormwood	CCB 1838	o	5 to 11	1,2,3,4,5,7,10. Often with 718. Leaves silvery both sides
812 135/52.1	Otanthus maritimus	Cottonweed (RDBk)	CCB 1838			3 Only other record 1896, gone by 1899
813 135/53.1	Achillea ptarmica	Sneezewort	VKM 1956	vr	6 to 12	2,7,9. Garden escape
814 135/53.3	Achillea millefolium	Yarrow	CCB 1838	a	3 to 12	
815 135/54.1	Chamaemelum nobile	Chamomile (Scarce)	CCB 1838	lf	5 to 8	6,11,12. (Cliffs)
816 135/55.2	Anthemis arvensis	Corn Chamomile	CCB 1838			Only other record. FF 1967 Trois Vaux
817 135/55.3	Anthemis cotula	Stinking Chamomile	EDM 1899			10 Only confirmed record
818 135/56.1	Chrysanthemum segetum	Corn Marigold	CCB 1838	vr	5 to 9	11,12. Formerly abundant in cultivated fields. Now rare
819 135/58.1	Leucanthemum vulgare	Oxeye Daisy	CCB 1838	la	2 to 11	Cliffs
820 135/58 lac x max	Leucanthemum x superbum	Shasta Daisy	DMcC 1961	vr	1 to 12	Frequent in gardens, rare escape
821 135/59.1	Matricaria recutita	Scented Mayweed	CMR 1956	vr	3 to 11	Very strongly sweet-scented. Phyllaries pale brown
822 135/59.2	Matricaria discoidea	Pineapple-weed	CMR 1954	c	4 to 1	Bare places & sandy tracks. Pineapple scent. Phyllaries, white margins
823 135/60.1	Tripleurospermum maritimum	Sea Mayweed	CMR 1954	c	3 to 12	Stems prostrate/decumbent. Strand line & above all round coast
824 135/60.2	Tripleurospermum inodorum	Scentless Mayweed	CCB 1838	c	1 to 12	Cultivated & waste ground. Leaf segments acute with bristle tips
825 135/62.1	Senecio cineraria	Silver Ragwort	JP 1977	r	6 to 9	1,2,3,4. Braye Beach. Probably a garden escape originally
826 35/62.10	Senecio jacobea	Ragwort	CCB 1838	a	4 to 1	On Mauvaises Herbes list

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 45

827 135/62.15	Senecio vulgaris	Groundsel	CCB 1838	f	1 to 12	
828 135/62.17	Senecio sylvaticus	Heath Groundsel	CCB 1838	lc	4 to 11	Mainly on cliffs & in old quarries. Erect, larger than above
828a 135/62.3	Senecio inequidens	Narrow-leaved Ragwort		o	5 to 7	11 Hampshire botanists group
829 135/70.1	Tussilago farfara	Colt's-foot	JDG 1937		5,14	Only other records AS 1958, DMcC/FLOWERS 1963
830 135/71.4	Petasites fragrans	Winter Heliotrope	EDM 1901	la	12 to 4	3,7,8
831 135/73.1	Calendula officinalis	Pot Marigold	DMcC 1957	f	1 to 12	2,3,7,8. Frequent garden escape
832 135/79.1	Helianthus annuus	Sunflower	BB 1988	r	8 to 10	7,9,12. Garden escape or remnants of planted seed crop
832a	Helianthus x laetifolius	Perennial Sunflower	BB 2000	r	8 to 10	7, 9. Verge in Picaterre by stream c 20 plants. Present for several years
833 135/79.2	Helianthus tuberosus	Jerusalem Artichoke	BB 1989	vr	8 to 10	7, Hedge in Le Petit Val
834 135/86.1	Eupatorium cannabinum	Hemp Agrimony	JP 1977	vr	6 to 9	4 Near Saye Farm. Only record in the wild

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

854 151/1.14	Juncus acutiflorus	Sharp-flowered Rush	EDM 1901	f	7 to 9	4. The latest flowering rush
855 151/1.15	Juncus bulbosus	Bulbous Rush	EDM 1899	vr	5 to 7	6,7. Amphibious Only other records 1934, 1987
856 151/1.20	Juncus maritimus	Sea Rush	CCB 1838	vla	7 to 8	4,5,6,10. Abundant in Berry's Quarry. Lowest bract exceeds inflorescence.
857 151/1.21	Juncus acutus	Sharp Rush (Scarce)	CCB 1838	o	5 to 9	5. (round Mannez pond). Stem very rigid
858 151/1.25	Juncus inflexus	Hard Rush	CCB 1838	lc	6 to 8	4,5,12. Abundant in Mannez quarry. Stem ridged, interrupted pith
859 151/1.26	Juncus effusus	Soft Rush	EDM 1901	f	6 to 10	4,5,7,11,13. Esp, round Mannez pond. Stem smooth, glossy
860 151/1.27	Juncus conglomeratus	Compact Rush	CCB 1838	vr	6 to 8	7 (Bonne Terre), 11 (Trois Vaux). Stems ridged, dull. Last record DMcC 1963
861 151/2.5	Luzula campestris	Field Woodrush	CCB 1838	a	3 to 6	Flowers in 1 sessile & 2 stalked clusters 3-12 in each
862 151/2.6	Luzula multiflora	Heath Woodrush	CCB 1848	o	4 to 7	4. Flowers in several stalked clusters, 8-18 in each

Cyperaceae

The Alderney Sedges need further investigation

863 152/3.1	Eleocharis palustris	Common Spike-rush	CCB 1838	vla	5 to 6, 10	5,10,11,13. Upper stem leaf-sheath truncated
864 152/3.4	<i>Eleocharis multicaulis</i>	Many-stalked Spike-rush	CCB 1838	r	6 to 7	5,7,11. Upper stem leaf-sheath oblique at apex

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

865 152/3.5	<i>Eleocharis quinqueflora</i>	Few-flowered Spike-rush	DMcC 1963	vr	6 to 7	As distinct from above 2 spp, lowest glume completely encircles base
866 152/4.1	Bolboschoenus maritimus	Sea Club-rush	EDM 1901	r	5 to 7	Burhou only. Refound FWS 1958, BB 1989-99
867 152/7.2	<i>Schoenoplectus tabernaemontani</i>	Grey Club-rush	EDM 1901	?		4. Possibly the only record
868 152/8.1	Isolepis setacea	Bristle Club-rush	EDM 1901	vlf	5 to 8	5,7,11,12. Bract much longer than inflorescence
869 152/8.2	Isolepis cernua	Slender Club-rush	CCB 1838	r	6 to 8	6,7,11. Bract barely longer than inflorescence
870 152/11.1	Cyperus longus	Galingale (Scarce)	CCB 1838	BB 27.1.08	r	New site 6 to 10 1,3,6,7 Streambed Fontaine David
870a 152/	Cyperus involucrata (C. alternifolius)	Umbrella Sedge	BB 7/1997	vr	6 to 10	7. Several clumps round both Platte Saline Ponds and seedlings in the grass. Garden escape.
871 152/12.1	Schoenus nigricans	Black Bog-rush	CCB 1838	vlf	5 to 6	6,11. (Next species also found in Clonque Valley by JDG 1937)
872 152/16.1	Carex paniculata	Greater Tussock-sedge	EDM 1901	r	5 to 8	7. Old Mill leat in Bonne Terre only. 21 large plants. <u>Photo</u> of site in January 2000 after clearance of willow scrub.
873 152/16.8b	<i>Carex muricata</i> subsp. <i>lamprocarpa</i>	Prickly Sedge	CCB 1838	r	6	5,6,13. Female glumes shorter & darker than utricles
874 152/16.9a	Carex divulsa subsp. <i>divulsa</i>	Grey Sedge	EDM 1901	r	6 to 7	7,8. Verge Route de Picaterre. Ripe spikes greenish-brown
875 152/16.9b	Carex divulsa subsp. <i>leersii</i>	Tall Spiked-sedge	ABJ 1933	r	6 to 7	7,8,14. Requires calcareous soil. Ripe spikes reddish-brown
876 152/16.10	Carex arenaria	Sand Sedge	CCB 1838	la	3 to 9	Sandy shores, dunes, etc
877 152/16.11	<i>Carex disticha</i>	Brown Sedge	ML 1996	1 site	6 to 8	5. Mannez Quarry near well head. Det. CJ. from BB specimen

878	152/16.16	Carex ovalis	Oval Sedge	ABJ 1932					'Mannez Quarry, one tuft' Only record
879	152/16.17	Carex echinata	Star Sedge	EDM 1901					Clanque cliffs' Only record
880	152/16.21	Carex lachenalii	Hare's-foot Sedge	ABJ 1932					(Mannez Quarry as C. leporina) Doubtful record because of habitat
881	152/16.23	Carex hirta	Hairy Sedge	EDM 1901	lc	4 to 8			Cliff valleys. Stem 3-sided. Lowest bract with pubescent sheath
882	152/16.27	Carex pseudocyperus	Cyperus Sedge	BB 1992			1 site	5 to 8	4. Longis pond
883	152/16.31	Carex pendula	Pendulous Sedge	DMcC 1963	vlf		4 to 8	7,8,	Line of Le Vallée
									stream. Seeds vigorously. Also planted near garden ponds
884	152/16.35	Carex flacca	Glaucous Sedge	(photo with leaves)	CCB 1838		f	4 to 8	2,3,4,5,9,11. Stems in loose glaucous tufts. Wet or dry grassland
885	152/16.36	Carex panicea	Carnation Sedge	CCB 1838			o	5 to 6	12,13,14. Cliff valleys. Prefers wet acid soils
886	152/16.41	Carex distans	Distant Sedge	ABJ 1933			r	5 to 7	3,4,13. Wet rocky places near sea. (With HKAS)
887	152/16.43	Carex extensa	Long-bracted Sedge	EDM 1901			r	6 to 7	5,13. Lowest bract much longer than inflorescence bent at rt angle
888	152/16.45	Carex flava	Large Yellow-sedge	CCB 1838					EDM 1901 'C in Cliff valleys'. Possibly the only records
889	152/16.46	Carex viridula	Yellow-sedge	JO 1953					3. Only record
890	152/16.51	Carex caryophylla	Spring Sedge	CCB 1838	vr	4 to 5			10. Habitat short grassland
891	152/16.73	Carex pulicaris	Flea Sedge	CCB 1838	vr	5 to 6			11

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
 Flowering period Squares where found and notes

Page 49

Poaceae	Formerly Graminae	Alderney's Grasses need more investigation. There are probably species not yet recorded							
Subfamily Bambusoideae									
892	153/ 1.1	Yushania anceps	Indian Fountain-bamboo	AHL 1968	vr				Planted. Small self-sown clump in quarry scree, Corblets, BB 8/97. Increased 2002
893	153/3.1	Pseudosasa japonica	Arrow Bamboo	AHL 1968					Planted
Subfamily Pooideae									
894	153/8.1	Nardus stricta	Mat-grass	AS 1960					Mannez Quarry. Doubtful record
895	153/12.1	Festuca pratensis	Meadow Fescue	CCB 1838					
896	153/12.2	Festuca arundinacea	Tall Fescue	EDM 1901	r	5 to 8			Quite plentiful in 8,9,11.
897	153/12.6	Festuca arenaria	Rush-leaved Fescue	ABJ 1933					5. Specimen in BM Only record
898	153/12.7a	Festuca rubra subsp. rubra	Red Fescue	CCB 1838	c	5 to 8	3,4,6,7,8,	10.	
899	153/12.7b	Festuca rubra subsp. juncea		EDM 1901					Possibly the only record
900	153/12.8	Festuca ovina	Sheep's-fescue	CCB 1838	c	5 to 9			The many subspecies have not been separated
901	153/12.10	Festuca filiformis	Fine-leaved Sheep's-fescue	DMcC 1973	o	5 to 6	1,7		
902	153/12.12	Festuca huonii	Houn's Fescue	BB 1987	r				4. Rocky scree Saye Bay
903	153/12.13	Festuca lemanii	Confused Fescue	CCB 1838					Possibly the only record
904	153/12.14	Festuca longifolia	Blue Fescue	CCB 1838					Possibly the only record
905	153/12.7 x 1	X Festulpia hubbardii	A hybrid Fescue	ABJ 1933					Other Festuca hybrids may exist
906	153/13.1	Lolium perenne	Perennial Rye-grass	CCB 1838	c	5 to 12			Spikelets with 4-14 florets
907	153/13.1 x 2	Lolium x boucheanum	A hybrid Rye-grass	PL 1976			r	5 to 9	7. Refound BB 1987
908	153/13.2	Lolium multiflorum	Italian Rye-grass	EDM 1901	vlf	5 to 9			Spikelets with 11-22 florets
909	153/?	Lolium temulentum	Darnel	CCB 1838					'St. Anne' Only record

[910](#) [153/14.1](#) [Vulpia fasciculata](#) Dune Fescue (Scarce) CCB 1838 f 5 to 7 3,4,5,7,8,10.
Spikelets 10-18mm. GCD 1907 as Vulpia membranacea

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 50

[911](#) [153/14.2](#) [Vulpia bromoides](#) Squirreltail Fescue CCB 1838 c 4 to 7 3,7,8. Spikelets 6-11mm
[912](#) [153/14.3](#) [Vulpia myuros](#) Rat's-tail Fescue CCB 1838 vr 4 to 7 9,10 Refound above Impôt

BB, 8.6.89
[913](#) [153/14.4b](#) [Vulpia ciliata](#) subsp. [ambigua](#) Bearded Fescue GCD 1906 Braye &

Corblets **Only record**

[914](#) [153/15.1](#) [Cynosurus cristatus](#) Crested Dog's-tail CCB 1838 c 4 to 8 All ex 1,8.
[915](#) [153/15.2](#) [Cynosurus echinatus](#) Rough Dog's-tail BB 1995 vr 6 to 7 4,5
[916](#) [153/17.1](#) [Briza media](#) Quaking-grass EDM 1901 r 6 to 7 3,6,9
[917](#) [153/17.2](#) [Briza minor](#) Lesser Quaking-grass BB 1986 vr 7 9
[918](#) [153/17.3](#) [Briza maxima](#) Greater Quaking-grass WFS 1986 r 1 to 12 2,7
[919](#) [153/18.1](#) [Poa infirma](#) Early Meadow-grass (RDBk) MMcW 1962 vlf 3 to 5 Culms to 10cm

(and on Burhou)

[920](#) [153/18.2](#) [Poa annua](#) Annual Meadow-grass CCB 1838 c 1 to 12 Culms to 20cm
[921](#) [Poa annua](#) var. [reptans](#) MH2 1978 **Only record, Burhou**
[922](#) [153/18.3](#) [Poa trivialis](#) Rough Meadow-grass CCB 1838 f 3 to 7 6,7. Culms to 70cm
[923](#) [153/18.4](#) [Poa humilis](#) Spreading Meadow-grass CCB 1838 f 5 to 7 7,9. Culms to 30cm
[924](#) [153/18.5](#) [Poa pratensis](#) Smooth Meadow-grass CCB 1838 c 5 to 7 9. Culms to 75cm
[925](#) [153/18.6](#) [Poa angustifolia](#) Narrow-leaved Meadow-grass TR 1990 1 site 5 to 7 8. Culms to 70cm

[926](#) [153/18.7](#) [Poa chaixii](#) Broad-leaved Meadow-grass EDM 1901 **10 Only record**

[928](#) [153/19.1](#) [Dactylis glomerata](#) Cock's-foot CCB 1838 a 1 to 12
[929](#) [153/20.1](#) [Catabrosa aquatica](#) Whorl-grass JES 1970 vr 6 to 8 Wet, sandy places near sea. **Possibly**

the only record

[930](#) [153/21.1](#) [Catapodium rigidum](#) Fern-grass EDM 1901 o 4 to 7 3,4,5,7. Walls and bare ground. Prefers alkaline soil. Very rigid

[931](#) [153/21.2](#) [Catapodium marinum](#) Sea Fern-grass CCB 1838 c 4 to 9 1,2,4,7. Very shiny.
Later flowering than above. Often reddish

[932](#) [153/23.1](#) [Parapholis strigosa](#) Hard-grass JES 1969 **Only record**

[933](#) [153/23.2](#) [Parapholis incurva](#) Curved Hard-grass HJMB 1957 **Only record**

[934](#) [153/24.2](#) [Glyceria fluitans](#) Floating Sweet-grass CCB 1838 vlf 5 to 8 7,12,13

[935](#) [153/24.2 x 4](#) [Glyceria x pedicellata](#) Hybrid Sweet-grass JO 1953 o 6 to 7 5

[936](#) [153/24.3](#) [Glyceria declinata](#) Small Sweet-grass JES 1969 vr 6 to 7 1,2

[937](#) [153/24.4](#) [Glyceria notata](#) Plicate Sweet-grass EDM 1901 o 5 to 7 7,9,10,13

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 51

[938](#) [153/26.1](#) [Helictotrichon pubescens](#) Downy Oat-grass EDM 1901 o 6 to 7 4,9,12

[939](#) [153/26.2](#) [Helictotrichon pratense](#) Meadow Oat-grass GCD 1906 Braye' **Only record**

[940](#) [153/27.1](#) [Arrhenatherum elatius](#) False Oat-grass CCB 1838 c 5 to 11 1,2,3,6,7,8,9,10,11

[941](#) [153/27.1b](#) [Arrhenatherum elatius](#) subsp. [bulbosum](#) False Oat-grass EDM 1901 c 5 to 11
1,2,7. Bulbils at base of stems

[942](#) [153/28.2](#) [Avena strigosa](#) Bristle Oat EDM 1901 r 6 to 7 9,12 Refound BB 1987

[943](#) [153/28.3](#) [Avena fatua](#) Wild-oat CCB 1838 r 6 to 9 4,10,12,13

[944](#) [153/28.4a](#) [Avena sterilis](#) subsp. [ludoviciana](#) Winter Wild-oat DMcC 1963

Determined by AM. **Only record**

945	153/28.5	Avena sativa	Oat	DMcC 1958	o	5 to 8	4,12,13.	Relic of cultivation
945a		Avena barbata	Slender Oat	BB/JG 30.7.10	vlf	5 to 7	13 NE Field before	Madonna stone
946	153/29.1	Gaudinia fragilis	French Oat-grass	ABJ 1933	r	5 to 7	4,8.	Whitegates. (With HKAS) Refound Roadside near Dairy DMcC 1963 & JEL 1971
947	153/30.1	Trisetum flavescens	Yellow Oat-grass	CCB 1838	o	6 to 8	4,8,9	
948	153/31.2	Koeleria macrantha	Crested Hair-grass	EDM 1901	f	6 to 7	4,5	
949	153/33.1	Holcus lanatus	Yorkshire-fog	CCB 1838	c	4 to 8		
950	153/33.2	Holcus mollis	Creeping Soft-grass	EDM 1901	r	7 to 9	7	
951	153/35.1	Aira caryophylla	Silver Hair-grass	CCB 1838	c	4 to 7	Stems to 25cm	
952	153/35.2	Aira praecox	Early Hair-grass	CCB 1838	c	4 to 5	Stems to 10cm	
953	153/37.1	Anthoxanthum odoratum	Sweet Vernal-grass	CCB 1838	c	4 to 9		
954	153/38.2	Phalaris aquatica	Bulbous Canary-grass	JP 1977				10 Only record
955	153/38.3	Phalaris canariensis	Canary-grass	WH 1881 BB 26.9.10	r	3-5 branched plants	Platte Saline	6 to 9 7,9,10. Probably a casual from bird seed. Panicle ovoid
956	153/38.4	Phalaris minor	Lesser Canary-grass	CRPA 1889	lf	6 to 11	1,7,8,12,13,14.	Panicle longer & more slender
957	153/39.1	Agrostis capillaris	Common Bent-grass	CCB 1838	c	5 to 9		
958	153/39.2	Agrostis gigantea	Black Bent	ABJ 1932	?	6 to 8	Refound FH 1977	
959	153/39.4	Agrostis stolonifera	Creeping Bent	CCB 1838	c	3 to 11		
960	153/39.7	Agrostis canina	Velvet Bent	EDM 1901	r	5 to 8	7	
961	153/41.1	Ammophila arenaria	Marram	CCB 1838	la	6 to 9	3,4,5,6.	Dunes & commons.
962	153/43.1	Lagurus ovatus	Hare's-tail	RM 1951	r	5 to 9	2,4,5,7,9	Refound JES 1966, BB 1990

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 52

963	153/44.1	Apera spica-venti	Loose Silky-bent (Scarce)	JES 1966				Only record ?
964	153/46.2	Polypogon viridis	Water Bent	GCD 1906	vr	6 to 7	3,8.	AS 1956 det. FLOWERS No site. Refound in a gutter in town MK 1995. Longis Garden centre, MBU 1999
965	153/47.1	Alopecurus pratensis	Meadow Foxtail	EDM 1901	r	4 to 6	8,12	
966	153/47.2	Alopecurus geniculatus	Marsh Foxtail	EDM 1901	r	4 to 8	11,12	Refound ABJ 1932
967	153/49.1	Phleum pratense	Timothy	EDM 1901	r	6 to 8	8,10.	Culms erect to 1.5m
968	153/49.2	Phleum bertolinii	Smaller Cat's-tail	GM 1969	vr			Culms to 50cm Cliff path BB 1989, not conf. Possibly the only records
969	153/49.5	Phleum arenarium	Sand Cat's-tail	CCB 1838	lc	5 to 7	1,3,4,5.	Sandy coast and commons. Rarely above 10-15cm
970	153/50.2	Bromus commutatus	Meadow Brome	GCD 1906				Only record ?
971	153/50.4a	Bromus hordeaceus subsp. hordeaceus	Soft Brome	CCB 1838	c	4 to 9		Culms to 80cm
972	153/50.4b	Bromus hordeaceus subsp. ferronii	Least Soft-brome	ABJ 1933	f	5 to 7	1,4,7,	Culms 10-15cm
973	153/50.4c	Bromus hordeaceus subsp. thominei		EDM 1901	lf	5 to 7	4, 5, 9, 10-14,	Culms 5-8cm (Marquand as B. nanus)
974	153/50.4 x 5	Bromus x pseudothominei	Lesser Soft-brome	GCD 1906	o	5 to 8	8,9,10,11.	Culms erect, to 80cm Refound FLOWERS 1984
974a	153/51.3	Bromopsis erecta	Upright Brome	EMcD 1995	vr	5 to 6	5.	Track just W of Hole-in-the-Wall. 6 plants First record
975	153/52.1	Anisantha diandra	Great Brome	EDM 1901	a	5 to 8	2,7,8,9,12,13	
976	153/52.2	Anisantha rigida	Ripgut Brome	EDM 1901	lc	5 to 8	Braye	
977	153/52.3	Anisantha sterilis	Barren Brome	CCB 1838	a	4 to 10		
978	153/52.5	Anisantha madritensis	Compact Brome	DMcC 1956	r	5 to 7		

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
 Flowering period Squares where found and notes

Page 54

Sparganiaceae

[1,005](#) [154/1.1a](#) [Sparganium erectum](#) Branched Bur-reed EDM 1901 vr 6 to 8 4,7 Refound FF 1967,
ML1996 in Longis pond

Typhaceae

1,006 [155/1.1](#) [Typha latifolia](#) Bulrush DMcC 1961 1 site 7 to 9 10. Tank on Essex Hill, probably
planted originally
1,007 [155/1.2](#) [Typha angustifolia](#) Lesser Bulrush BB 1987 1 site 6 to 9 5. Mannez pond. Male &
female flowers separated by stem

Liliaceae **Now includes former Amaryllidaceae**

[1,008](#) [158/6.2](#) [Kniphofia praecox](#) Greater Red-hot-poker BB 1987 r 1 to 12 5,7,10. Scattered
garden escape. Determined J. Taylor

1,009 [Kniphofia tuckii](#) A Red-hot-poker BB 1989 vr 1 to 12 5,9 Determined J. Taylor

1,010 [158/10.3](#) [Tulipa gesneriana](#) Garden Tulip DB 1994 4 2, Garden escape or throwout **Only
record**

1,011 [158/13.1](#) [Convallaria majalis](#) Lily-of-the-valley DMcC 1963 vr 2 to 7 9. Garden escape. Refound
BB 1988

[1,012](#) [158/18.2](#) [Ornithogallum angustifolium](#) Star-of-Bethlehem CCB 1838 r 2 to 6 4,5,8,9

1,013 [158/19.6](#) [Scilla peruviana](#) Portuguese Squill TWA 1958 1 site 6 to 7 4. Single plant persisting
many years since about 15 dug up c. 1980

[1,014](#) [158/19.7](#) [Scilla autumnalis](#) Autumn Squill CCB 1838 la 7 to 10 E. & S. Coasts, Corblets
Quarry & Mannez Garenne

[1,015](#) [158/20.2](#) [Hyacinthoides non-scripta](#) Bluebell CCB 1838 c 3 to 6 Less robust than H.
hispanica. Tip of raceme bent over. Also with white flowers

[1,016](#) [158/20.2 x 3](#) [Hyacinthoides non-scripta x H. hispanica](#) A hybrid Bluebell DMcC 1973 f 3 to 5
5, 10, 11. Probably frequent, characters intermediate difficult to determine.. Cross fertile

[1,017](#) [158/20.3](#) [Hyacinthoides hispanica](#) Spanish Bluebell CMR 1950 la 3 to 5/6 Tip of raceme
straight. Occasionally with white flowers

1,018 [158/23.2](#) [Muscari armeniacum](#) Garden Grape-hyacinth BB 1997 vr 3 to 4 9. Scrub at car
parking area W. of Impôt. 7-10 plants well established

1,019 [158/23.4](#) [Muscari comosum](#) Tassel Hyacinth CH /RW 1976 1 site 5 to 6 8. above quarry below Butes.
Still there 1997

1,020 [158/24.1](#) [Allium schoenoprasum](#) Chives NVM 1967 4 to 8 Garden escape **Only record**

1,021 [158/24.3](#) [Allium roseum subsp. bulbiferum](#) Rosy Garlic NVM 1982 vr 4 to 6 7. Two plants
found on Platte Saline, JB 5/1998

[1,022](#) [158/24.4](#) [Allium neapolitanum](#) Neapolitan Garlic BB1988 BB2010 vr Several clumps at new sites
along E. coast 5 to 7 8,9. 1962-82 records error for following sp. Spathe 1 large, broad Sq. 5

1,022a [158/24.6](#) [Allium moly](#) Yellow Garlic BB 4.6.10 vr 1st. record
6 to 7 7. Platte Saline 3x2m patch on recently disturbed soil

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
 Flowering period Squares where found and notes

Page 55

[1,023](#) [158/24.5](#) [Allium subhirsutum](#) Hairy Garlic FLI 1962 r 3 to 6 4,5,6,7,8,9,10. Spathe 3, small

[1,024](#) [158/24.7](#) [Allium triquetrum](#) Three-cornered Garlic EDM 1901 a 9 to 7 Single patch in
1900. On Mauvais Herbes list since 1952

1,025 [158/24.15](#) [Allium spherocephalon](#) Round-headed Leek BB 1987 vr 6 to 8 7. Inflorescence has
flowers only, no bulbils

[1,026](#) [158/24.16](#) [Allium vineale](#) Wild Onion DMcC 1967 r 5 to 8 3,4,5. Inflorescence has flowers &
bulbils or bulbils only

1,026a	158/27.1	Agapanthus praecox	African Lily	BB 13 7 09	vr	2 sites widely separated	1st records	6 to 8
7,11. Tourgis hill behind Dolmen and on Giffoine								
1,027	158/28.1	Tristagma uniflorum	Spring Starflower	BB 1988		2 sites	2 to 4	4,10. Garden escape. Fort Essex & Route des Mielles in asphalt road
1,028	158/29.1	Amaryllis belladonna	Jersey Lily	JP 1985	1 site	7 to 11	8, Roadside bank near school	
1,029	158/31.1	Leucojum aestivum	Summer Snowflake	MMcW 1962	2 sites	2 to 5	8,9 Found in The Terrace	BB 1989
1,030	158/33	Narcissus spp. ag.	Not indigenous. Mostly relics of cultivation in the 1948-60's.				Several	
unidentified spp. are quite Common								
1,031	158/33.1 x 3	Narcissus x medioluteus	Primrose-peerless	KF 1957		o	3 to 5	
1,032	158/33.3	Narcissus poeticus	Pheasant's-eye Daffodil	CMR 1954		o	11 to 5	4,5,6,7,10,12
1,033	158/33.5	Narcissus pseudonarcissus	Daffodil. (The English native species)		CMR 1954		1 site	
3 to 5 4. Field beside Fort Corblets. Probably planted 1948-50								
1,033.1	158/33.7	Narcissus minor	Spanish Daffodil. (Another wild species, SW Europe)		BB 1998			
1 site 3 to 5 4. Field beside Fort Corblets. Probably planted 1948-50								
1,034	158/33.?	Narcissus papyraceus	Paper White (Tazetta group)		BHR 1957		f	1 to 5
Widespread throughout island. Relic of 1950's cultivation								
1,034.1	158/33.?	Narcissus 'primo'	'Grande primo citronière' (Another Tazetta group sp.)		BB 1998			
a 2 to 4 Widespread around island. Relic of 1950's cultivation. By far the commonest multiheaded sp. Field margins, hedges, verges, bramble scrub, waste ground. Literally thousands everywhere, March 2000								
1,035	158/34.1	Asparagus officinalis subsp. officinalis	Garden Asparagus	DMcC 1957	vr		5 to 7	
9. Garden escape								
1,036	158/35.1	Ruscus aculeatus	Butcher's Broom	CCB 1838		o	12 to 4	5,9,10,12,13,14. Cliffs
1,037	158/	Polygonatum multiflorum	Solomon's Seal	BB 1993		vr	4 to 6	7, In scrub at St. Vignalis' Gdn. Probably planted, still there 1999
1,038	158/37.?	Crinum powelli		BB 1994	1 site	5 to 10	3. Large clump in dunes behind Saye Bay, W.	end for c. 20 yrs.

Iridaceae

1,039	159/5 fil x tin	Iris x hollandica	Dutch Iris	BB 1990	vr	3 to 4	7, Roadside hedge	Watermill Farm. ?Relic of cultivation. Still there 2002
-------	-----------------	-----------------------------------	------------	---------	----	--------	-------------------	---

List sequence No., with Kent number in red Scientific name English name Finder/date Frequency
Flowering period Squares where found and notes

Page 56

1,040	159/5.1	Iris germanica	Bearded Iris	BB 1989	vr	7 to 8	5. Verge near Lighthouse for many years. Gone 1998
1,041	159/5.3	Iris pseudacorus	Yellow Iris	CCB 1838	v/c	4 to 7	1,4,7,10,12. Now only round Longis & Platte Saline Ponds
1,042	159/5.8	Iris foetidissima	Stinking Iris	EDM 1901	f	6 to 7	Abundant in Sq.4. F. in Vallée and on Giffoine. O. in other damp spots
1,043	159/7.1	Romulea columnae	Sand Crocus (RDBk)	CCB 1838		v/a	3 to 5 Cliffs, rim of Berry's quarry and E. coast
1,044	159/9.2b	Gladiolus communis	Eastern Gladiolus	CMR 1954		o	4 to 7 Scattered round coasts
1,043.1	159/10.1	Ixia paniculata	Tubular corn-lily	BB 2000	vr	5 to 6	10. 3 plants on roadside bank, Bluestone Hill, 23.5.00
1,045	159/13.3 x aur	Crocsmia x crocosmiiflora	Montbretia	CMR 1950		f	7 to 9 Scattered, cliffs & quarries

Agavaceae

1,046	160/2.1	Agave americana	Century Plant	In flower in Alderney 2011	BB 1986	vr	?	
5, 7. Planted many years ago, has suckered								
1,047	160/3.1	Cordylone australis	Cabbage Palm	BB 1986		lf	5 to 8 3,7,8,10. Planted in the 1930's, and in many gardens. Self-seeds	

1,048 [160/4.1](#) [Phormium tenax](#) New Zealand Flax BB 1987 r 6 to 8 7,8. Garden escape. Field on cliffs sq, 7 ?planted

1,049 [160/](#) [Trachycarpus fortunei](#) Chusan Palm RW 1987 vr 5 to 6 8,13. Planted in several gardens

Orchidaceæ

[1,050](#) [162/7.1](#) [Spiranthes spiralis](#) Autumn Lady's-tresses EDM 1901 c 7 to 10 Scattered in short turf on dunes and cliffs

[1,051](#) [162/14.1](#) [Anacamptis pyramidalis](#) Pyramidal Orchid EDM 1901 lc 5 to 10 1,2,3,4,5,7,8,9,10. Short sandy turf

[1,052](#) [162/18.1](#) [Dactylorhiza fuchsii](#) Common Spotted-orchid EDM 1901 vr 6 to 8 7. Leaves with dark elongated spots

[1,053](#) [162/18.2](#) [Dactylorhiza maculata](#) Heath Spotted Orchid CMR 1954 vr 6 to 8 3, 10. single plant BB Essex Hill 1989. Leaves with dark round spots

[1,055](#) [162/18.4](#) [Dactylorhiza praetermissa](#) Southern Marsh-orchid EDM 1900 r 5 to 8 7 (BonneTerre), 11. A few each St. Vignalis bog & old Mill leat. No spots

[1,056](#) [162/20.3](#) [Orchis morio](#) Green-winged Orchid EDM 1902 vlf 4 to 6 4, 5, 12. Mainly Mannez Garenne & E. coast

1057.1 [162/20.2](#) [Orchis mascula](#) Early-purple Orchid JM 1954 BB 27.3.2008 vr vr Several plants by field gate next to Mannez House 2 to 4 Longid, lsrge patch near TheKennels VKM 1957, "Absent" DMcC 1975

[1,058](#) [162/23.3](#) [Ophrys apifera](#) Bee Orchid SH 1949 JB2 14.6.08 BB 2 plants there 29.6.11 2 sites New site, grassy bank above harbour 6 to 7 4. Now a single colony, 1-7 plants. Not seen 1983-1991 Refound 1992 & up to 25 plants most years to 2011

This ends the main plant list.

Total 1090 species 17.2.2012